

University of Massachusetts Amherst

From the Selected Works of Raymond S Bradley

2011

High-resolution paleoclimatology

Raymond S Bradley, *University of Massachusetts - Amherst*



Available at: https://works.bepress.com/raymond_bradley/83/

Chapter 1

High-Resolution Paleoclimatology

Raymond S. Bradley

Abstract High resolution paleoclimatology involves studies of natural archives as proxies for past climate variations at a temporal scale that is comparable to that of instrumental data. In practice, this generally means annually resolved records, from tree rings, ice cores, banded corals, laminated speleothems and varved sediments. New analytical techniques offer many unexplored avenues of research in high resolution paleoclimatology. However, critical issues involving accuracy of the chronology, reproducibility of the record, frequency response to forcing and other factors, and calibration of the proxies remain. Studies of proxies at high resolution provide opportunities to examine the frequency and magnitude of extreme events over time, and their relationships to forcing, and such studies may be of particular relevance to societal concerns.

Keywords Climate dynamics · Natural archives · Paleoclimate · Proxies

1.1 Introduction

Paleoclimatology uses natural archives to reconstruct climate in the pre-instrumental period. The longest instrumental records are from Western Europe, and a few of these extend back into the early eighteenth (or even late seventeenth) century. However, for most regions, continuous instrumental measurements rarely extend beyond the early nineteenth century, with some remote (desert or polar) regions having barely 50 years of observations (Fig. 1.1). Consequently, our instrumental perspective on climate variability is extremely limited. In particular, it is unlikely

R.S. Bradley (✉)

Department of Geosciences, Climate System Research Center, University of Massachusetts, Amherst, MA 01003-9297, USA
e-mail: rbradley@geo.umass.edu




Fig. 1.1 Approximate earliest date of continuous instrumental records, which defines the need for high-resolution proxy-based data prior to these dates

that we understand the full spectrum of variability of the most important climate modes (such as the El Niño/Southern Oscillation [ENSO], Pacific Decadal Oscillation [PDO], North Atlantic Oscillation [NAO]. etc). High-resolution paleoclimatology addresses this issue by focusing on climate proxies that can be resolved at seasonal to annual resolutions. These proxy records may extend back continuously from the present, or provide discrete windows into the past, to shed light on modes of variability in earlier times. By providing data at a resolution comparable to that of the instrumental record, high-resolution paleoclimatology plays an important role in resolving anthropogenic effects on climate. Specifically, it helps to place contemporary climate variability in a long-term perspective (*detection*, in the parlance of the Intergovernmental Panel on Climate Change [IPCC]), and it enables climatic changes to be examined in terms of forcing mechanisms (*attribution*). High-resolution paleoclimatology also provides targets (either time series or maps of past climatic conditions) with which models (general circulation models [GCMs] or energy balance models [EBMs]) can be tested and validated, and it offers the opportunity to explore climate dynamics (modes of variability, abrupt climate changes, climate system feedbacks) over long periods of time. Thus, high-resolution paleoclimatology naturally interfaces with, and complements, the research priorities of the climate dynamics community.

This figure will be printed in b/w

46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90

1 High-Resolution Paleoclimatology

1.2 Data Sources for High-Resolution Paleoclimatology

The critical requirements for high-resolution paleoclimatology are that:

- An accurate chronology can be established; this generally requires replication of the archive being sampled.
- The archive can be sampled in detail, ideally at seasonal to annual resolutions, but at least at the resolution of a few years.
- The parameter being measured is reasonably well understood in terms of its relationship to climate (i.e., its mechanistic and seasonal response) so that it can be calibrated in terms of climate, by using the instrumental record as a yardstick for interpreting the paleorecord.
- The relationship between the proxy and climate observed today has been similar in the past (the principle of uniformitarianism).
- The record captures variance of climate over a wide range of frequencies, or at least the window of variance that the proxy does capture is known.

In the next section, these issues are examined with reference to the main archives that are available for high-resolution paleoclimatology: tree rings, corals, speleothems, ice cores, and varved sediments. This examination is followed by a discussion of the opportunities and challenges in high-resolution paleoclimatology, with particular reference to dendroclimatology.

1.3 Chronology and Replication

An accurate timescale is essential in high-resolution paleoclimatology. A chronology is commonly obtained by counting annual increments, by using variations in some parameter to mark the passage of time. This might be the cyclical ^{18}O maximum in a coral record, registering the sea surface temperature (SST) minimum over each annual cycle; or the presence of a 'clay cap' in varved lacustrine sediments, marking each winter's sediment layer; or the width of a tree ring between the large, open-walled spring cells that form each year. However, simply counting these recurrent features in a sample (even if they are counted several times by different analysts) does not guarantee an accurate chronology. The best procedure is to replicate the record by using more than one sample (core), to eliminate potential uncertainties due to 'missing' layers and to avoid misinterpretation of dubious sections. On this matter, dendroclimatic studies have a clear and unambiguous advantage over most other paleoclimate proxies. Duplicate cores are easily recovered, and cross-dating using one or more samples is routinely done. Tree-ring chronologies are thus as good as a natural chronometer can be, at least for those regions where there is an annual cycle of temperature or rainfall and trees are selected to record such changes in their growth. However, for those vast areas of equatorial and tropical forests, where trees are not under climatic stress and so do not produce annual rings,

136 establishing a chronology has been far more challenging. Recent analytical
137 improvements using continuous flow isotope mass spectrometry have made feasi-
138 ble the almost continuous sampling of wood, so that annual changes in isotopic
139 properties can be identified, even in wood that appears to be undifferentiated in
140 its growth structure (Evans and Schrag 2004; Poussart et al. 2004). This technique
141 opens up the possibility of using trees for paleoclimatic reconstruction in regions
142 that were hitherto unavailable. However, replication of samples from nearby trees is
143 still necessary to reduce chronological uncertainties in these newer records.

144 In the case of most other high-resolution proxies, replication is rarely carried
145 out. This is generally related to the cost of sample recovery (in terms of logistics
146 or time) or because of the analytical expense of duplicating measurements. Most
147 coral records, for example, are based on single transects through one core, though
148 the veracity of the chronology may be reinforced through the measurement of mul-
149 tiple parameters, each of which helps confirm the identification of annual layering
150 in the coral. Similarly, in ice cores, multiparameter glaciochemical analyses can
151 be especially useful in determining a secure chronology (McConnell et al. 2002a;
152 Souney et al. 2002). In addition, in some locations more than one core may be
153 recovered to provide additional ice for analysis and to help resolve uncertainties in
154 chronology (Thompson 1993). It may also be possible to identify sulfate peaks in the
155 ice, related to explosive volcanic eruptions of known age. Such chronostratigraphic
156 horizons can be very helpful in confirming an annually counted chronology (Stenni
157 et al. 2002). Varved sediments are sometimes analyzed in multiple cores, but sample
158 preparation (such as impregnation of the sediments with epoxy, thin section prepa-
159 ration, etc.) is expensive and very time-consuming, so duplication is not commonly
160 done. Where radioactive isotopes from atmospheric nuclear tests conducted in the
161 late 1950s and 1960s can be identified in sediments (and in ice cores), such horizons
162 can be useful time markers. Tephra layers (even finely dispersed cryptotephra) can
163 be useful in confirming a sedimentary chronology if the tephra can be geochemi-
164 cally fingerprinted to a volcanic eruption of known age (e.g., Pilcher et al. 2005).
165 Finally, where annual layer counting is not feasible—as in many speleothems—
166 radioactive isotopes (^{210}Pb , ^{14}C , and uranium-series) can be used to obtain mean
167 deposition/accumulation rates, though there may have been variations in those rates
168 between dated levels.

169
170

171 1.4 High-Resolution Sampling

172

173 Advances in analytical techniques have now made sub-annual sampling and mea-
174 surements fairly routine in most high-resolution proxies. Whereas tree rings were
175 generally measured in terms of total annual increments, densitometry now enables
176 measurements of wood density and incremental growth in early and latewood sec-
177 tions of each annual ring. Image analysis provides further options in terms of
178 analyzing cell growth parameters (Panyushkina et al. 2003). Isotopic dendroclimatic
179 studies require subannual sampling resolution to determine growth increments. In
180 corals, such detailed sampling is now routine; often 10 or more samples will be

1 High-Resolution Paleoclimatology

181 obtained per annual increment (e.g., Mitsuguchi et al. 1996; Quinn and Sampson
182 2002). Stalagmite research has rarely achieved such detail, with sampling intervals
183 (in most studies) of a few years at best. However, some studies have established
184 chronologies by counting annual layers on polished sections under a microscope,
185 and new analytical approaches (using an electron microprobe, secondary ionization
186 mass spectrometry [SIMS], or excimer laser ablation–inductively coupled plasma–
187 mass spectrometry [ELA-ICP-MS]) have made it feasible to identify annual layers
188 through seasonal changes in trace elements (such as Mg, Ca, Sr, Ba, and U), along
189 multiple transects of a sample (e.g., Fairchild et al. 2001; Desmarchelier et al. 2006).
190 Image analysis of varved sediments (via impregnated thin sections examined under
191 a petrographic or scanning electron microscope) can reveal intra-annual sediment
192 variations that may be associated with seasonal diatom blooms or rainfall events
193 (Dean et al. 1999). In ice cores, it is now possible to make continuous multipa-
194 rameter measurements, providing extremely detailed time series (McConnell et al.
195 2002a, b). Thus, in most natural archives available for high-resolution paleoclima-
196 tology, detailed measurements can be made both to define annual layers or growth
197 increments and to characterize changes therein. However, it is not necessarily the
198 case that an annual layer fully represents conditions over the course of a year. Much
199 of the sediment in a varve, for example, may result from brief periods of runoff.
200 Similarly, annual layers in an ice core represent only those days when snowfall
201 occurred. Indeed, they may not even do that, if snow was subsequently lost through
202 sublimation or wind scour. Coral growth increments may result from more continu-
203 ous growth, and trees may also grow more continuously, at least during the growing
204 season. Speleothems accumulate from water that has percolated through the overly-
205 ing regolith, and so short-term variations related to individual rainfall episodes are
206 likely to be ‘smoothed out.’ Nevertheless, there is some evidence that extreme rain-
207 fall episodes can be detected in the carbon isotopes of speleothems in areas where
208 the throughflow of water is rapid (Frappier et al. 2007).

1.5 Relationships Between Natural Archives and Climate


213 Extracting a climatic signal from individual archives requires an understanding of
214 the climatic controls on them. Analysis of the temporal relationships between vari-
215 ables may provide a statistical basis for calibration, but a theoretical basis for such
216 a relationship is also required, to direct some light into the statistical black box.
217 This may require in situ process-based studies to understand the factors control-
218 ling the proxy signal. Even if such studies are short-term, they can provide valuable
219 insights into how climate influences the system being studied, and hence improve
220 our understanding of the paleoclimatic record. For example, studies of meteorolo-
221 gical conditions at the ice-coring site on Sajama, Bolivia, demonstrated strong
222 seasonality in snow accumulation, with much of the snowfall that accumulated late
223 in the accumulation season being subsequently lost through sublimation (Hardy
224 et al. 2003). Consequently, the ice core record is made up of sections of snow that
225 accumulated for (at most) a few months each year, demonstrating that division of

226 such records into 12 monthly increments is not appropriate (cf. Thompson et al.
227 2000a). Similarly, hydrological studies in the Arctic have shown that in some lakes,
228 much of the runoff and associated sediment may be transferred into the lake over the
229 course of only a few weeks. For example, measurements at Sophia Lake (Cornwallis
230 Island, Nunavut, Canada) showed that 80% of the runoff and 88% of the annual
231 sediment flux occurred in the first 33 days of the 1994 melt season (Braun et al.
232 2000). This sediment was subsequently distributed across the lake floor, forming an
233 annual increment (varve), but the climatic conditions that mobilized the sediment
234 were brief and perhaps unrepresentative of the summer season (and the year as a
235 whole). Other studies of arctic lakes indicate that watersheds containing glaciers
236 provide more continuous runoff and associated sediment flux throughout each summer,
237 and thus provide a better proxy for summer climatic conditions (e.g., Hardy
238 et al. 1996). Thus, understanding the environment from which the proxy archive is
239 extracted is critically important for proper interpretation of the paleoclimate record.
240 Process-based studies (often derided as simply ‘monitoring’) have also provided
241 insights into climatic controls on corals, showing strong nonlinearities at high water
242 temperatures (Lough 2004). In situ measurements within caves, aimed at gain-
243 ing a better understanding of paleoclimate records, are now also being carried out
244 (e.g., McDonald et al. 2004; Cruz et al. 2005). By comparison, dendroclimatology
245 is far advanced because ecophysiological studies of tree growth have a long history.
246 Consequently, factors influencing tree growth increments are well understood
247 (Fritts 1996; Schweingruber 1996; Vaganov et al. 2006), providing a very strong
248 foundation for paleoclimatic studies using tree rings.


251 1.6 Uniformitarianism

253 Perhaps because of the rapidity of recent climate change, many archives are no
254 longer responding to climate in a manner that typifies much of the past. This phe-
255 nomenon was first noted by Briffa et al. (1998), who showed that some trees that
256 were formerly strongly influenced by temperature were no longer so influenced,
257 or at least not to the same extent. Figure 1.2 shows the geographical distribution
258 of this effect. Briffa et al. (2004) speculated that this response might be related to
259 recent increases in ultraviolet radiation resulting from the loss of ozone at high ele-
260 vations. Others have argued it might reflect the fact that trees in some areas have
261 reached a threshold, perhaps now being affected more by drought stress than was
262 formerly the case. Whatever the reason, it raises the question of whether such con-
263 ditions might have occurred in the past, and if so, whether it would be possible
264 to recognize such a ‘decoupling’ of the proxy archive from the (‘normal’) climate
265 driver. Paleoclimate reconstruction is built on the principle of uniformitarianism, in
266 which the present is assumed to provide a key to the past. If modern conditions (dur-
267 ing the calibration period) are not typical of the long term, this assumption will be
268 invalid. It is thus important to resolve the reasons for such changes and determine if
269 additional parameters (such as cell growth features) might provide clues about when
270 such stresses may have overwhelmed the typical climate response.

1 High-Resolution Paleoclimatology



Regression with difference series (infilled)



This figure will be printed in b/w

AQI

Fig. 1.2 (a) Instrumental temperatures (*red, heavier line*) and tree-ring density reconstructions of temperature (*black, thinner line*) averaged over all land grid boxes north of 50°N, smoothed with a 5-year low-pass filter. (b) Map showing where the average temporal pattern of divergence between tree-ring density chronologies and mean warm season temperatures is most apparent. The smoothed difference between the *black* and *red* curves in (a) were regressed against the local difference curves produced from the averages of data in each grid box. Where the regression slope coefficients are progressively >1.0 (the *yellow, orange and red* boxes, which are generally the most northerly locations), the greater is the local difference between density and temperature. In the areas shown *blue and light purple* (areas further south), the difference is apparent but of lower magnitude. The areas shown as *dark purple* (basically the most southern regions) do not show the divergence [note change in scale on color bar] (from Briffa et al. 2004). On-line version shows these figures in color

316 On a related point, it is clear that many natural archives are being detrimentally
317 affected by recent changes in climate. Thus, many high-elevation ice caps in the
318 tropics have been affected by surface melting and strong sublimation, so that the
319 recent isotopic record has been degraded or even lost entirely (Thompson et al.
320 2000b). Similarly, corals in many areas were greatly affected by exceptionally high
321 sea surface temperatures associated with the 1997–1998 El Niño (Wilkinson et al.
322 1999). Many century-old *Porites* colonies in the Great Barrier Reef were killed at
323 this time.

324 325 326 327 **1.7 Frequency Response**

328
329 High-resolution records may have certain low-frequency characteristics that differ
330 from the spectrum of the climatic environment in which they are situated. Such
331 effects may be due to long-term biological growth (in the case of trees, and per-
332 haps corals), compaction (ice, sediments), non-climatic changes in depositional
333 environments (lake sediments, speleothems), and other proxy records. This issue
334 is especially important as efforts are made to extend paleoclimatic reconstructions
335 further back in time, to reveal changes in climate over thousands of years. Sediments
336 are certainly affected by compaction, but this effect can be relatively easily corrected
337 for by examining changes in density. This is also true in ice cores. Diffusion of iso-
338 topes within firn leads to a reduction in the amplitude of isotopic values that must
339 also be considered. Deposition rates in speleothems are determined by radiocar-
340 bon or uranium series dates, and such analysis is generally sufficient to determine
341 if deposition has been continuous over time. Certainly, there are no compression
342 issues to be concerned with here, so in that sense speleothems do offer a very good
343 option for identifying low-frequency changes in climate. This is illustrated well in
344 the Dongge Cave record of Wang et al. (2005) (Fig. 1.3). The record shows an under-
345 lying low-frequency decline in monsoon precipitation, related to orbital forcing, on
346 which decadal- to centennial-scale variations are superimposed, which appear to be
347 (at least in part) related to variations in solar irradiance.

348 The issue of determining low-frequency changes in climate has been most prob-
349 lematical in dendroclimatology. The biological growth function of trees must first
350 be removed before climatic information can be extracted. When this procedure is
351 done, some low-frequency information may be lost. Furthermore, since most tree-
352 ring series are short, assembling a composite long time series from many short
353 records makes it even more problematical to obtain low-frequency information
354 over timescales longer than the typical segment length (Cook et al. 1995). New
355 approaches to standardization of tree-ring series have been developed, and these
356 help to preserve more low-frequency information than do more traditional methods.
357 However, such approaches require very large datasets and so cannot be applied in all
358 cases. Another approach involves combining different proxies, some that may con-
359 tain more low-frequency information with others that capture well higher-frequency
360 information, so that together they cover the full spectrum of climate variability

1 High-Resolution Paleoclimatology




Fig. 1.3 (a) $\delta^{18}\text{O}$ time series of a Dongge Cave (China) stalgmite (*thin line*). Six vertical shaded bars denote the timing of Bond events 0–5 in the North Atlantic. Two *vertical gray bars* (without numbers) indicate two other notable weak Asian monsoon periods that can be correlated to ice-rafted debris events. Higher frequency variability appears to be related to solar (irradiance) forcing. NCC is the Neolithic Culture of China, which collapsed at the time indicated. (b) Age-depth relationship. *Black error bars* show ^{230}Th dates with 2σ errors. Two different age-depth curves are shown, one employing linear interpolation between dated depths and the second slightly modified by tuning to INTCAL98 within the ^{230}Th dating error (from Wang et al. 2005). On-line version shows this figure in color

This figure will be printed in b/w

(Moberg et al. 2005). This approach has much promise, and further fine-tuning will likely lead to a better understanding of large-scale climate variability over recent millennia.

1.8 High-Resolution Proxies: Challenges and Opportunities

High-resolution paleoclimatic records provide unique opportunities to better understand the climate system because they extend the limited sampling interval that is available from short instrumental records. This longer perspective is especially important for studies of rare events, such as explosive volcanic eruptions or the occurrence of extreme climatic conditions such as droughts or floods. Ice cores reveal (through sulfate and electrical conductivity measurements) that there have been much larger explosive volcanic eruptions in the past than during the period of instrumental records (Zielinski et al. 1994; Castellano et al. 2005); by identifying these events, it is then possible to explore the relationship between eruption size and location and the subsequent climatic effects (e.g., D'Arrigo and Jacoby 1999). Many dendroclimatic studies have recognized the connection between explosive eruptions and cold growing season conditions, which sometimes have led to frost damage in trees (e.g., LaMarche and Hirschboeck 1984; Baillie and Munro 1988; Briffa et al. 1990; D'Arrigo et al. 2001). Proxy records of volcanic forcing also provide a much larger database of eruption events than is available for the instrumental

406 period; compositing climatic conditions following such events increases the signal-
407 to-noise ratio, giving a clearer view of the climate system response to such events.
408 Thus Fischer et al. (2007) were able to show that summer conditions in Europe have
409 tended to be both cold and dry after major tropical volcanic eruptions; but in winter,
410 a positive NAO circulation has generally been established, resulting in mild, wet
411 conditions in northern Europe and well below average precipitation in the Alps and
412 Mediterranean region.

413 Dendroclimatic research has been especially important in documenting the fre-
414 quency, geographical extent, and severity of past drought episodes, as well as
415 periods of unusually high rainfall amounts; such studies have been especially exten-
416 sive in the United States (e.g., Stahle and Cleaveland 1992; Hughes and Funkhouser
417 1998; Cook et al. 2004). These studies have shown that there has often been a
418 strong connection between severe droughts in the southwestern United States and
419 the occurrence of La Niña episodes, although the precise geographical pattern of
420 each drought has varied over time (Stahle et al. 2000; Cole et al. 2002). Tree-
421 ring research has also been applied to reconstructing modes of circulation in the
422 past, such as the North Atlantic Oscillation (Cook et al. 1998; Cullen et al. 2001),
423 Pacific Decadal Oscillation (Gedalof and Smith 2001), and Atlantic Multidecadal
424 Oscillation (AMO) (Gray et al. 2004). In all of these cases, the paleoclimatic recon-
425 structions have expanded our understanding of the spectrum of variability of these
426 modes of circulation and provided insight into how large-scale teleconnections (and
427 interactions between Atlantic- and Pacific-based circulation regimes) may lead to
428 persistent, large-amplitude anomalies over North America and other regions.

429 Great strides have been made in constructing hemispheric- and global-scale
430 patterns of past climate variability by combining many different types of high-
431 resolution paleoclimatic records, using a variety of statistical methods (Mann et al.
432 1998, 1999, 2005; Moberg et al. 2005; Rutherford et al. 2005). These studies have
433 demonstrated the importance of volcanic and solar forcing, and of the increasingly
434 dominant effects of anthropogenic forcing over the last 150 years. Nevertheless,
435 such studies rely largely on the most extensive database of paleoclimatic recon-
436 structions that is currently available—that provided by dendroclimatology. On the
437 one hand, this is good because the physiological basis for how trees respond to cli-
438 mate is well understood, thanks to decades of careful studies, and tree rings provide
439 the most accurate chronologies available. However, the use of tree rings in long-
440 term paleoclimate reconstructions is dogged by questions of uniformitarianism (a
441 question not unique to dendroclimatology, of course), but more significantly by the
442 difficulty of resolving the full spectrum of climate variability from overlapping, rela-
443 tively short, tree-ring series. This matter can be resolved by obtaining longer records
444 where possible, expanding the tree-ring database to improve data density back in
445 time, and developing new statistical approaches; all these methods are necessary
446 to ensure that long-term paleoclimatic reconstructions are as reliable as possible.
447 New isotopic and image analysis techniques applied to tree growth may add further
448 information about past climate variations in regions that were formerly off-limits to
449 dendroclimatologists, thereby extending the geographical domain for large-scale cli-
450 mate reconstruction. New proxies, especially from lake sediments and speleothems,

1 High-Resolution Paleoclimatology

will likely further supplement this expansion of high-resolution records, providing records with more robust low-frequency characteristics that can be combined with proxies that are exceptionally good at capturing high-frequency climate variability (e.g., Moberg et al. 2005). In this way, the next decade of high-resolution paleoclimatology will likely see paleoclimatic reconstructions with far less uncertainty, covering more geographical regions, and providing meaningful estimates of climate sensitivity before the ‘Anthropocene’.

Acknowledgements I gratefully acknowledge the support of my research by the National Oceanic and Atmospheric Administration (NOAA, NA050AR4311106), the National Science Foundation (NSF, ATM-0402421), and the U.S. Department of Energy (DOE, DE-FG02-98ER62604).

References

- Baillie MGL, Munro MAR (1988) Irish tree-rings, Santorini and volcanic dust veils. *Nature* 332:344–346
- Braun C, Hardy DR, Bradley RS, Retelle M (2000) Streamflow and suspended sediment transport into Lake Sophia, Cornwallis Island, Nunavut, Canada. *Arctic Antarctic Alpine Res* 32: 456–465
- Briffa KR, Bartholin TS, Eckstein D, Jones PD, Karlen W, Schweingruber FH, Zetterberg P (1990) A 1400 year tree-ring record of summer temperatures in Fennoscandia. *Nature* 346:434–439
- Briffa KR, Osborn TJ, Schweingruber FH (2004) Large-scale temperature inferences from tree rings: a review. *Global Planet Change* 40:11–26
- Briffa KR, Schweingruber FH, Jones PD, Osborn TJ, Shiyatov SG, Vaganov EA (1998) Reduced sensitivity of recent tree-growth to temperatures at high northern latitudes. *Nature* 391:678–682
- Castellano E, Becagli S, Hansson M, Hutterli M, Petit JR, Rampino MR, Severi M, Steffensen JP, Traversi R, Udisti R (2005) Holocene volcanic history as recorded in the sulfate stratigraphy of the European Project for Ice Coring in Antarctica Dome C (EDC96) ice core. *J Geophys Res* 110:D06114. doi:10.1029/2004JD005259
- Cole JE, Overpeck JT, Cook ER (2002) Multiyear La Niña events and persistent drought in the contiguous United States. *Geophys Res Lett* 29:1647. doi:10.1029/2001GL013561
- Cook ER, Briffa KR, Meko DM, Graybill DS, Funkhouser G (1995) The ‘segment length curse’ in long tree-ring chronology development for paleoclimatic studies. *Holocene* 5:229–237
- Cook ER, D’Arrigo RD, Briffa KR (1998) The North Atlantic Oscillation and its expression in circum-Atlantic tree ring chronologies from North America and Europe. *Holocene* 8:9–17
- Cook ER, Meko DM, Stahle, DW, Cleaveland MK (1999) Drought reconstructions for the continental United States. *J Climate* 12:1145–1162
- Cook ER, Woodhouse CA, Eakin CM, Meko DM, Stahle DW (2004) Long-term aridity changes in the western United States. *Science* 306:1015–1018
- Cruz FW Jr, Karmann I, Viana O Jr, Burns SJ, Ferrari JA, Vuille M, Moreira MZ, Sai NF (2005) Stable isotope study of cave percolation waters in subtropical Brazil: implications for paleoclimate inferences from speleothems. *Chem Geol* 220:245–262
- Cullen HM, D’Arrigo RD, Cook ER, Mann ME (2001) Multiproxy reconstructions of the North Atlantic Oscillation. *Paleoceanography* 16:27–39
- D’Arrigo RG, Jacoby JC (1999) Northern North American tree-ring evidence for regional temperature changes after major volcanic events. *Climatic Change* 41:1–15
- D’Arrigo R, Frank D, Jacoby G, Pederson N (2001) Spatial response to major volcanic events in or about A.D. 536, 934 and 1258: frost rings and other dendrochronological evidence from Mongolia and northern Siberia. *Climatic Change* 49:239–246
- Dean JM, Kemp AES, Bull D, Pike J, Patterson G, Zolitschka B (1999) Taking varves to bits: scanning electron microscopy in the study of laminated sediments and varves. *J Paleolimnology* 22:121–136

- 496 Desmarchelier J, Hellstrom JM, McCulloch MT (2006) Raid trace element analysis of speleothems
497 by ELA-ICP-MS. *Chem Geol* 231:102–117
- 498 Evans MN, Schrag DP (2004) A stable isotope-based approach to tropical dendroclimatology.
499 *Geochim Cosmochim Acta* 68:3295–3305
- 500 Fairchild IJ, Baker A, Borsato A, Frisia S, Hinton RW, McDermott F, Tooth AF (2001) Annual
501 to sub-annual resolution of multiple trace-element trends in speleothems. *J Geol Soc London*
502 158:831–841
- 503 Fischer, EM, Luterbacher J, Zorita E, Tett SFB, Casty C, Wanner H (2007) European climate
504 response to tropical volcanic eruptions over the last half millennium. *Geophys Res Lett* 34,
505 doi:10.1029/2006GL027992
- 506 Frappier A, Sahagian D, Carpenter SJ, González LA, Frappier BR (2007) Stalagmite stable isotope
507 record of recent tropical cyclone events. *Geology* 35:111–114
- 508 Fritts HC (1996) *Tree rings and climate*. Academic Press, San Diego
- 509 Gedalof Z, Smith D (2001) Inter-decadal climate variability and regime-scale shifts in Pacific North
510 America. *Geophys Res Lett* 28:1515–1518
- 511 Gray ST, Graumlich LJ, Betancourt JL, Pederson GT (2004) A tree ring-based reconstruction
512 of the Atlantic Multidecadal Oscillation since A.D. 1567. *Geophys Res Lett* 31,
513 doi:10.1029/2004GL019932
- 514 Hardy DR, Bradley RS, Zolitschka B (1996) The climatic signal in varved sediments from Lake
515 C-2, northern Ellesmere Island, Canada. *J Paleolimnology* 16:227–238
- 516 Hardy DR, Vuille M, Bradley RS (2003) Variability of snow accumulation and isotopic
517 composition on Nevado Sajama, Bolivia. *J Geophys Res-Atmospheres* 108: D22, 4693.
518 doi:10.1029/2003JD003623
- 519 Hughes MK, Funkhouser G (1998) Extremes of moisture availability reconstructed from tree rings
520 for recent millennia in the Great Basin of western North America. In: Innes M, Beniston JL
521 (eds) *The impacts of climate variability on forests*. Springer, Berlin, Heidelberg, New York, pp
522 99–107
- 523 LaMarche VC, Hirschboeck K (1984) Frost rings in trees as records of major volcanic eruptions.
524 *Nature* 307:121–126
- 525 Lough JM (2004) A strategy to improve the contribution of coral data to high-resolution
526 paleoclimatology. *Palaeogeog Palaeoclimatol Palaeoecol* 204:115–143
- 527 Mann ME, Bradley RS, Hughes MK (1998) Global-scale temperature patterns and climate forcing
528 over the past six centuries. *Nature* 378:266–270
- 529 Mann ME, Bradley RS, Hughes MK (1999) Northern Hemisphere temperatures during the past
530 millennium: inferences, uncertainties, and limitations. *Geophys Res Lett* 26:759–762
- 531 Mann ME, Rutherford S, Wahl E, Ammann C (2005) Testing the fidelity of methods used in proxy-
532 based reconstructions of past climate. *J Climate* 18:4097–4107
- 533 McConnell JR, Lamorey GW, Hutterli MA (2002a) A 250-year high-resolution record of Pb flux
534 and crustal enrichment in central Greenland. *Geophys Res Lett* 29:2130–2133
- 535 McConnell JR, Lamorey GW, Lambert SW, Taylor KC (2002b) Continuous ice-core chemical
536 analyses using inductively coupled plasma mass spectrometry. *Environ Sci Technol* 36:7–11
- 537 McDonald J, Drysdale R, Hill D (2004) The 2002–2003 El Niño recorded in Australian cave drip
538 waters: implications for reconstructing rainfall histories using stalagmites. *Geophys Res Lett*
539 31: L22202. doi:10.1029/2004GL020859
- 540 Mitsuguchi T, Matsumoto E, Abe O, Uchida T, Isdale PJ (1996) Mg/Ca thermometry in coral
541 skeletons. *Science* 274:961–963
- 542 Moberg A, Sonechkin DM, Holmgren K, Datsenko NM, Karlén W (2005) Highly variable
543 Northern Hemisphere temperatures reconstructed from low- and high-resolution proxy data.
544 *Nature* 433:613–617
- 545 Panyushkina IP, Hughes MK, Vaganov EA, Munro MAR (2003) Summer temperature in north-
546 eastern Siberia since 1642 reconstructed from tracheids dimensions and cell numbers of *Larix*
547 *cajanderi*. *Can J Forest Res* 33:1–10

1 High-Resolution Paleoclimatology

- 541 Pilcher J, Bradley RS, Francus P, Anderson L (2005) A Holocene tephra record from the Lofoten
542 Islands, Arctic Norway. *Boreas* 34:136–156
- 543 Poussart PF, Evans MN, Schrag DP (2004) Resolving seasonality in tropical trees: multi-decade
544 high-resolution oxygen and carbon isotope records from Indonesia and Thailand. *Earth Planet
Sci Lett* 218:301–316
- 545 Quinn TE, Sampson D (2002) A multi-proxy approach to reconstructing sea surface conditions
546 using coral skeleton geochemistry. *Paleoceanography* 17:1062. doi:10.1029/2000PA000528
- 547 Rutherford S, Mann ME, Osborn TJ, Bradley RS, Briffa KR, Hughes MK, Jones PD (2005) Proxy-
548 based Northern Hemisphere surface temperature reconstructions: sensitivity to methodology,
predictor network, target season, and target domain. *J Climate* 18:2308–2329
- 549 Schweingruber FH (1996) *Tree rings and environment. Dendroecology.* Haupt, Berne
- 550 Souney JM, Mayewski PA, Goodwin ID, Meeker LD, Morgan V, Curran MAJ, van Ommen TD,
551 Palmer AS (2002) A 700-year record of atmospheric circulation developed from the Law Dome,
552 East Antarctica. *J Geophys Res* 107: D22, 4608. doi:10.1029/2002JD002104
- 553 Stahle DW, Cleaveland MK (1992) Reconstruction and analysis of spring rainfall over the
554 southeastern U.S. for the past 1000 years. *Bull Am Meteorol Soc* 73:1947–1961
- 555 Stahle DW, Cook ER, Cleaveland MK, Therrell MD, Meko DM, Grissino-Mayer HD, Watson E,
556 Luckman BH (2000) Tree-ring data document 16th century megadrought over North America.
Eos 81(12):121,125
- 557 Stenni B, Proposito M, Gragnani R, Flora O, Jouzel J, Faour S, Frezzotti M (2002) Eight centuries
558 of volcanic signal and climate change at Talos Dome (East Antarctica). *J Geophys Res* 107:D9.
doi:10.1029/2000JD000317
- 559 Thompson LG (1993) Ice core evidence from Peru and China. In: Bradley RS, Jones PD (eds)
560 *Climate since AD 1500.* Routledge, London, pp 517–548
- 561 Thompson LG, Henderson KA, Mosley-Thompson E, Lin P-N (2000a) The tropical ice core
562 record of ENSO. In: Diaz HF, Markgraf V (eds) *El Niño and the Southern Oscillation: mul-
563 tiscala variability and global and regional impacts.* Cambridge University Press, Cambridge,
pp 325–356
- 564 Thompson LG, Mosley-Thompson E, Henderson K (2000b) Ice core paleoclimate records in South
565 America since the Last Glacial Maximum. *J Quat Sci* 15:377–394
- 566 Vaganov SG, Hughes MK, Shaskin AV (2006) *Growth dynamics of conifer tree rings.* Springer,
567 Berlin, Heidelberg, New York
- 568 Wang Y, Cheng H, Edwards LR, He Y, Kong X, An Z, Wu J, Kelly MJ, Dykoski CA, Li X (2005)
569 The Holocene Asian monsoon: links to solar changes and North Atlantic climate. *Science*
308:854–857
- 570 Wilkinson C, Linden O, Cesar H, Hodgson G, Rubens J, Strong AE (1999) Ecological and socioe-
571 conomic impacts of 1998 coral mortality in the Indian Ocean: An ENSO impact and a warning
572 of future change? *Ambio* 28(2):188–196
- 573 Zielinski GA, Mayewski PA, Meeker LD, Whitlow SI, Twickler SM, Morrison M, Meese DA,
574 Gow AJ, Alley RB (1994) Record of volcanism since 7000 BC from the GISP2 Greenland ice
575 core and implications for the volcano-climate system. *Science* 264:948–952
- 576
- 577
- 578
- 579
- 580
- 581
- 582
- 583
- 584
- 585

Chapter 1

586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630

Q. No.	Query
AQ1	Since figure 1.2 will be printed in black and white, please provide the colour references in grey scale format. Please avoid the usage of colour codes

UNCORRECTED PROOF