

Sichuan University

From the Selected Works of G.C. Wu

Winter December 20, 2013

Variational iteration method for the Burgers' flow with fractional derivatives—New Lagrange multipliers

Guo-Cheng Wu, *Neijiang Normal University*
Dumitru Baleanu



Available at: <https://works.bepress.com/gcwu/7/>



Variational iteration method for the Burgers' flow with fractional derivatives—New Lagrange multipliers



Guo-Cheng Wu ^{a,b,*}, Dumitru Baleanu ^{c,d,e,*}

^a College of Mathematics and Information Science, Neijiang Normal University, Neijiang 641112, China

^b College of Water Resources and Hydropower, Sichuan University, Chengdu 610065, China

^c Department of Mathematics and Computer Sciences, Cankaya University, 06530 Balgat, Ankara, Turkey

^d Institute of Space Sciences, Magurele-Bucharest, Romania

^e Department of Chemical and Materials Engineering, Faculty of Engineering, King Abdulaziz University, Jeddah, Saudi Arabia

ARTICLE INFO

Article history:

Received 24 May 2012

Received in revised form 12 November 2012

Accepted 18 December 2012

Available online 5 January 2013

Keywords:

Fractional differential equation

Caputo derivative

Variational iteration method

ABSTRACT

The flow through porous media can be better described by fractional models than the classical ones since they include inherently memory effects caused by obstacles in the structures. The variational iteration method was extended to find approximate solutions of fractional differential equations with the Caputo derivatives, but the Lagrange multipliers of the method were not identified explicitly. In this paper, the Lagrange multiplier is determined in a more accurate way and some new variational iteration formulae are presented.

© 2013 Elsevier Inc. All rights reserved.

1. Introduction

The diffusion process has been observed in many real physical systems such as highly ramified media in porous systems, anomalous diffusion in fractals and heat transfer close to equilibrium. Fractional calculus is a power tool for finding solution of non-linear problems. Some numerical methods [1–6] and analytical methods [7–12] have been developed for fractional differential equations (FDEs).

The variational iteration method (VIM) [13,14] was extended to FDEs and has been one of the methods used most often. Generally speaking, the use of the variational iteration method (VIM) follows the three steps: (a) to establish the correction functional; (b) identification of the Lagrange multipliers; (c) determination of the initial iteration. Obviously, the step (b) is crucial to derive a variational iteration formula.

Noting the applications of the VIM [13,15] only handled the term of fractional derivatives as restricted variations, we consider a more general FDE

$${}_0^C D_t^\alpha u + R[u] + N[u] = f(t), \quad (1)$$

where ${}_0^C D_t^\alpha$ is the Caputo derivative, $R[u]$ is a linear term and $N[u]$ is a nonlinear one. Momani and Inc et al. [16–20] applied the VIM to the above equation and suggested a variational iteration formula

$$\begin{cases} u_{n+1} = u_n + \int_0^t \lambda(t, \tau) ({}_0^C D_t^\alpha u_n + R[u_n] + N[u_n] - f(\tau)) d\tau, & 0 < t, 0 < \alpha \leq 1, \\ \lambda(t, \tau) = -1, \end{cases} \quad (2)$$

* Corresponding authors. Address: College of Mathematics and Information Science, Neijiang Normal University, Neijiang 641112, China (G.-C. Wu). Address: Department of Mathematics and Computer Sciences, Cankaya University, 06530 Balgat, Ankara, Turkey (D. Baleanu).

E-mail addresses: wuguocheng2002@yahoo.com.cn (G.-C. Wu), dumitru@cankaya.edu.tr (D. Baleanu).

where the function $\lambda(t, \tau)$ is called the Lagrange multiplier.

This paper gives a new way to identify the Lagrange multiplier and improves the variational iteration formula (2) as

$$\begin{cases} u_{n+1} = u_n + \int_0^t \lambda(t, \tau) ({}^C_0 D_t^\alpha u_n + R[u_n] + N[u_n] - f(\tau)) d\tau, & 0 < t, \quad 0 < \alpha, \\ \lambda(t, \tau) = \frac{(-1)^\alpha (\tau - t)^{\alpha-1}}{\Gamma(\alpha)}. \end{cases} \quad (3)$$

The above iteration formula is also valid for differential equations when α is an arbitrary positive integer.

2. Preliminaries

Definition 2.1. The Caputo derivative is given as

$${}^C_0 D_t^\alpha u = \frac{1}{\Gamma(m - \alpha)} \int_0^t \frac{1}{(t - \tau)^{\alpha - m + 1}} u^{(m)}(\tau) d\tau, \quad 0 < t, \quad m = [\alpha] + 1. \quad (4)$$

Definition 2.2. The Riemann–Liouville (R–L) integration of u is defined as

$${}_0 I_t^\alpha u(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t - \tau)^{\alpha-1} u(\tau) d\tau, \quad 0 < t, \quad 0 < \alpha. \quad (5)$$

Definition 2.3. Laplace transform of the term ${}^C_0 D_t^\alpha u$ is given as

$$L[{}^C_0 D_t^\alpha u] = s^\alpha \bar{u}(s) - \sum_{k=0}^{m-1} u^{(k)}(0^+) s^{\alpha-1-k}, \quad m - 1 < \alpha \leq m, \quad (6)$$

where L is Laplace transform and $\bar{u}(s) = L[u(t)]$.

Assuming $\bar{h}(s) = L[h(t)]$ and $\bar{g}(s) = L[g(t)]$, the convolution theorem is

$$h(t) * g(t) = \int_0^t h(t - \tau) g(\tau) d\tau, \quad (7)$$

and

$$\bar{h}(s) \bar{g}(s) = L[h(t) * g(t)]. \quad (8)$$

The detail properties of fractional calculus and Laplace transform can be found in [21–23], respectively.

3. Some new Lagrange multipliers

Theorem 3.1. If the correction functional for Eq. (1) is established via the R–L integration

$$u_{n+1} = u_n + {}_0 I_t^\alpha \lambda(t, \tau) [{}^C_0 D_t^\alpha u_n + R[u_n] + N[u_n] - f(\tau)], \quad (9)$$

the terms $R[u_n]$ and $N[u_n]$ are restricted variations, the Lagrange multiplier can be identified as

$$\lambda(t, \tau) = -1. \quad (10)$$

Proof. Take Laplace transform (6) on the both sides of Eq. (9)

$$\bar{u}_{n+1}(s) = \bar{u}_n(s) + L[{}_0 I_t^\alpha \lambda(t, \tau) ({}^C_0 D_t^\alpha u_n + R[u_n] + N[u_n] - f(\tau))]. \quad (11)$$

Consider the term

$${}_0 I_t^\alpha \lambda_0 {}^C_0 D_t^\alpha u_n = \frac{1}{\Gamma(\alpha)} \int_0^t (t - \tau)^{\alpha-1} \lambda(t, \tau) {}^C_0 D_t^\alpha u_n(\tau) d\tau. \quad (12)$$

Setting the Lagrange multiplier $\lambda(t, \tau) = \lambda(X) / X_{X=t-\tau}$, Eq. (12) is the convolution of the function $a(t) = \frac{\lambda(t) t^{\alpha-1}}{\Gamma(\alpha)}$ and the term ${}_0^C D_t^\alpha u_n(t)$. The terms $R[u_n]$ and $N[u_n]$ are considered as restricted variations which implies $\delta R[u_n] = 0$ and $\delta N[u_n] = 0$, respectively. Make the correction functional (11) stationary with respect to $\bar{u}_n(s)$ and take the classical variation derivative δ on the both sides of Eq. (11). Then Eq. (11) can be calculated as

$$\delta \bar{u}_{n+1}(s) = \delta \bar{u}_n(s) + \delta \left[\bar{a}(s) s^\alpha \bar{u}_n(s) - \sum_{k=0}^{m-1} u^{(k)}(0^+) s^{\alpha-1-k} \right] = (1 + \bar{a}(s) s^\alpha) \delta \bar{u}_n(s). \quad (13)$$

From Eq. (13), we can obtain the equation

$$1 + \bar{a}(s) s^\alpha = 0, \quad (14)$$

which results in

$$\bar{a}(s) = -\frac{1}{s^\alpha} \text{ and } a(t) = -\frac{t^{\alpha-1}}{\Gamma(\alpha)}. \quad (15)$$

□

As a result, the Lagrange multiplier can be identified as

$$\lambda(t, \tau) = a(t - \tau) \Gamma(\alpha) (t - \tau)^{1-\alpha} = -1. \quad (16)$$

The iteration formula (9) reads

$$u_{n+1} = u_n - {}_0I_t^\alpha [{}_0^C D_t^\alpha u_n + R[u_n] + N[u_n] - f(\tau)]. \quad (17)$$

We can check our iteration formula's validness through the relaxation oscillator equation [24]

$${}_0^C D_t^\alpha u + \omega^\alpha u = 0, \quad u(0) = 1, \quad u'(0) = 0, \quad 0 < t, \quad 1 < \alpha \leq 2, \omega > 0. \quad (18)$$

The iteration formula of Eq. (18) can be given as

$$\begin{cases} u_{n+1} = u_n - {}_0I_t^\alpha ({}_0^C D_t^\alpha u_n + \omega^\alpha u_n), \\ u_0 = 1. \end{cases} \quad (19)$$

As a result, we can obtain the series solution

$$\begin{aligned} u_0(t) &= 1, \\ u_1(t) &= 1 - \frac{\omega^\alpha t^\alpha}{\Gamma(1+\alpha)}, \\ u_2(t) &= 1 - \frac{\omega^\alpha t^\alpha}{\Gamma(1+\alpha)} + \frac{\omega^{2\alpha} t^{2\alpha}}{\Gamma(1+2\alpha)}, \\ &\dots \end{aligned}$$

For $n \rightarrow \infty$, $u_n(t) = \sum_{k=0}^n \frac{(-\omega t)^{k\alpha}}{\Gamma(1+k\alpha)}$ rapidly tends to the exact solution $E_\alpha(-(\omega t)^\alpha)$ which is the Mittag-Leffler function.

Theorem 3.2. If the correction functional for Eq. (1) is established via the Riemann integration

$$u_{n+1} = u_n + \int_0^t \lambda(t, \tau) [{}_0^C D_t^\alpha u_n + R[u_n] + N[u_n] - f(\tau)] d\tau, \quad (20)$$

and the terms $R[u_n]$ and $N[u_n]$ are restricted variations, the Lagrange multiplier can be identified as

$$\lambda(t, \tau) = \frac{(-1)^\alpha (\tau - t)^{\alpha-1}}{\Gamma(\alpha)}. \quad (21)$$

From Eq. (17), we can derive

$$\begin{aligned} u_{n+1} &= u_n - {}_0I_t^\alpha [{}_0^C D_t^\alpha u_n + R[u_n] + N[u_n] - f(\tau)] = u_n - \int_0^t \frac{(t - \tau)^{\alpha-1}}{\Gamma(\alpha)} [{}_0^C D_t^\alpha u_n + R[u_n] + N[u_n] - f(\tau)] d\tau \\ &= u_n + \int_0^t \frac{(-1)^\alpha (\tau - t)^{\alpha-1}}{\Gamma(\alpha)} [{}_0^C D_t^\alpha u_n + R[u_n] + N[u_n] - f(\tau)] d\tau. \end{aligned}$$

As a result, for Eq. (20), the Lagrange multiplier is derived

$$\lambda(t, \tau) = \frac{(-1)^\alpha (\tau - t)^{\alpha-1}}{\Gamma(\alpha)},$$

which completes the proof of (21).

One can check that the following iteration formula (See Eq. (25a) in [25]) for the ordinary differential equation (ODE) $\frac{d^m u}{dt^m} + R[u] + N[u] = f(t)$,

$$\begin{cases} u_{n+1} = u_n + \int_0^t \lambda(t, \tau) \left(\frac{d^m u}{d\tau^m} + R[u_n] + N[u_n] - f(\tau) \right) d\tau, & 0 < \alpha, \\ \lambda(t, \tau) = \frac{(-1)^m (\tau-t)^{m-1}}{(m-1)!}, \end{cases} \tag{22}$$

is a special case of Eq. (3). We can conclude that our iteration formula is a “uniform” one for both ODEs and FDEs.

Remarks:

- (I) The iteration formula is also valid for approximately solving FDEs of arbitrary order in sense of the R–L derivative ${}^{RL}D_t^\alpha$ and the sequential derivatives. The differences are the initial iterations.
- (II) The Lagrange multiplier presented in (3) is a simplest one. If we consider more terms in $R[u]$ in Eq. (1), more explicit Lagrange multipliers can be identified. For example, consider the FDE

$${}_0^C D_t^\alpha u + \omega^\alpha u + N[u] = f(t), 0 < \alpha \text{ and } {}^{RL}D_t^\alpha u + \omega^\alpha u + N[u] = f(t), 0 < \alpha. \tag{23}$$

Similarly, the following variational iteration formulae can be given as

$$\begin{cases} u_{n+1} = u_n + \int_0^t \lambda(t, \tau) ({}_0^C D_\tau^\alpha u_n + \omega^\alpha u_n + N[u_n] - f(\tau)) d\tau, \\ \lambda = -(t - \tau)^{\alpha-1} E_{\alpha, \alpha}(-\omega^\alpha (t - \tau)^\alpha), \end{cases} \tag{24}$$

and

$$\begin{cases} u_{n+1} = u_n + \int_0^t \lambda(t, \tau) ({}_0^{RL} D_\tau^\alpha u_n + \omega^\alpha u_n + N[u_n] - f(\tau)) d\tau, \\ \lambda = -(t - \tau)^{\alpha-1} E_{\alpha, \alpha}(-\omega^\alpha (t - \tau)^\alpha). \end{cases} \tag{25}$$

For $\alpha = 1$ and $\alpha = 2$, Eqs. (24) and (25) reduce to the results in (see the iteration formula in [25])

$$\begin{cases} u_{n+1} = u_n + \int_0^t \lambda(t, \tau) \left(\frac{du_n}{d\tau} + \omega u_n + N[u_n] - f(\tau) \right) d\tau, \\ \lambda = -(t - \tau)^{\alpha-1} E_{\alpha, \alpha}(-\omega^\alpha (t - \tau)^\alpha) \Big|_{\alpha=1} = -e^{\omega(\tau-t)}, \end{cases}$$

and

$$\begin{cases} u_{n+1} = u_n + \int_0^t \lambda(t, \tau) \left(\frac{d^2 u_n}{d\tau^2} + \omega^2 u_n + N[u_n] - f(\tau) \right) d\tau, \\ \lambda = -(t - \tau)^{\alpha-1} E_{\alpha, \alpha}(-\omega^\alpha (t - \tau)^\alpha) \Big|_{\alpha=2} = \frac{\sin(\omega(\tau-t))}{\omega}, \end{cases}$$

where $E_{\alpha, \beta}(t)$ is the Mittag-Leffler function with two parameters.

The Lagrange multipliers (24) and (25) lead to approximate solutions of higher accuracies than the results from the variational iteration formula (3). For example, only with one step, one can derive the exact solution of (18).




Fig. 1. $g = \sin(2\pi x)$, $\alpha = 0.9$, $\nu = 0.1$.

(III) The simplest variational iteration formula (17) can reduce to the Volterra integral equation and the analysis of the convergence and the existence of the solutions can be found in [26].

4. Approximate solutions of the Burgers' flow with fractional derivatives

The classical Burgers equation often appears in traffic flow and gas dynamics. Recently, some researchers considered various fractional Burgers equation to model the diffusion behaviors of the flow through porous medium [27–31]. In this section, the VIM is applied to the time-fractional Burgers equation [10]

$${}_0^C D_t^\alpha u + u \frac{\partial u}{\partial x} = \nu \frac{\partial^2 u}{\partial x^2}, \quad 0 < t, \quad 0 \leq x \leq 1, \quad 0 < \alpha \leq 1, \quad (26)$$

where u is the flow's velocity and ν is the viscosity coefficient. It is revealed that the effect of the fractional derivative accumulates slowly to give rise to a significant dissipation.




Fig. 2. $g = \sin(2\pi x)$, $\alpha = 0.9$, $\nu = 0.5$.




Fig. 3. $g = \sin(2\pi x)$, $\alpha = 0.9$, $\nu = 0.9$.

We consider the Burgers equations with the initial condition $u(x, 0) = g(x)$. From Eq. (17), we can have

$$\begin{cases} u_{n+1} = u_n - \int_0^t \frac{(t-\tau)^{\alpha-1}}{\Gamma(\alpha)} \left(\zeta D_\tau^\alpha u_n + u_n \frac{\partial u_n}{\partial x} - \nu \frac{\partial^2 u_n}{\partial x^2} \right) d\tau, & 0 < \alpha \leq 1, \\ u_0 = u(x, 0) = g(x), & u(0, t) = u(1, t) = 0. \end{cases} \tag{27}$$

More generally, for the time-fractional couple Burgers equations of fractional order in [32]

$$\begin{cases} \zeta D_t^\alpha u + 2u \frac{\partial u}{\partial x} - \frac{\partial^2 (uv)}{\partial x^2} = 0, & 0 < \alpha, \\ \zeta D_t^\beta v + 2v \frac{\partial v}{\partial x} - \frac{\partial^2 (uv)}{\partial x^2} = 0, & 0 < \beta, \end{cases} \tag{28}$$

the variational iteration formula can be given as

$$\begin{cases} u_{n+1} = u_n + \int_0^t \lambda_1(t, \tau) \left(\zeta D_\tau^\alpha u_n + 2u_n \frac{\partial u_n}{\partial x} - \frac{\partial^2 u_n v_n}{\partial x^2} \right) d\tau, & \lambda_1(t, \tau) = \frac{(-1)^\alpha (\tau-t)^{\alpha-1}}{\Gamma(\alpha)}, & 0 < \alpha, \\ v_{n+1} = v_n + \int_0^t \lambda_2(t, \tau) \left(\zeta D_\tau^\beta v_n + 2v_n \frac{\partial v_n}{\partial x} - \frac{\partial^2 u_n v_n}{\partial x^2} \right) d\tau, & \lambda_2(t, \tau) = \frac{(-1)^\beta (\tau-t)^{\beta-1}}{\Gamma(\beta)}, & 0 < \beta, \end{cases} \tag{29}$$




Fig. 4. $g = \sin(2\pi x)$, $\alpha = 0.5$, $\nu = 0.5$.




Fig. 5. $g = \sin(2\pi x)$, $\alpha = 0.9$, $\nu = 0.5$.




Fig. 6. $g = \sin(2\pi x)$, $\alpha = 0.99$, $\nu = 0.5$.

For Eq. (26), the successive approximate solutions can be obtained

$$\begin{aligned}
 u_0 &= g(x) = g, \\
 u_1 &= g - (gg' - \nu g^{(2)}) \frac{t^\alpha}{\Gamma(\alpha+1)}, \\
 u_2 &= g - (gg' - \nu g^{(2)}) \frac{t^\alpha}{\Gamma(1+\alpha)} + \left(2gg'^2 + g^2g^{(2)} - 2\nu gg^{(3)} - 4\nu g'g^{(2)} + \nu^2 g^{(4)} \right) \frac{t^{2\alpha}}{\Gamma(1+2\alpha)} \\
 &\quad - (gg' - \nu g^{(2)}) \left(g'^2 + gg^{(2)} - \nu g^{(3)} \right) \frac{\Gamma(1+2\alpha)}{\Gamma^2(1+\alpha)} \frac{t^{3\alpha}}{\Gamma(1+3\alpha)}, \\
 &\vdots
 \end{aligned}$$

where g' denotes $\frac{dg}{dx}$ and $g^{(m)} = \frac{d^m g}{dx^m}$.

We get the approximate solution u_2 as the second term approximation. For $g(x) = \sin(2\pi x)$ and the fractional order $\alpha = 0.9$, Figs 1–3 show the velocity of the flow with various viscosity coefficients ν . For $g(x) = \sin(2\pi x)$ and the viscosity coefficient $\nu = 0.5$, Figs 4–6 illustrate the velocity at different fractional orders.

5. Conclusions

FDEs have been proven to be a useful tool to describe the nonlocal diffusion of the flow in porous media. As one of the analytical methods in FDEs, the existing applications of the VIM in FDEs handled the terms of fractional derivatives as restricted variations or directly employed the one for ODEs. So the Lagrange multipliers determined in that way are not good enough to obtain the approximate solutions of high accuracies. The main reason is that it is difficult for one to use the integration by parts to derive the Lagrange multipliers explicitly.

In this study, the Lagrange multiplier is identified by Laplace transform and such a drawback is overcome. The VIM for FDEs is completed now.

Furthermore, with the result in this study, the following aspects can be considered in future work:

- New analytical methods employing other linearized techniques, for example, the Adomian decomposition series, which can handle the nonlinear terms of FDEs and improve the accuracies of approximate solutions;
- To develop numerical algorithms of fractional partial differential equations based on the VIM which can fully use the merits of the method;
- To consider other applications of the VIM in new non-classical models such as fractional fuzzy equations, fractional time-delay models, the fractional q -difference equations and other dynamical equations on time scales.

Acknowledgment

The first author would like to feel grateful to the referees' suggestions. This work is financially supported by the Key Program of NSFC (No. 51134018).

References

- [1] K. Diethelm, N.J. Ford, A.D. Freed, A predictor–corrector approach for the numerical solution of fractional differential equations, *Nonlinear Dyn.* 29 (2002) 3–22.
- [2] K. Diethelm, N.J. Ford, Multi-order fractional differential equations and their numerical solution, *Appl. Math. Comput.* 154 (2004) 621–640.
- [3] F.W. Liu, V. Anh, I. Turner, Numerical solution of the space fractional Fokker–Planck equation, *J. Comput. Appl. Math.* 166 (2004) 209–219.
- [4] C. Tadjeran, M.M. Meerschaert, H.P. Scheffler, A second-order accurate numerical approximation for the fractional diffusion equation, *J. Comput. Phys.* 213 (2006) 205–213.
- [5] W. Deng, Short memory principle and a predictor–corrector approach for fractional differential equations, *J. Comput. Appl. Math.* 206 (2007) 174–188.
- [6] D. Baleanu, K. Diethelm, E. Scalas, J. Trujillo, *Fractional Calculus Models and Numerical Methods*, Complexity, Nonlinearity and Chaos, World Scientific, Boston, Mass, USA, 2012.
- [7] N.T. Shawagfeh, Analytical approximate solutions for nonlinear fractional differential equations, *Appl. Math. Comput.* 131 (2002) 517–529.
- [8] S. Momani, Z. Odibat, Homotopy perturbation method for nonlinear partial differential equations of fractional order, *Phys. Lett. A* 365 (2007) 345–350.
- [9] P. Zhuang, F.W. Liu, V. Anh, I. Turner, New solution and analytical techniques of the implicit numerical method for the anomalous subdiffusion equation, *SIAM J. Numer. Anal.* 46 (2008) 1079–1095.
- [10] C.P. Li, Y.H. Wang, Numerical algorithm based on Adomian decomposition for fractional differential equations, *Comput. Math. Appl.* 57 (2009) 1672–1681.
- [11] J. Hristov, Approximate solutions to fractional subdiffusion equations, *Eur. Phys. J. Special Topics* 193 (2011) 229–243.
- [12] J.S. Duan, R. Rach, D. Baleanu, A.M. Wazwaz, A review of the Adomian decomposition method and its applications to fractional differential equations, *Commun. Fract. Calc.* 3 (2012) 73–99.
- [13] J.H. He, Approximate analytical solution for seepage flow with fractional derivatives in porous media, *Comput. Methods Appl. Mech. Eng.* 167 (1998) 57–68.
- [14] J.H. He, Variational iteration method – a kind of non-linear analytical technique: some examples, *Int. J. Non-Linear Mech.* 34 (1999) 699–708.
- [15] S. Abbasbandy, An approximation solution of a nonlinear equation with Riemann–Liouville’s fractional derivatives by He’s variational iteration method, *J. Comput. Appl. Math.* 207 (2007) 53–58.
- [16] S. Momani, Z. Odibat, Analytical approach to linear fractional partial differential equations arising in fluid mechanics, *Phys. Lett. A* 355 (2006) 271–279.
- [17] M. Inc, The approximate and exact solutions of the space- and time-fractional Burgers equations with initial conditions by variational iteration method, *J. Math. Anal. Appl.* 345 (2008) 476–484.
- [18] M.R. Yulita, M.S.M. Noorani, I. Hashim, Variational iteration method for fractional heat- and wave-like equations, *Nonlinear Anal. Real World Appl.* 10 (2009) 1854–1869.
- [19] Y. Nawaz, Variational iteration method and homotopy perturbation method for fourth-order fractional integro–differential equations, *Comput. Math. Appl.* 61 (2011) 2330–2341.
- [20] M.G. Sakar, F. Erdogan, A. Yildirim, Variational iteration method for the time-fractional Fornberg–Whitham equation, *Comput. Math. Appl.* 63 (2012) 1382–1388.
- [21] I. Podlubny, *Fractional Differential Equations*, Academic press, San Diego, 1999.
- [22] L. Debnath, D. Bhatta, *Integral Transforms and their Applications*, Chapman & Hall/CRC, 2006.
- [23] H. Sheng, Y. Li, Y.Q. Chen, Application of numerical inverse Laplace transform algorithms in fractional calculus, *J. Franklin Inst.* 348 (2011) 315–330.
- [24] F. Mainardi, Fractional relaxation–oscillation and fractional diffusion–wave phenomena, *Chaos Solitons Fractals* 7 (1996) 1461–1477.
- [25] J.H. He, X.H. Wu, Variational iteration method: new development and applications, *Comput. Math. Appl.* 54 (2007) 881–894.
- [26] K. Diethelm, *The Analysis of Fractional Differential Equations: An Application–Oriented Exposition using Differential Operators of Caputo type*, Springer, 2010.
- [27] M.D. Ruiz-Medina, J.M. Angulo, V.V. Anh, Scaling limit solution of a fractional Burgers equation, *Stochastic Process. Appl.* 93 (2001) 285–300.
- [28] T. Hayat, M. Khan, S. Asghar, On the MHD flow of fractional generalized Burgers’ fluid with modified Darcy’s law, *Acta. Mech. Sin.* 23 (2007) 257–261.
- [29] C. Xue, J. Nie, W. Tan, An exact solution of start-up flow for the fractional generalized Burgers’ fluid in a porous half-space, *Nonlinear Anal.-Theory Methods Appl.* 69 (2008) 2086–2094.
- [30] M. Khan, S.H. Ali, H. Qi, On accelerated flows of a viscoelastic fluid with the fractional Burgers’ model, *Nonlinear Anal.-Real World Appl.* 10 (2009) 2286–2296.
- [31] S.H.A.M. Shah, Some helical flows of a Burgers’ fluid with fractional derivative, *Meccanica* 45 (2010) 143–151.
- [32] Y. Chen, H.L. An, Numerical solutions of coupled Burgers equations with time-and space-fractional derivatives, *Appl. Math. Comput.* 200 (2008) 87–95.