

Howard University

From the Selected Works of Hemayet Ullah

June, 2001

G Protein Regulation of Ion Channels and Abscisic Acid Signaling in Arabidopsis Guard Cells

Hemayet Ullah, *Howard University*

Xi-Qing Wang

Alan M Jones

Sarah M Assmann



Available at: <https://works.bepress.com/hemayet-ullah/4/>

G Protein Regulation of Ion Channels and Abscisic Acid Signaling in *Arabidopsis* Guard Cells

Xi-Qing Wang,¹ Hemayet Ullah,² Alan M. Jones,²
Sarah M. Assmann^{1*}

The phytohormone abscisic acid (ABA) promotes plant water conservation by decreasing the apertures of stomatal pores in the epidermis through which water loss occurs. We found that *Arabidopsis thaliana* plants harboring transferred DNA insertional mutations in the sole prototypical heterotrimeric GTP-binding (G) protein α subunit gene, *GPA1*, lack both ABA inhibition of guard cell inward K^+ channels and pH-independent ABA activation of anion channels. Stomatal opening in *gpa1* plants is insensitive to inhibition by ABA, and the rate of water loss from *gpa1* mutants is greater than that from wild-type plants. Manipulation of G protein status in guard cells may provide a mechanism for controlling plant water balance.

Heterotrimeric G proteins are key regulators of ion channels in animal cells (1, 2). Upon activation, the G protein α subunit ($G\alpha$) binds GTP, resulting in separation of the α subunit from the $\beta\gamma$ subunit pair ($G\beta\gamma$). $G\alpha$ and $G\beta\gamma$ can both interact with downstream components of signaling pathways (2, 3). Among the important downstream effectors are K^+ and Ca^{2+} channels, which are regulated by G proteins via both cytosolic signaling cascades and membrane-delimited pathways (1, 2). G protein-mediated ion-channel regulation is an integral component of vision, taste, smell, and hormonal signaling in mammalian systems (2–4).

In higher plants, guard cell ion-channel regulation controls stomatal apertures. Stomatal opening relies on increases in K^+ , Cl^- , malate²⁻, and sucrose in the guard cell symplast to drive water influx and cell swelling. These processes result in an outbowing of the guard cell pair and an increase in pore aperture. During stomatal opening, K^+ uptake is mediated by inwardly rectifying K^+ channels. During inhibition of stomatal opening by the plant hormone abscisic acid (ABA), these channels are inhibited (5–7). In guard cells, ABA activates phospholipases C and D (8–10) and can elevate cytosolic calcium levels via inositol 1,4,5-trisphosphate or other pathways (8, 11–14). Cytosolic Ca^{2+} elevation inhibits inwardly rectifying K^+ channels (6, 7, 15) and activates slow anion channels

that mediate Cl^- and malate²⁻ efflux (15, 16). In mammalian systems, certain phospholipases C and D are regulated by heterotrimeric G proteins. In the *Arabidopsis thaliana* genome, there is only one prototypical $G\alpha$ gene, *GPA1* (17, 18), and this gene is expressed in guard cells (Fig. 1A) (19). Thus, we hypothesized that *GPA1* may regulate ion channels (20) and ABA response (21) in this cell type. We used two independent *Arabidopsis* lines harboring the recessive transferred DNA (T-DNA) knockout alleles *gpa1-1* or *gpa1-2* (22) to test these hypotheses.

Guard cells isolated from homozygous *gpa1-1* and *gpa1-2* plants (23, 24) failed to express full-length *GPA1* transcripts (Fig. 1B), as expected (19, 22). In contrast to the response of wild-type plants, stomata of *gpa1* mutant plants showed no inhibition of stomatal opening (25, 26) by ABA (Fig. 2A). We next used patch clamp techniques to test whether sensitivity of the inward K^+ channels to inhibition by ABA had also been altered in the *gpa1* mutants (27). Just as for stomatal opening, the inward K^+ channels of the mutant plants were ABA insensitive (Fig. 2, B and C). Thus, ABA inhibition of inward K^+ channels and stomatal opening require the presence of functional *GPA1*.

ABA activation of slow anion channels (28–30) is thought to be another component of ABA inhibition of stomatal opening. Anion efflux occurring upon anion channel opening should depolarize the membrane and hinder K^+ uptake. However, the hypothesis that G proteins regulate anion channels in plants had no experimental precedence, and few mammalian anion channels are G protein regulated. Accordingly, we tested whether

ABA activation of slow anion channels was altered in the *gpa1* mutants. We initially used an established protocol for recording the whole-cell anion channel response to ABA, which uses a cytosolic solution with moderate Ca^{2+} and strong pH buffering capacities (29, 31). We found that *GPA1* does regulate plant anion channels: In both *gpa1* mutants, ABA activation of anion channels was abolished (Fig. 3, A and B).

Activation of anion channels also promotes stomatal closure by mediating loss of anionic solutes and membrane depolarization that drives K^+ efflux (28–30, 32, 33). Thus, we expected that ABA promotion of stomatal closure (34) would also be eliminated in *gpa1* mutant plants, but it was not (Fig. 3C) (35). The uncoupling of ABA inhibition of stomatal opening (Fig. 2A) and ABA promotion of stomatal closure (Fig. 3C) demonstrates that these two effects are not simply the reverse of one another (28).

Consideration of the anion channel and stomatal closure experiments led to the interpretation that a parallel or compensatory pathway mediating ABA promotion of stomatal closure was present in the intact *gpa1* guard cells (Fig. 3C) but was nonfunctional under the conditions of the patch clamp experiments (Fig. 3, A and B). One candidate member of such a pathway is cytosolic pH. Cytosolic pH was strongly buffered in the protocol of Fig. 3, A and B, yet ABA elevates cytosolic pH in intact guard cells (34, 36).




Fig. 1. RT-PCR analysis of *GPA1* expression in guard cells of wild-type and *gpa1* mutant lines. (A) *GPA1* is more highly expressed in guard cell protoplasts (GCP) than in mesophyll cell protoplasts (MCP) of wild-type *Arabidopsis*. Different cycle numbers were performed as indicated to assess *GPA1* expression levels. Expected band sizes of 605 bp (*GPA1*) and 1 kb (*Actin*) are observed; the identity of *GPA1* products was confirmed by sequencing. (B) *GPA1* is expressed in wild-type guard cells and not in the guard cells of *gpa1* knockout mutants, as assessed by RT-PCR. *GPA1* primers (lanes 2, 4, and 6) bracketed the T-DNA insertion site; thus, no PCR product was expected from the *gpa1* lines. Actin bands indicate relative cDNA amounts.

¹Biology Department, Pennsylvania State University, 208 Mueller Laboratory, University Park, PA 16802–5301, USA. ²Department of Biology, University of North Carolina, Chapel Hill, NC 27599–3280, USA.

*To whom correspondence should be addressed. E-mail: sma3@psu.edu


REPORTS

Therefore, we next obtained electrophysiological measurements under weak cytosolic pH buffering (31) that would permit observa-

tion of anion channel regulation by ABA-induced alterations in pH. Normal ABA activation of anion channels was observed in all

genotypes (Fig. 3, D and E), consistent with the observation of ABA-induced stomatal closure in epidermes of mutant plants (Fig.


Fig. 2. Mutations *gpa1-1* and *gpa1-2* cause ABA insensitivity of stomatal opening and inward K^+ channel regulation. (A) Comparison of ABA (20 μ M) inhibition of stomatal opening in wild-type, *gpa1-1*, and *gpa1-2*. Error bars represent \pm SE from three independent trials; $n = 40$ apertures per trial. Only wild-type plants showed significant ABA inhibition of stomatal opening (** $P \leq 0.01$, Student's t test). Photographs show representative stomata from the three genotypes after ABA treatment (open bars, -ABA; hatched bars, +ABA). (B) Whole-cell recordings of guard-cell inward K^+ currents in the absence or presence of 50 μ M ABA. K^+ currents were recorded from a holding potential of -76 mV with voltage steps from -216 to $+64$ mV in $+20$ -mV increments. ABA was added immediately after achieving the whole



cell configuration. Recordings were obtained after 10 min. (C) Current/voltage relations (error bars represent \pm SE) showed no ABA inhibition of inward K^+ currents in *gpa1-1* and *gpa1-2* guard cells versus significant ABA inhibition in wild-type guard cells at voltages ≤ -156 mV ($P \leq 0.01$, Student's t test). Voltage protocols were as for (B). $n = 9$ to 15 cells.

cell configuration. Recordings were obtained after 10 min. (C) Current/voltage relations (error bars represent \pm SE) showed no ABA inhibition of inward K^+ currents in *gpa1-1* and *gpa1-2* guard cells versus significant ABA inhibition in wild-type guard cells at voltages ≤ -156 mV ($P \leq 0.01$, Student's t test). Voltage protocols were as for (B). $n = 9$ to 15 cells.

Fig. 3. Mutants *gpa1-1* and *gpa1-2* exhibit complex ABA-induced stomatal closure and ABA regulation of anion currents. (A) ABA (50 μ M) activation of anion currents in *gpa1-1* and *gpa1-2* guard cells under cytosolic conditions of moderate Ca^{2+} buffering and strong pH buffering (31). Slow anion channel currents were recorded from a holding potential of $+30$ mV with voltage steps from -145 to $+35$ mV in $+30$ -mV increments. Currents were identified as slow anion currents by their kinetics and reversal potential (48). (B) Current-voltage relations as recorded in (A), showing that ABA failed to activate anion channel currents in *gpa1-1* and *gpa1-2* guard cells in the presence of strong cytosolic pH buffering [10 mM HEPES-tris (pH 7.5)]. Values are \pm SE; $n = 8$ to 16 cells. Only wild-type cells showed significant ABA stimulation of anion current ($P \leq 0.01$ at voltages ≤ -25 mV). (C) All genotypes showed significant stomatal closure in response to 20 μ M ABA ($P \leq 0.01$, Student's t test). Error bars represent \pm SE from 3 independent trials; $n = 40$ per trial, (open bars, -ABA; hatched bars, +ABA). (D) ABA activation of anion channel currents with a pipette solution identical to that for (A) and (B) except for weak cytosolic pH buffering [0.1 mM HEPES-tris (pH 7.5)] (31). Voltage protocols were as for Fig. 3A. (E) Current-voltage relations (error bars represent \pm SE) showed significant ($P \leq 0.01$ at voltages ≤ -25 mV) ABA



activation of anion channel currents in all genotypes under these conditions. $n = 6$ to 9 cells. (F) Stomatal closure response to ABA in the presence of an intracellular pH clamp imposed by the membrane-permeant weak acid butyrate. Changes in stomatal aperture reflect final minus initial aperture. Error bars represent \pm SE from three independent trials; $n = 40$ per trial. Butyrate significantly reversed ABA responses in all genotypes ($P \leq 0.01$). The butyrate response was significantly greater in *gpa1-1* lines than in wild type ($P \leq 0.01$) and did not differ significantly between *gpa1-1* and *gpa1-2*. Hatched bars, ABA; solid bars, ABA + 1 mM Na-butyrate.

activation of anion channel currents in all genotypes under these conditions. $n = 6$ to 9 cells. (F) Stomatal closure response to ABA in the presence of an intracellular pH clamp imposed by the membrane-permeant weak acid butyrate. Changes in stomatal aperture reflect final minus initial aperture. Error bars represent \pm SE from three independent trials; $n = 40$ per trial. Butyrate significantly reversed ABA responses in all genotypes ($P \leq 0.01$). The butyrate response was significantly greater in *gpa1-1* lines than in wild type ($P \leq 0.01$) and did not differ significantly between *gpa1-1* and *gpa1-2*. Hatched bars, ABA; solid bars, ABA + 1 mM Na-butyrate.

REPORTS

3C). Thus, the pH-independent pathway of ABA action (Fig. 3, A and B) requires *GPA1*, whereas the pH-dependent pathway (Fig. 3, D and E) does not. Consistent with this model, block of the pH-dependent pathway in epidermal peels by a cytosolic pH clamp with butyric acid, a membrane-permeant weak acid (34, 35), opposed ABA-induced stomatal closure in all three genotypes (Fig. 3F).

Stomatal apertures in leaves taken directly from the growth environment (24) were ~20% greater in *gpa1* mutant lines than in wild-type plants, suggesting that stomatal sensitivity to background levels of endogenous ABA is shifted in the *gpa1* mutants. Water loss (37, 38) is greater from *gpa1* leaves than from wild-type leaves (Fig. 4), confirming that elimination of *GPA1* impacts water relations *in planta*. This information may contribute to efforts to engineer plants with improved stomatal regulation.

Previous pharmacological studies of guard cell function with G protein activators such as guanosine 5'-O-(3'-thiotriphosphate) (GTP- γ -S), and inactivators such as guanosine 5'-O-(2'-thiodiphosphate) (GDP- β -S) (18), suggested that active G proteins inactivate the inward K⁺ channels via both cytosolic and membrane-delimited pathways to inhibit stomatal opening (21, 39–41). Such data were challenged, however, by other pharmacological studies suggesting that G protein activation could stimulate stomatal opening (42) or had both stimulatory and inhibitory effects on inward K⁺ currents (43). These results can be clarified by the precision afforded by T-DNA mutagenesis (44), in which one specific protein is eliminated with retention of an otherwise wild-type genetic complement. Reverse genetics also has allowed us to directly test and support the hypothesis




Fig. 4. Water loss is greater from *gpa1* mutant leaves. Water loss is expressed as the percentage of initial fresh weight (37). Values are mean \pm SE (error bars) of measurements with three individual plants per genotype. One of three independent trials is shown. Regression analysis confirmed that the wild-type curve differs significantly from the *gpa1-1* and *gpa1-2* responses at $P \leq 0.01$. Circles, wild type; squares *gpa1-1*; triangles, *gpa1-2*.

that ABA signaling in guard cells uses G protein activation (21). There is evidence that plant G proteins are involved in responses to light (45), pathogens (46), and several hormones (18, 47), including ABA as shown here. It will be of interest to unravel how plant G protein pathways can couple receptors with their cognate downstream effects for such diverse and multiple signals, given that the *Arabidopsis* genome contains only *GPA1* as a prototypical G α subunit gene.

References and Notes

- A. M. Brown, L. Birnbaumer, *Annu. Rev. Physiol.* **52**, 197 (1990).
- K. Wickman, D. E. Clapham, *Physiol. Rev.* **75**, 865 (1995).
- J. R. Hepler, A. G. Gilman, *Trends Biochem. Sci.* **17**, 383 (1992).
- G. M. Sheperd, *Cell* **67**, 845 (1991).
- M. R. Blatt, *Planta* **180**, 445 (1990).
- F. Lemtri-Chlieh, E. A. C. MacRobbie, *J. Membr. Biol.* **137**, 99 (1994).
- X.-Q. Wang, W.-H. Wu, S. M. Assmann, *Plant Physiol.* **118**, 1421 (1998).
- Y. S. Lee *et al.*, *Plant Physiol.* **110**, 987 (1996).
- T. Jacob, S. Ritchie, S. M. Assmann, S. Gilroy, *Proc. Natl. Acad. Sci. U.S.A.* **96**, 12192 (1999).
- I. Staxen *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **96**, 1779 (1999).
- S. Gilroy, N. D. Read, A. J. Trewavas, *Nature* **346**, 769 (1990).
- M. R. McAinsh, C. Brownlee, A. M. Hetherington, *Nature* **343**, 186 (1990).
- C. P. Leckie, M. R. McAinsh, G. J. Allen, D. Sanders, A. M. Hetherington, *Proc. Natl. Acad. Sci. U.S.A.* **95**, 15837 (1998).
- E. A. C. MacRobbie, *Proc. Natl. Acad. Sci. U.S.A.* **97**, 12361 (2000).
- J. I. Schroeder, S. Hagiwara, *Nature* **338**, 427 (1989).
- G. J. Allen, K. Kuchitsu, S. P. Chu, Y. Murata, J. I. Schroeder, *Plant Cell* **11**, 1785 (1999).
- H. Ma, M. F. Yanofsky, E. M. Meyerowitz, *Proc. Natl. Acad. Sci. U.S.A.* **87**, 3821 (1990).
- H. Ma, *Plant Mol. Biol.* **26**, 1611 (1994).
- RNA was extracted from guard cell (purity percentage of 99:1, guard cell protoplasts:mesophyll cell protoplasts) and mesophyll cell protoplast preparations. Reverse transcription followed by polymerase chain reaction (PCR) was performed [SuperScript first-strand synthesis system for reverse transcriptase (RT)-PCR, Gibco BRL, Life Technologies]. The forward primer started at 662 base pairs (bp) in the *GPA1* cDNA clone (forward; 5' ggc tgc tga aat cga aag ac 3'); the reverse primer started at 1266 bp (reverse; 5' gtc cac cca cgt caa aca at 3').
- G. S. Aharon, A. Gelli, W. A. Snedden, E. Blumwald, *FEBS Lett.* **424**, 17 (1998).
- S. M. Assmann, in *Signal Transduction in Plant Growth and Development*, D. P. S. Verma, Ed. (Springer, Vienna, 1996), pp. 39–61.
- H. Ullah *et al.*, *Science* **292**, 2066 (2001).
- L. A. Romano, H. Miedema, S. M. Assmann, *Plant Cell Physiol.* **39**, 1133 (1998).
- We used 4- to 6-week-old plants grown under short days. Seeds were germinated in a standard medium (23) in 7.5% (w/v) phytigel (Sigma), with kanamycin (50 mg/liter) for mutants. After 10 days, robust seedlings were transferred to Metromix 250 potting mixture (Scotts, Marysville, OH).
- J. Leymarie, G. Lasceve, A. Vavasseur, *Aust. J. Plant Physiol.* **25**, 785 (1998).
- Light-induced stomatal opening was assayed according to (25). Normalization of aperture widths by stomatal length (16) mirrored the results shown here for widths alone (48).
- Guard cell protoplasts were isolated from rosette leaves of 4- to 5-week-old, nonbolting plants [after (7)], except that the second enzyme solution contained 1.3% (w/v) Cellulase RS (Yakult Honsha, Tokyo) and 0.0075% (w/v) Pectolyase Y-23 (Seishin Pharmaceutical, Tokyo). Solutions used in patch-clamp experiments on K⁺ channels were as follows: 10 mM MES (tris, pH 6.0), 10 mM K-glutamate, 4 mM MgCl₂, and 1 mM CaCl₂ [osmolality of 540 mmol/kg (bath solution)] and 10 mM Hepes-tris (pH 7.8), 80 mM K-glutamate, and 20 mM KCl [osmolality of 560 mmol/kg (pipette solution)]. Fresh adenosine triphosphate (ATP) (5 mM from a 0.5 M Mg ATP stock solution in 0.5 M tris) was added daily to the pipette solution. Final osmolalities were obtained by addition of sorbitol. Time-activated currents were calculated as the difference between average steady-state current between 3000 and 3800 ms and instantaneous current at 20 ms.
- J. Li, X.-Q. Wang, M. B. Watson, S. M. Assmann, *Science* **287**, 300 (2000).
- Z.-M. Pei, K. Kuchitsu, J. M. Ward, M. Schwarz, J. I. Schroeder, *Plant Cell* **9**, 409 (1997).
- A. Grabov, J. Leung, J. Giraudat, M. R. Blatt, *Plant J.* **12**, 203 (1997).
- The initial pipette solution for anion channel recording was as follows: 150 mM CsCl, 2 mM MgCl₂, 6.7 mM EGTA, 3.35 mM CaCl₂, 10 mM Hepes-tris (pH 7.5), with osmolality of 540 mmol/kg (29). Fresh 5 mM Mg-ATP and 5 mM tris-GTP were added daily (29). When we studied the role of pH in anion channel regulation by ABA, the pH buffer in the pipette solution was 0.1 mM Hepes-tris (pH 7.5). The bath solution was 30 mM CsCl, 2 mM MgCl₂, 5 mM CaCl₂, 10 mM Mes-tris (pH 5.6), with osmolality of 560 mmol/kg. Protoplasts were pretreated for at least 1.5 hours with ABA, and 50 μ M ABA was added to bath and pipette solutions (29). Anion currents were recorded 11 to 14 min after achieving the whole-cell configuration. Steady-state currents were averaged between 52.5 and 62.5 s.
- E. A. C. MacRobbie, *J. Exp. Bot.* **48**, 515 (1997).
- R. Hedrich, H. Busch, K. Raschke, *EMBO J.* **9**, 3889 (1990).
- M. R. Blatt, F. Armstrong, *Planta* **191**, 330 (1993).
- For Fig. 3C, stomatal closure was assayed according to (29), except that after ABA application (20 μ M), leaves were incubated for 3 hours before aperture measurement. For Fig. 3F, we assayed stomatal closure with the protocol of (34), although with minor modification. Leaves were incubated for 2 hours under 450 μ mol m⁻² s⁻¹ of light, followed by the addition of 1 mM Na-butyrate and 20 μ M ABA. Apertures were measured 2 hours later. The wild-type butyrate response (48) was saturated with 1 mM Na-butyrate.
- H. R. Irving, C. A. Gehring, R. W. Parish, *Proc. Natl. Acad. Sci. U.S.A.* **89**, 1790 (1992).
- J. Leung, S. Merlot, J. Giraudat, *Plant Cell* **9**, 579 (1997).
- On the basis of (37), 5 to 6 young leaves were excised from an individual plant, and their total fresh weight was measured during incubation in a chamber through which dry air was passed.
- K. Fairley-Grenot, S. M. Assmann, *Plant Cell* **3**, 1037 (1991).
- W.-H. Wu, S. M. Assmann, *Proc. Natl. Acad. Sci. U.S.A.* **91**, 6310 (1994).
- F. Armstrong, M. R. Blatt, *Plant J.* **8**, 187 (1995).
- H. J. Lee, E. B. Tucker, R. C. Crain, Y. Lee, *Plant Physiol.* **102**, 95 (1993).
- W. K. Kelly, J. E. Esser, J. I. Schroeder, *Plant J.* **8**, 479 (1995).
- P. J. Krysan, J. C. Young, M. R. Sussmann, *Plant Cell* **11**, 2283 (1999).
- K. M. Warpeha, H. E. Hamm, M. M. Rasenick, L. S. Kaufman, *Proc. Natl. Acad. Sci. U.S.A.* **88**, 8925 (1991).
- A. Gelli, V. J. Higgins, E. Blumwald, *Plant Physiol.* **113**, 269 (1997).
- M. Ueguchi-Tanaka *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **97**, 11638 (2000).
- X.-Q. Wang, S. M. Assmann, unpublished observations.
- Supported by USDA grants 98-35304-6681 and 2001-35304-09916 and NSF grant MCB-9874438 to S.M.A. We thank Z.-M. Pei for advice on patch-clamping *Arabidopsis* guard cells.

16 January 2001; accepted 10 April 2001