

**USDA National Wildlife Research Center**

---

**From the Selected Works of Larry Clark**

---

2014

## Chapter 7: The Chemical Senses in Birds

Larry Clark



Available at: [https://works.bepress.com/larry\\_clark/58/](https://works.bepress.com/larry_clark/58/)

# The Chemical Senses in Birds

**Larry Clark**

*United States Department of Agriculture, Animal and Plant Health Inspection Service, Wildlife Services, National Wildlife Research Center, Fort Collins, CO, USA*

**Julie Hagelin**

*Institute of Arctic Biology, University of Alaska Fairbanks, Fairbanks, AK, USA; Alaska Department of Fish and Game, Fairbanks, AK, USA*

**Scott Werner**

*United States Department of Agriculture, Animal and Plant Health Inspection Service, Wildlife Services, National Wildlife Research Center, Fort Collins, CO, USA*

## 7.1 CHEMICAL SENSES

The chemical senses generally fall into three categories: chemesthesis (irritation and pain), olfaction (smell), and gustation (taste). Traditionally, the emphasis in describing responsiveness to chemical stimuli has been placed on taste and smell. The reality is more complex. For example, the sensory afferents for chemesthetic perception are in close proximity with olfactory receptors in the nasal cavity and with gustatory receptors in the oral cavity. Because external chemical stimuli can be processed by multiple sensory systems, there has been a great deal of confusion in the literature on the importance of individual sensory modalities. Generally, the principal mediating sensory modality may be related to stimulus type, concentration, and presentation. However, when perception of external chemical stimuli occurs via the integrated perception across modalities, the combined perceptual quality is commonly referred to as flavor.

## 7.2 CHEMESTHESIS

Chemesthesis is the perception of chemically induced pain. The first neural mediator of noxious stimuli is the nociceptor (Woolf and Ma, 2007). These primary sensory neurons are the interface between the internal and external environments. Nociceptors have cell bodies located in the dorsal root ganglion, a peripheral axon that innervates tissues, and a central axon that enters the spinal cord to transfer information to the central nervous system. Nociceptors have three functions: (1) detection of potentially damaging external noxious stimuli, which is useful in warning an animal

to the risk of injury; (2) detection of endogenous inflammatory stimuli, which is useful in initiating and promoting behaviors conducive to healing and repair; and (3) detection of neural damage and ectopic firing. This latter function is a pathological condition of chronic pain. Nociceptors have high thresholds for exogenous stimuli, presumably because it would be maladaptive to defensively respond to every external assault. Nociceptors have low thresholds for endogenous stimuli. This is an adaptive response to promote healing once damage has occurred (Patapoutian et al., 2009).

A major component of the chemesthetic system is the trigeminal nerve (TN). The TN is the principal somatic sensory nerve of the head, and its primary function is the coding of mechanical and thermal stimuli. However, the TN also contains chemoreceptive fibers that mediate the detection of chemical irritants (Silver and Maruniak, 1981). The somatosensory system is the primary somatic sensory system of the rest of the body. Like the TN, the somatosensory system primarily codes for mechanical and thermal stimuli, but it does have sensory afferents that are chemosensory (Gentle, 2011; Necker, 2000; Wild, 1985).

### 7.2.1 Trigeminal and Somatosensory Nerves

The morphological organization of the peripheral TN in birds is not very different from that found in mammals (Dubbeldam and Karten, 1978; Dubbeldam and Veenman, 1978; Gottschaldt, 1985). The TN is the fifth cranial nerve in birds, arising from the rostralateral medulla near the caudal surface of the optic lobe (Getty, 1975; Schrader, 1970).

The TN travels along the trochlear nerve (IV), entering a fossa in the floor of the cranial cavity where the trigeminal ganglion (TG) is found. The TG is subdivided into a smaller medial ophthalmic region and a larger lateral maxillomandibular region, from which the nerve splits into three branches. In the chicken (*Gallus gallus domesticus*) the ophthalmic branch innervates the frontal region, the eyeball, upper eyelid, conjunctiva, glands in the orbit, the rostrorodorsal part of the nasal cavity, and the tip of the upper jaw. The ophthalmic branch as a communicating ramus with the trochlear nerve serves for motor control of the eye region. This aspect can provide for reflexive response to irritating stimuli to the ocular region. The larger medial ramus accompanies the olfactory nerve into the nasal fossa via the medial orbitonasal foramen. The maxillary branch provides sensory input from the integument of the crown, temporal region, rostral part of the external ear, upper and lower eyelids, the region between the nostrils and eye, conjunctival mucosa, the mucosal part of the palate, and the floor of the medial wall of the nasal cavity. The mandibular branch provides sensory input from the skin and rhamphotheca of the lower jaw, intermandibular skin, wattles, oral mucosa of rostral floor of the mouth, and the palate near the angle of the mouth (Getty, 1975; Schrader, 1970).

### 7.2.2 Performance Characteristics of Nociceptors

Pain and irritation perception begin with activation of primary sensory nociceptors. In birds, chemosensitive fibers in the TN and somatosensory nerves are similar to mammalian afferents. Most are unmyelinated C-type polymodal nociceptors with conduction velocities of 0.3–1 m/s. However, some myelinated A-delta high-threshold mechanoreceptors with conduction velocities of 5–40 m/s also respond to chemical stimuli. The discharge patterns and conduction velocities for the chicken, mallard (*Anas platyrhynchos*), and pigeon (*Columba livia*) are similar to those observed in mammals (Gentle, 1989; Necker, 1974).

Although birds have slightly different neural architecture relative to mammals, the underlying functions of neural connections have been evolutionarily preserved (Butler and Cotterill, 2006; Dugas-Ford et al., 2012; Güntürkün, 2012). This also applies to the underlying physiological and biochemical processes of chemically induced pain. Generally, birds have the same classes of neuropeptides as mammals, but their structures are not totally homologous. Avian endogenous pain-promoting substances such as substance P, 5-HT, histamine, bradykinin, and acetylcholine evoke inflammation and pain-related behaviors in chickens, pigeons, rats, dogs, and guinea pigs (Szolcsanyi et al., 1986; Gentle and Hill, 1987; Gentle and Hunter, 1993; Koda et al., 1996; Hu et al., 2002; Ohta et al., 2006). Prostaglandins that modulate the pain response in mammals also serve this


function in birds, and their effects can be abolished by prostaglandin biosynthesis inhibitors such as aspirin-like analgesics (Clark, 1995).

Despite these physiologically mediated similarities, there are profound differences in how birds and mammals respond to exogenous chemical stimuli. In mammals, chemicals such as capsaicin are potent trigeminal irritants. These irritants deplete substance P from afferent terminals and the dorsal root ganglion, producing an initial sensitization followed by desensitization to further chemical stimulation (Szolcsanyi, 1982). In contrast, birds are insensitive to capsaicin (Mason and Maruniak, 1983; Szolcsanyi et al., 1986). Peripheral presentation of capsaicin to pigeons and chickens does not cause release of substance P in avian sensory afferents (Pierau et al., 1986; Szolcsanyi et al., 1986; Sann et al., 1987). These taxon-specific responses to exogenous chemical stimuli underscore taxonomic differences in both endogenous neuropeptides and receptors, whose significance has been implicated in the evolutionary ecology of the taxa (Mason et al., 1991; Clark, 1998; Tewksbury and Nabhan, 2001).

### 7.2.3 Receptor Mechanisms

Nonselective transient receptor potential (TRP) cation channels are involved in sensory neuron activation events, neurotransmitter release, release of inflammatory mediators, and other aspects of pain transduction (Cortright et al., 2007; Figure 7.1). Most of what is known about TRP channels is derived from work done on mammals (Holzer, 2011). However, increasingly more comparative evolutionary similarities and differences are being characterized for other taxa (Saito and Shingai, 2006; Saito et al., 2011). TRPV1 (initially called VR1) was first cloned in mammals and found to respond to the exogenous vanilloid, capsaicin (Caterina et al., 1997), as well as endogenous agonists, anandamide, and 12-HPETE, which are structurally similar to capsaicin (Zygmunt et al., 1999; Hwang et al., 2000). TRPV1 is also activated by heat (>43 °C) and acid (pH ≤ 6). The sensation that TRPV1 activation evokes in humans via these polymodal nociceptors is one of tingling and burning, like the sensation produced by capsaicin found in chili peppers. Like its mammalian counterpart, the TRP receptor in birds (cTRPV1, chick dorsal root ganglion) responds to high temperatures (≥45 °C) and extracellular acid solution (pH ≤ 4). However, cTRPV1 is different, showing a 68% identity and 79% similarity to rat TRPV1. These differences in receptor composition manifest as a poor response to capsaicin (Jordt and Julius, 2002) and explain the behavioral differences in capsaicin sensitivity between birds and mammals; mammals are behaviorally sensitive to capsaicin and birds are not (Mason et al., 1991; Norman et al., 1992).

Currently, 28 TRP channels, grouped into six functional subfamilies, have been characterized. The subfamilies are




**FIGURE 7.1 Changes in transient receptor potential (TRP) channels produced by inflammation.** Endogenous sensitizers act on receptors expressed by nociceptors to activate intracellular signal transduction pathways. Pathways phosphorylate TRP channels, altering trafficking to the membrane, thresholds, and kinetics. Growth factors, such as nerve growth factor (NGF), are retrogradely transported to the cell body of the nociceptors. Through intracellular signaling pathways, expression of TRP channels is increased and they are transported to the peripheral terminal. Changes in transcription and translation of TRP channels and other proteins can switch the chemical phenotype of the neurons from their state in naive conditions to an altered state during inflammation. B2, bradykinin receptor; ERK, extracellular signal-regulated kinase; ETAR, endothelin receptor type A; GDNF, glial-cell-derived neurotrophic factor; NK1, neurokinin receptor 1; PAR2, protease-activated receptor 2; PGE2, prostaglandin E2; PI3K, phosphoinositide 3-kinase; PK, protein kinase; PKR, prokineticin receptor; TNF $\alpha$ , tumor necrosis factor  $\alpha$ ; TNFR1, TNF receptor 1; TRKA, tyrosine kinase receptor A. *Adapted from Patapoutian et al. (2009).*

responsive to exogenous compounds that code for qualitative perceptual similarities (e.g., the “hotness” of capsaicin, the “burn” of cinnamon oil, the “coolness” of menthol, the irritation of mustard oil; Holzer, 2011). Although the specific homologies for other TRP channels in birds are generally not known, based on behavioral responsiveness to a variety of mammalian irritants, it is anticipated that TRP channel receptor molecules in birds would be structurally similar and/or have similar expression in nociceptors to that found in mammals for cinnamon oil, allicin (garlic/onion), and menthol and divergent for mustard oil and anthranilate (grape) compounds (Clark, 1998; Stucky et al., 2009).

Digital fluorescence imaging of intracellular calcium  $[Ca^{2+}]_i$  in vitro preparations of chicken and rat trigeminal dorsal root ganglia show that there are separate and overlapping populations of neurons that are sensitive to

the well-described avian irritant, methyl anthranilate, and capsaicin (Kirifides et al., 2004). In the chicken, 48% of neurons responded to methyl anthranilate, whereas only 16% responded to capsaicin. Moreover, there was a greater change in  $[Ca^{2+}]_i$  to equimolar concentrations of methyl anthranilate (78%) relative to capsaicin (43%). Increases in  $[Ca^{2+}]_i$  were dependent upon extracellular calcium for both methyl anthranilate and capsaicin. However, responses to methyl anthranilate, but not capsaicin, were dependent on extracellular sodium. This suggests different transduction mechanisms for the two compounds. Together, these observations provide further rationale for the observed behavioral differences in birds to these two compounds. Starlings (*Sturnus vulgaris*) demonstrate congenital avoidance to methyl anthranilate but not capsaicin, although they could be trained to avoid capsaicin in conditioned



**FIGURE 7.2** Consumption of food treated with capsaicin (CAP), methyl capsaicin (MCAP), vanillyl acetamide (VNAC), veratryl acetamide (VRAC), and veratryl amine (VRAM) in rats and starlings. Note the general inverse relationship of consumption as structure changes, suggesting functional receptor differences in the two taxa. Bird repellents are more basic and rigid (planar) than mammal aversive compounds. Concentration applied: 1000 ppm. Consumption of 4 g of untreated food is control baseline intake for both species. *Data adapted from Mason et al. (1991).*


avoidance paradigms, and that avoidance was contingent upon an intact ophthalmic branch of the TN (Mason and Clark, 1995). These observations also suggest that while birds can perceive capsaicin, although somewhat poorly, it is not coded as pain, highlighting the importance of central processing in the perceptual interpretation of peripheral signals.

#### 7.2.4 Chemical Structure–Activity Relationships to Irritants

Despite the apparent insensitivity of birds to capsaicin, they can respond to other vanilloid compounds (Figure 7.2). Aromatic compounds that are considered aversive by birds are qualitatively characterized as having an aromatic heterocyclic core, high degree of basicity, high degree of lipophilicity, and a high degree of electronegativity (Figure 7.3). The core aromatic heterocycle of a repellent compound is enhanced by substitutions that affect electron donation: amino > methoxy > methyl > hydroxyl groups. Resonance of lone pairs of electrons enhances repellency as a function of substituent position: ortho > para > meta. Acidic substituents in the electron withdrawing group detract from aversive qualities of the compound. Steric effects and extreme

delocalization of lone pairs of electrons, as might occur in meta isomers and aromatic structures with multiple substituted electron donating groups, tend to interfere with repellency (Mason et al., 1989; Clark, 1991a; Clark and Shah, 1991, 1994; Clark et al., 1991; Shah et al., 1991).

Quantitative structure–activity relationships of aromatic compounds and repellency are consistent with earlier qualitative studies. The aversive properties of 14 derivatives of cinnamic acid compounds are characterized by heat of formation (DH(f)), polarizability (XY and YY), and superdelocalizability (Sr). All of these descriptors are electronic (Watkins et al., 1999). These findings generally align with a reanalysis of the quantitative structure–activity relationships of the 117 compounds described above (Clark, 1997). Canonical analysis of the relationship of physicochemical, topological, and electrostatic descriptors and the response shape of the four-parameter fluid intake curve showed that 94% of variance in the response profile could be accounted for by five parameters: polarizability, ES2, ANC, KAPPA2, and CHI2. Polarizability is the relative susceptibility of the electron cloud of a molecule to be distorted by presence of an external electric field. Owing to distortion, an induced electric dipole moment appears. Temporary dipoles induce dipoles in other molecules, resulting in van der



**FIGURE 7.3** The relative reduction of fluid intake for solutions as a function of chemical class, which assumes a benzene parent structure with the nomenclatural taxonomy defined by the principal electron withdrawing group. Fluid intake is the asymptotic minimum intake in one-bottle 6-h drinking trials ( $R_{min}$ ). Strongly aversive solutions (where  $R_{min}$  is not statistically distinguishable from zero) have  $R_{min} \leq 0.2$ . Moderately aversive solutions have  $0.2 < R_{min} \leq 0.4$ , weakly aversive solutions have  $0.4 < R_{min} \leq 0.6$ , and solutions with  $R_{min} > 0.6$  are not aversive at all (not statistically different from water controls). Median  $R_{min}$  (solid bars),  $R_{min}$  25–75th percentile (shaded box),  $R_{min}$  5–95th percentile (capped line), and the range of  $R_{min}$  (open symbols). Adapted from Clark (1997).

Waals intermolecular forces by orienting the temporary and induced dipoles with each other. ES2 is an electrotopological descriptor that describes electronic interactions between molecules. ANC is a partial negative electronic charge descriptor of electrostatic potential that influences molecular interactions. CHI2 and KAPP2 are valence connectivity and shape descriptors that may describe the rigidity of the molecule and accessibility of the molecule to receptor systems. The importance of electronic features of molecules is consistent with studies of TRPA1 channel modulation and activation of cysteine-reactive chemicals. TRP channel activation was found to be more dependent on chemical reactivity relative to molecular shape (Hinman et al., 2006; Macpherson et al., 2007). However, the importance of gaining access to proximity of the TRP channels owing to influences of molecular flexibility and shape still remains to be more fully explored.

### 7.2.5 Responses to Respiratory Stimuli

Changes in carbon dioxide concentration in the nasopharynx region can cause species-specific changes in reflexive breathing in birds (Hiestand and Randall, 1941). However, concentrations of carbon dioxide that are sufficiently high to be irritating to mammals have no effect on blood

pressure, heart rate, tidal volume, breathing frequency, upper airway resistance, or lower airway resistance in geese (*Anser anser*) and chickens. Geese and chickens respond differently than mammals to exposure to sulfur dioxide, but in a similar manner when exposed to ammonia and phenyl diguanide (Callanan et al., 1974; McKeegan et al., 2005).

### 7.2.6 Nasal and Respiratory Irritation and Interaction of Olfaction and Chemesthesis

The TN is important in the perception of odors (Tucker, 1971; Silver and Maruniak, 1981; Keverne et al., 1986). Electrophysiological evidence shows that the TN responds to odors, although it is generally less sensitive than the olfactory nerve (Tucker, 1963). Behavioral assays yield similar results. Pigeons trained to respond to odors fail to respond after olfactory nerve transections. However, odor responding can be reinstated if the odor concentration is increased (Michelsen, 1959; Henton, 1969; Henton et al., 1966). Odor sensitivity of pigeons decreased by 2–4 log units (vapor saturation) after olfactory nerve transection (Walker et al., 1979).

Although olfaction can modulate responding to chemical irritants, it is relatively unimportant (Clark, 1995). In European starlings, avoidance of anthranilate compounds

was partially a consequence of olfactory cues. When the olfactory nerves were transected, avoidance was only mildly diminished. When the ophthalmic branches of the TN were transected, the starlings became insensitive to the aversive properties of the anthranilates (Mason et al., 1989).

### 7.2.7 Behavioral Responses to Irritants

Many aromatic molecules are aversive to birds (Kare, 1961; Mason et al., 1989; Crocker and Perry, 1990; Clark and Shah, 1991, 1993; Crocker et al., 1993). Several lines of evidence suggest that a variety of compounds have intrinsic properties that cause them to be aversive on a purely sensory basis. First, the aversive quality is unlearned; that is, avoidance occurs upon initial contact (Clark and Shah, 1991). Second, there is no evidence that consumption is altered by gastrointestinal feedback; intake of fluid treated with those sensory stimuli is constant over time (Clark and Mason, 1993). Third, unlike mammals, birds seem unable to associate the aversive quality of the stimulus with other chemosensory cues, suggesting that conditioned flavor avoidance learning does not occur (Clark, 1996; Clark and Avery, 2013). Fourth, birds do not habituate to the stimulus; avoidance persists in the absence of reinforcement (Clark and Shah, 1994).

### 7.2.8 Applications

Current interest in chemesthetic function and properties in birds is largely focused in four areas: (1) the evolutionary phylogenetic relationships of receptor mediated perception of noxious stimuli and its consequence to the foraging ecology of birds (Clark, 1998; Tewksbury and Nabhan, 2001); (2) the applicability of using aversive compounds in modulating feeding behavior of birds to develop repellents for prevention of crop damage or otherwise mitigating against damage caused by birds (Mason and Clark, 1997; Clark and Avery, 2013); (3) efforts to gain a better understanding of pathologic pain caused by “debeaking” and promotion of animal welfare in domestic chicken production through better management methods or development of appropriate analgesics (Kuenzel, 2007; Gentle, 2011); and (4) discovery of better analgesics for management of pain in veterinary clinical settings.

## 7.3 OLFACTION

### 7.3.1 Morphology of Olfactory System

Air entering a bird’s nasal cavity passes through a series of mucous-covered, invaginated chambers called nasal conchae. Nasal conchae influence air flow dynamics and direct odors to the caudal-most chamber, which contains the chemically sensitive olfactory epithelium (reviewed in

Roper, 1999; see also: Bang, 1960, 1961, 1963, 1964, 1965, 1966; Bang and Cobb, 1968). The surface of the olfactory epithelium is composed of receptor cells, which detect odorous compounds and occur at the ends of olfactory nerve dendrites. Each receptor cell is surrounded by a cluster of supporting cells and ends in a knob bristling with 6–15 cilia that extend into the lumen. The length of cilia varies by species. Black vultures, for example, have cilia of 40–50  $\mu\text{m}$ , whereas domestic fowl have cilia of 7–10  $\mu\text{m}$  (Shibuya and Tucker, 1967). To gain access to the cilia of receptor cells, odor molecules must diffuse through a mucous membrane. Cilia themselves provide no transport function. Rather, secretions covering cilia provide rapid flow for odor molecules. Olfactory gland secretions must be removed and replaced to maintain diffusion and avoid receptor habituation to odorant molecules. Traction of nearby respiratory cilia facilitates removal of secretions.

The extent of scrolling of caudal conchae correlates with the surface area of olfactory epithelium and the relative size of the olfactory bulb, which is the region of the brain that processes odor input (Bang and Cobb, 1968; Bang, 1971; Bang and Wenzel, 1985; reviewed in Roper, 1999; Hagelin, 2007a). Avian orders with relatively larger olfactory bulbs have lower detection thresholds, indicating they are more sensitive to certain odorous compounds than those with relatively small olfactory bulbs (Clark et al., 1993; Table 7.1, Figure 7.4). Elaborated olfactory systems typically belong to species with demonstrated reliance on odor cues in the field (Stager, 1964; Hutchison and Wenzel, 1980; Hagelin, 2004) and, in some species, correlate positively with the number of olfactory receptor genes (Steiger et al., 2008). Fossil evidence also indicates olfactory bulb size was relatively large early in bird evolution, revealing a previously unrecognized emphasis on smell (Zelenitsky et al., 2011).

Although a larger olfactory bulb size or greater scrolling of receptor epithelium likely indicates greater functional capacity (e.g., more cells and neural circuits; Meisami, 1991), it is important not to dismiss avian species with relatively “unelaborate” olfactory systems (Hagelin, 2007b). Both field and laboratory tests indicate that several taxa with relatively small olfactory bulbs can discriminate between and/or adaptively employ certain odors, such as those related to breeding and nesting (e.g., crested auklets (*Aethia cristatella*) Hagelin et al., 2003; European starlings Clark and Mason, 1985; Gwinner and Berger, 2008; Corsican Blue Tit (*Parus caeruleus ogliastreae*) Petit et al., 2002).


### 7.3.2 Innervation of Olfactory Receptors

Olfactory receptor cells from each nasal cavity transmit information via the olfactory nerve to the olfactory bulb, located in the anterior region of each brain hemisphere. Each olfactory bulb is composed of concentric cell layers.

**TABLE 7.1** Summary of Mean Ratios of Ipsilateral Olfactory Bulb Diameter to Cerebral Hemisphere Diameter and Their Standard Errors (SE) for Several Orders of Birds

Order	N	Ratio	SE	Order	N	Ratio	SE
Anseriformes	4	19.4	1.5	Psittaciformes	2	8.0	1.4
Apodiformes	8	12.3	1.9	Falconiformes	5	17.4	2.6
Apterygiformes	1	34.0	0.0	Charadriiformes	9	16.4	0.9
Caprimulgiformes	3	23.3	0.7	Galliformes	3	14.2	1.4
Columbiformes	2	20.0	1.4	Piciformes	5	11.4	1.3
Cuculiformes	4	19.5	0.6	Passeriformes	25	13.3	0.7
Gruiformes	14	22.2	0.9	Pelecaniformes	4	12.1	1.6
Gaviformes	1	20.0	0.0	Coraciiformes	5	14.5	1.6
Podicipediformes	2	24.5	1.8	Sphenisciformes	1	17.0	0.0
Procellariiformes	10	29.1	1.4				

Sample sizes indicate the number of species (N).  
 Source: Data adapted from Bang and Cobb (1968).



**FIGURE 7.4** Relationship between olfactory detection threshold and relative size of the olfactory bulb for different orders of birds. Adapted from Clark and Shah (1993).

Incoming olfactory nerve fibers constitute the outer layer. Branching nerve terminals penetrate into the adjacent, glomerular layer, where they connect with dendrites of mitral and tufted cells in spherical arborizations called glomeruli. The perikarya of these cells are in the deeper mitral cell layer, where their axons leave to project to many areas of the forebrain.

Like other vertebrates, the olfactory bulbs of birds are bilaterally symmetrical; each is associated with its own (ipsilateral) brain hemisphere. The layering of different cell types within avian olfactory bulbs is qualitatively similar to reptiles, in that well-defined cell layers (like those of mammals) are lacking (Allison, 1953; Andres, 1970). However, there are many interneuron connections in the cell layers

between the mitral and glomerular regions. There are no direct connections between the two (contralateral) olfactory bulbs (Rieke and Wenzel, 1978).

Although birds clearly have olfactory bulbs, they appear to lack an accessory olfactory system (Rieke and Wenzel, 1974, 1978). Both olfactory and accessory olfactory structures commonly occur in other vertebrates. The accessory olfactory system is frequently linked to conspecific scent stimuli that modulate social behavior (e.g., reproduction, aggression). However, there is good evidence for mammals that both the main olfactory and accessory olfactory systems can detect and process overlapping sets of odor stimuli (Keller et al., 2009). Accessory olfactory structures include the vomeronasal organ and accessory olfactory bulb. It is possible that accessory olfactory bulbs in birds occur during early embryonic development only, but are lost later on (Matthes, 1934). This idea, however, has received little scientific attention.

### 7.3.3 Olfactory Neuronal Response

Electrophysiological responses to odor stimuli are taken as definitive evidence of olfactory capacity. These can be recorded from a single “unit” (neuron) or multiunit nerve fibers. Recordings of black vultures indicate that the electro-olfactogram appears primarily during inspiration, which coincides with peak spike activity (Shibuya and Tucker, 1967). Electrophysiological recordings of mammals, amphibians, reptiles, and birds all show similar responses, irrespective of the size of a species’ olfactory apparatus (Tucker, 1965; Shibuya and Tonosaki, 1972).

Single-unit responses from within the olfactory bulb of domestic chickens show widely variable rates of spontaneous firing (mean 4.9 spikes/s, range 0.1–32.4 spikes/s) prior to odor exposure (McKeegan, 2002). Odor stimulation modifies spontaneous firing via excitation or inhibition. Avian firing rates appear to fall in between rates reported for mammals and reptiles (McKeegan, 2002, 2009). Single units of chickens responded to two or more odors and revealed surprising sensitivity to biologically relevant scents associated with captivity (e.g., hydrogen sulfide). Responses to extremely low (<0.5 ppm) stepwise changes in concentration to hydrogen sulfide revealed a level of fine-tuning not previously reported for other vertebrates (McKeegan et al., 2002). Continuous presentation of a stimulus can result in physiological adaptation of both single-unit (McKeegan and Lippens, 2003) and nerve-unit recordings, like mammals. Recovery can be achieved within a few minutes of rest.

Olfactory nerve fibers are unmyelinated, which produces slow conduction velocities of about 1.5 m/s (Macadar et al., 1980). Interestingly, transected olfactory nerves (which experimentally inhibit olfaction) can repair and recover full physiological capacity within 30 days (Tucker et al., 1974). Although healed nerves are scarred and smaller, recordings

and autonomic reflexes to odorants did not differ between controls and nerves that had been cut at least 6 months earlier (Tucker, 1971; Tucker et al., 1974).

Another means of quantifying olfactory neural responses involves calcium imaging (Restrepo et al., 1995). This method uses fluorescence to quantify changes in the flux of calcium ions associated with neural activation (i.e., signal transduction) of a single olfactory receptor neuron (ORN). Jung et al. (2005) tested responses of acutely dissociated ORNs from olfactory epithelium of embryonic domestic chicks. Avian ORNs were placed in Ringer’s solution containing liquid solutions of odorants. The fluorescence patterns, which correspond to increases or decreases in  $\text{Ca}^{2+}$  concentration, were remarkably similar to those of other vertebrates (mammals and fish) that had been tested with the same set of odorants (Jung et al., 2005).

### 7.3.4 Laboratory Detection Thresholds, Discrimination, and Seasonal Change

Physiological responses (e.g., change in respiration or heart rate) to novel odor stimuli have been observed (Wenzel and Sieck, 1972). Habituation to the stimulus under this paradigm, however, is problematic. Operant and classical conditioning paradigms that use positive or negative reinforcement (Michelsen, 1959; Henton et al., 1966; Henton, 1969) are usually poor at determining olfactory thresholds or discrimination (Calvin et al., 1957). However, two process learning paradigms, such as cardiac conditioning, have proven to be a successful technique for detection, discrimination, and threshold testing (Rescorla and Solomon, 1967; Walker et al., 1986; Clark and Mason, 1989; Clark and Smeraski, 1990; Clark, 1991a; Clark et al., 1993). During cardiac conditioning, an odor (the conditional stimulus) is paired with an aversive experience, such as a shock (the unconditional stimulus). Heart rate is compared before and after stimulus presentation during training until a level of cardiac acceleration is reliably achieved, indicating a bird has learned to associate the odor in anticipation of a shock. Thereafter, tests of detection or odor discrimination can proceed. Most birds tested with this paradigm have shown olfactory capabilities comparable to mammals (Davis, 1973). Even passerines, with the least developed olfactory system, demonstrate behavioral responsiveness to odors (Clark and Mason, 1987; Clark and Smeraski, 1990; Clark, 1991a; Clark et al., 1993) (Table 7.2).

European starlings offer an interesting case study of olfactory structure, function, and seasonality. Male starlings incorporate green plants that are rich in aromatic volatiles into nests, some of which act as a fumigant against parasites and pathogens (Clark and Mason, 1985, 1987, 1988; Clark, 1991b; Gwinner, 1997; Gwinner et al., 2000; Gwinner and Berger, 2005). Starlings are most sensitive to, and can discriminate between, plant odors during spring only, rather than

TABLE 7.2 Summary of Selected Behavioral Olfactory Threshold Data for Different Species of Birds

Species	Ratio <sup>1</sup>	Stimulus	Threshold (ppm)		Source
			Min	Max	
Rock dove ( <i>Columba livia</i> )	18.0	<i>n</i> -Amyl acetate	0.31	29.8	Henton (1969), Henton et al. (1966), Walker et al. (1979), Walker et al. (1986)
		Benzaldehyde	0.47	00.75	Walker et al. (1986)
		Butanethiol	13,820	–	Snyder and Peterson (1979)
		Butanol	0.17	–	Walker et al. (1986)
		<i>n</i> -Butyl acetate	0.11	2.59	Henton (1969), Walker et al. (1986)
		Butyric acid	2.59	–	Henton (1969)
		Ethanethiol	10,080	–	Snyder and Peterson (1979)
		Heptane	0.29	0.38	Stattelman et al. (1975)
		Hexane	1.53	2.98	Stattelman et al. (1975)
		Pentane	16.45	20.76	Stattelman et al. (1975)
Chicken ( <i>Gallus gallus</i> )	15.0	Heptane	0.31	0.57	Stattelman et al. (1975)
		Hexane	0.64	1.00	Stattelman et al. (1975)
		Pentane	1.58	2.22	Stattelman et al. (1975)
Northern bobwhite ( <i>Colinus virginianus</i> )	–	Heptane	2.14	3.49	Stattelman et al. (1975)
		Hexane	3.15	4.02	Stattelman et al. (1975)
		Pentane	7.18	10.92	Stattelman et al. (1975)
Black-billed magpie ( <i>Pica pica</i> )	–	Butanethiol	13,416	–	Snyder and Peterson (1979)
		Ethanethiol	8400	–	Snyder and Peterson (1979)
European starling ( <i>Sturnus vulgaris</i> )	9.7	Cyclohexane	2.50	–	Clark and Smeraski (1990)
Cedar waxwing ( <i>Bombycilla cedrorum</i> )	–	Cyclohexane	6.80	86.46	Clark (1991a)
Tree swallow ( <i>Tachycineta bicolor</i> )	15.0	Cyclohexane	73.42	–	Clark (1991a)
Brown-headed cowbird ( <i>Molothrus ater</i> )	7.0	Ethyl butyrate	0.76	–	Clark and Mason (1989)
Catbird ( <i>Dumetella carolinensis</i> )	–	Cyclohexane	35.14	–	Clark et al. (1993)
Eastern phoebe ( <i>Sayornis phoebe</i> )	–	Cyclohexane	35.61	–	Clark et al. (1993)
European goldfinch ( <i>Carduelis carduelis</i> )	–	Cyclohexane	13.05	–	Clark et al. (1993)
Great tit ( <i>Parus major</i> )	–	Cyclohexane	34.10	–	Clark et al. (1993)
Black-capped chickadee ( <i>Parus atricapillus</i> )	3.0	Cyclohexane	59.95	–	Henton (1969)

<sup>1</sup>The ratio of the longest axis of the olfactory bulb to that of the ipsilateral cerebral hemisphere.

in summer and fall. Spring is coincident with nest building and suggests a hormonal influence (Clark and Smeraski, 1990).

Birds treated with testosterone (T), a hormone that enlarges song-learning nuclei of the brain and alters behavior, exhibited enlarged olfactory bulbs year-round, indicating a proximate effect on bulb structure. However, perception of plant odor in T-implanted males was greatest during spring only, indicating that perception was independent of T-treatment and olfactory bulb volume. One hypothesized but untested mechanism is that an increase in receptor cell density in starling olfactory epithelium occurs in spring (DeGroof et al., 2010).

### 7.3.5 Development

Volatile compounds diffuse through avian eggshell (Rahn et al., 1979), providing an opportunity for odor exposure within the egg (Tolhurst and Vince, 1976; Sneddon et al., 1998). Many vertebrates, including birds, detect and learn chemical information as embryos (e.g., humans: Schaal et al., 2000; Mennella et al., 2001; other mammals: Hepper, 1988; Bilko et al., 1994; amphibians: Mathis et al., 2008; birds: Porter and Picard, 1998; Bertin et al., 2012). Early exposure can cause changes in neuroanatomy, which alters chemosensory perception in a way that can adaptively shape responses later in life (e.g., to food, mates, etc.) (Todrank et al., 2011).

Studies of domestic chickens, the avian model for development, indicate that odor detection can occur before or after young pierce the egg's air sac and begin breathing air (Tolhurst and Vince, 1976; Bertin et al., 2012; Hagelin et al., 2013). ORNs are functional 6 days prior to air-breathing (on embryonic developmental day 13; Lalloué et al., 2003), when nasal passages are full of amniotic fluid. Embryos at this stage swallow frequently, facilitating fluid movement, similar to mammals in utero (Sneddon et al., 1998). Airbreathing begins approximately 2 days prior to hatching, on embryonic developmental day 19 (Tolhurst and Vince, 1976).

The magnitude of embryonic response varies relative to stimulus concentration and timing of exposure (Bertin et al., 2010). Later developmental stages show relatively greater responses to odors (Gomez and Celli, 2008; Bertin et al., 2012). Detectable stimuli include artificial odors (Sneddon et al., 1998), as well as naturally occurring scents, such as nest materials (Gwinner and Berger, 2008), food-related odors (Burne and Rogers, 1999; Cunningham and Nevitt, 2011), and compounds found in plumage scent of at least one alcid species (Hagelin et al., 2013).

### 7.3.6 Field Studies and Behavioral Ecology

Like other vertebrates, birds detect and respond adaptively to odors (reviewed in Roper, 1999; Hagelin, 2007a;

Balthazart and Taziaux, 2009; Caro and Balthazart, 2010). Hagelin (2007a) made a distinction between environmentally derived odors (e.g., food, predators) and those produced by birds themselves (e.g., body odors, fecal odor, preen gland secretions). The latter can have social and reproductive implications. This section considers examples of adaptive olfactory responses to environmental odors as well as bird-derived scents.

The use of olfactory cues for locating food has been documented for numerous species, such as procellariids, vultures, corvids, hummingbirds, honeyguides, parrots, and kiwis (Roper, 1999). Turkey vultures (*Cathartes aura*), for example, are attracted to ethyl-mercaptan, a volatile associated with decomposed carcasses (Stager, 1964, 1967), and locate food without visual cues (Houston, 1986). Procellariiforms also forage over considerable distances. Black-footed albatrosses (*Diomedea nigripes*) respond to bacon grease over 31 km away (20 miles; Miller, 1942), whereas Leach's storm petrel (*Oceanodroma leucorhoa*) home to scent targets at a distance of 1–12 km (Clark and Shah, 1992). Some procellariiformes also respond to a compound that is correlated with prey called dimethyl-sulfide (DMS) (Nevitt et al., 1995). DMS smells like rotten seaweed and results from the breakdown of metabolic products of marine algae (phytoplankton). Petrels, however, do not feed on phytoplankton. Rather, DMS concentrates in locations where a bird's prey (zooplankton, such as krill) is actively grazing on phytoplankton. Grazing by zooplankton lyses phytoplankton cells and thereby creates a DMS odor plume, which some birds follow to locate food (Nevitt, 2011).

With regard to predators, the scent of urine and/or feces has an aversive effect on some avian species (blue tits (*Cyanistes caeruleus*), Amo et al., 2008; house finches (*Carpodacus mexicanus*), Roth et al., 2008; red junglefowl (*Gallus gallus*), Zidar and Løvlie, 2012), but not all (eastern blue birds (*Sialia sialis*), Godard et al., 2007; house wren (*Troglodytes aedon*), Johnson et al., 2011). Application of predator odor can also deter breeding ducks and songbirds (Eicholz et al., 2012; Forsman et al., 2013). Responses appear to be innate rather than learned (Amo et al., 2011b), although sleeping birds are unreactive (Amo et al., 2011a).

Odors are also germane to avian orientation and navigation (reviewed in Wallraff, 2005; Gagliardo, 2013). Homing pigeons, for example, exhibit larger olfactory bulbs than nonhoming breeds (Rehkämper et al., 1988, 2008). Investigators have also altered pigeon homing behavior via experimental disruption of the olfactory system. Manipulations include olfactory nerve transection (Papi et al., 1971; Gagliardo et al., 2006, 2009), anesthesia of olfactory mucosa (Wallraff, 1988), ablating the central piriform cortex of the brain (Papi and Casini, 1990), and nostril plugging. The last of these manipulations indicates that pigeons rely more on their right nostril for olfactory information (Gagliardo et al., 2007, 2011). ZENK, an

immediate early gene expressed in olfactory neurons, also implicates the use of olfaction during the process of homing (Patzke et al., 2010).

Emerging evidence for passerine species further supports olfaction during migration. For example, adult gray catbirds (*Dumetella carolinensis*) rendered temporarily anosmic (by washing the olfactory tissues with zinc sulfate) oriented differently from adult controls but similarly to juvenile birds, which were migrating for the first time and therefore unable to navigate (Holland et al., 2009). With regard to cellular mechanisms, black-headed buntings (*Emberiza melanocephala*) increase activation of olfactory tissues (as measured by c-fos immunoreactivity) during migration. These birds exhibit a seasonally enhanced emphasis on olfaction while migrating, compared to visual systems (Rastogi et al., 2011).

Many birds produce a variety of odorous compounds (Table 7.3; reviewed in Campagna et al., 2011). For example, a seabird colony, with its dense numbers of birds, burrows, and feces, makes for a potent chemosensory experience. Pioneering work by Grubb (1974) on Leach's storm petrel showed differential return rates to nest sites after surgical manipulation, indicative of olfactory-based homing: 91% for controls, 74% for sham surgery, and 0% for olfactory nerve section. Several petrel species have since been shown to discriminate between the odor of their own nest and conspecific burrows (Mínguez, 1997; De León et al., 2003; Bonadonna et al., 2003a,b). Attraction to home nest

odor is also reported for passerines (Caspers and Krause, 2010; Krause and Caspers, 2012).

Avian chemical substances are linked with a variety of social contexts (reviewed in Hagelin, 2007a; Hagelin and Jones, 2007; Balthazart and Taziaux, 2009; Caro and Balthazart, 2010). Uropygial gland secretions, for example, show some level of hormonal control and exhibit individual, sex, and age-specific patterns (e.g., Procellariiformes: Mardon et al., 2010, 2011; Anseriformes: Kolattukudy et al., 1987; Galliformes: Karlsson et al., 2010; passerines: Whittaker et al., 2010; Whelan et al., 2010; Shaw et al., 2011; Amo et al., 2012a). Pioneering work by Balthazart and Schoffeniels (1979) indicated male mallards decreased social displays and sexual behavior toward females when their olfactory nerves were sectioned, suggesting that intact olfactory system is critical to courtship and mating. Crested auklets produce a seasonally elevated scent associated with a stereotyped behavior that focuses on the scented region of the body (the nape). Auklets are attracted to natural feather odor, a chemical cocktail of odor compounds, and scented decoys, which suggests odor has a social function (Hagelin et al., 2003; Jones et al., 2004; Hagelin, 2007a). Odorous compounds of crested auklets can also negatively impact ectoparasites in experimental tests (Douglas, 2008, 2013).

Procellariiform seabirds show a surprising level of body odor discrimination, in that they are attracted to mate odors and avoid self-odor (Antarctic petrel (*Pachyptila desolata*) Bonadonna and Nevitt, 2004; blue petrels (*Halobaena caerulea*), Mardon and Bonadonna, 2009). Furthermore, preference for the odor of unrelated individuals over those of kin was recently discovered (European storm petrel (*Hydrobates pelagicus*), Bonadonna and Sanz-Aguilar, 2012). Such results suggest that body odors could provide a mechanism for inbreeding avoidance, known as self-referent phenotype matching (Mateo and Johnston, 2000). This may be particularly important in petrels which are a long-lived philopatric species that mates for life. Petrels are also likely to encounter kin on their natal breeding grounds that they have never met before (Bonadonna and Nevitt, 2004; Bonadonna and Sanz-Aguilar, 2012). Recent evidence for passerines suggests that conspecific odor may provide relevant social information. Bird responses to scent correlated with social rank (house finch, Amo et al., 2012b), sex (European starling, Amo et al., 2012a), and body size (dark-eyed junco (*Junco hyemalis*), Whittaker et al., 2011).

### 7.3.7 Summary

Every bird tested has exhibited a functional sense of smell (Bang and Wenzel, 1985). The extent of olfactory development also is on par with that found in mammals. However, ornithologists have largely overlooked the role of olfaction in avian biology. Many birds adaptively employ

**TABLE 7.3** Some Avian Orders Considered To Be Very Odorous by Ornithologists

Order	Common Name	Number of Species <sup>1</sup>
Procellariiformes	Petrels, shearwaters, diving petrels	16
Ciconiiformes	Herons, storks, new world vultures	12
Anseriformes	Ducks, geese, swans, screamers	49
Charadriiformes	Sandpipers, gulls, auks	23
Psittaciformes	Parrots	14
Cuculiformes	Cuckoos	16
Coraciiformes	Kingfishers, rollers, hoopoes, woodhoopoes	14
Piciformes	Woodpeckers, barbets, toucans	33
Passeriformes	Grackles, starlings, ravens, finches, honeycreepers	46

<sup>1</sup>Data compiled from Weldon and Rappole, 1997.

environmental odors; they also produce and respond to conspecific scents. Although passerines have a relatively poorly developed olfactory anatomy, they nonetheless show some degree of olfactory acuity. Other species, such as procellariiformes, have olfactory systems that are acutely sensitive to odor cues and capable of a surprisingly detailed level of conspecific odor discrimination. Given the broad range of contexts that implicate avian olfaction, future interdisciplinary research that compares olfactory mechanisms in birds to better-known vertebrate systems, such as mammals and fish, holds exciting promise.

## 7.4 GUSTATION

### 7.4.1 Taste Receptors

Relative to other vertebrates, birds have fewer taste receptors and taste receptor genes (Berkhoudt, 1985; Shi and Zhang, 2005) (Table 7.4). Notwithstanding these observations, birds have a well-developed system for gustation with functional significance for their behavior, ecology, and evolution. Taste receptors are located in taste buds throughout the oral cavity. The greatest concentration of avian taste receptors is found around salivary glands in the soft epithelium of the palate, the posterior tongue, and the oropharynx (Bath, 1906; Lindenmaier and Kare, 1959; Saito,

1966; Ganchrow and Ganchrow, 1985). Afferent taste signals in birds are carried in the glossopharyngeal nerve (cranial nerve IX; Duncan, 1960). The glossopharyngeal nerve innervates the posterior buccal and pharyngeal areas (Kare and Mason, 1986). Unlike mammals, the facial nerve (VII) does not innervate the avian tongue (Wenzel, 1973). Rather, glossopharyngeal afferents in birds enter the medulla and join fibers from the facial (including chorda tympani) and vagus nerves (X) to form a well-developed *fasciculus solitarius* (Lindenmaier and Kare, 1959). The chorda tympani innervates taste buds adjacent to the anterior mandibular salivary glands, situated in the buccal epithelium of the lower jaw (Kare and Mason, 1986).

### 7.4.2 Response to Sweet

Birds have a well-developed sense of taste that generally corresponds to their feeding habits. Frugivorous and omnivorous birds tend to perceive and prefer sweet more so than species in other foraging guilds. For example, European starlings prefer 0.5–5% D-fructose solutions (w/v) to distilled water (Espaillat and Mason, 1990). Sugar detection thresholds of cockatiels (*Nymphicus hollandicus*) is 0.36 M sucrose, 0.40 M fructose and 0.16 M glucose (Matson et al., 2000, 2001). The sugar detection thresholds of broad-billed hummingbirds (*Cyanthus latirostris*) is between 1.31 and 1.54 mM sucrose, 0.87–1.31 mM fructose, 1.54–1.75 mM glucose and 1.75–3.5 mM of a 1:1 mixture of fructose and glucose (Medina-Tapia et al., 2012). Interestingly, the sweet taste receptor gene *Tas1r2* is absent in all bird genomes sequenced thus far, irrespective of their diet (Zhao and Zhang, 2012), suggesting that additional avian receptors may exist for sweet.

The order of preference among nectarivorous passerines is sucrose = glucose + fructose = fructose > glucose > xylose (Lotz and Nicolson, 1996). Lesser double-collared sunbirds (*Nectarinia chalybea*) and Cape sugarbirds (*Promerops cafer*) absorb sucrose, glucose, and fructose from ingested food at nearly 100% efficiency, but xylose was excreted (Lotz and Nicolson, 1996; Jackson et al., 1998a,b). Although nectar composition and concentration are often considered independently, these characteristics may have a synergistic effect on the sugar preferences of nectar-feeding birds (Schondube and Martinez del Rio, 2003).

Sugar preferences among nectarivorous and frugivorous birds are concentration-dependent. Although nectarivorous birds in Africa prefer sucrose when offered a choice of 0.25 M solutions of glucose, fructose, and sucrose, no preference among these sugars was observed when their concentration was increased to 0.73 M; the dietary choices in these species indicate the birds had either reached a limit where they had sufficient energy intake or they were affected by

**TABLE 7.4** Abundance of Taste Buds among Vertebrate Species<sup>1</sup>

Species	Taste Buds	Source
Domestic chick (day-old)	5–12	Lindenmaier and Kare (1959)
Domestic chicken (3 months)	24	Lindenmaier and Kare (1959)
Blue tit	24	Gentle (1975)
Bullfinch	41–42	Duncan (1960)
Pigeon	59	Moore and Elliot (1946)
Japanese quail	62	Warner et al. (1967)
European starling	200	Bath (1906)
Parrot	300–400	Bath (1906)
Domestic cat (juvenile)	473	Elliot (1937)
Lizard	550	Schwenk (1985)
Bat	800	Moncrieff (1946)
Domestic cat (adult)	2755	Robinson and Winkles (1990)
Human	6974	Miller and Reedy (1990)
Rabbit	17,000	Moncrieff (1946)
Pig	19,904	Chamorro et al. (1993)
Ox	35,000	Moncrieff (1946)
Catfish	100,000	Hyman (1942)

<sup>1</sup>Modified from Kare and Mason (1986) and Mason and Clark (2000).

post-ingestion constraints (Downs and Perrin, 1996; Downs, 1997). House finches demonstrated no preference for equicaloric, 2% solutions of hexoses (1:1 mixture of fructose and glucose) and sucrose, and strong preference manifest for hexoses but not sucrose at 4, 6, and 10% concentrations; energetics, rather than sucrase deficiency, may determine finches' sugar preferences (Avery et al., 1999).

Studies of unrelated, nectarivorous birds (including a generalist, nonpasserine nectarivore) have demonstrated a distinct switch from hexose preference at low concentrations to sucrose preference at higher concentrations (Lotz and Schondube, 2006; Fleming et al., 2008; Brown et al., 2010a,c). Sucrose preference at higher concentrations may possibly be explained by taste perception due to differences in solution osmolality or a degree of imprinting due to experience with natural nectar compositions. Village weavers (i.e., generalist passerine nectarivores; *Ploceus cucullatus*) preferred hexose solutions at 5% and 10% sucrose equivalents (SE), yet no sugar preference was observed at 15, 20, and 25% SE (Odendaal et al., 2010). In contrast, dark-capped bulbuls (*Pycnonotus tricolor*), an opportunistic nectarivore, significantly preferred hexose solutions, irrespective of concentration (5–25%), when given a choice between equicaloric hexose and sucrose solutions (Brown et al., 2010b). Interestingly, malachite sunbirds (*Nectarinia famosa*) demonstrated either sucrose preference, no preference, or hexose preference when offered equimolar, equiweight, or equicaloric paired solutions of sucrose and hexose, respectively (Brown et al., 2008).

The bananaquit (*Coereba flaveola*) strongly prefers the most concentrated sucrose solution when the lowest concentration ranged from 276 to 522 mM. From 522 to 1120 mM sucrose concentrations, bananaquits adjust their volumetric food intake to maintain constant energy intake. At a sucrose concentration of 276 mM, however, bananaquits did not maintain their rate of energy intake by increasing food consumption (Mata and Bosque, 2004). Although nectarivorous birds generally prefer concentrated over dilute sugar solutions, the concentration difference that they can discriminate is smaller at low concentrations relative to high concentrations; this pattern may be a consequence of the functional form of intake responses that often results in decelerating sugar intakes with increasing sugar concentration (Martinez del Rio et al. 2001; Leseigneur and Nicolson, 2009). With regard to gender-specific food intake among nectarivorous birds, males take longer to digest than females when fed on sucrose-rich nectars as opposed to hexose-rich nectars; therefore, they can allow themselves a relatively lower digestive capacity (Markman et al., 2006). The digestive transit rates of Cape white-eyes (*Zosterops virens*) fed artificial fruit were faster for glucose- than sucrose-based diets, irrespective of concentration; increased food intake with decreasing glucose concentration and no significant differences in food intake with differing sucrose concentrations were observed (Wellmann

and Downs, 2009). Indeed, nectar ingestion rate is determined by viscosity, and total food intake is primarily modulated by sugar concentration (Köhler et al., 2010).

Sugar preference and selection among nectarivorous and frugivorous birds are likely to have coevolutionary effects on flowering and fruit-bearing plants. Among 58 wild fruits studied in Hong Kong, all fruit species contained glucose, all but one contained fructose, and only 11 species contained sucrose; birds are known to eat 29 of these species without detectable sucrose and four with sucrose (Ko, 1996). From a comparative analysis of glucose, fructose, and sucrose in the nectar and fruit juice of 525 tropical and subtropical plant species, passerine nectars and fruits had low sucrose and high hexose content, respectively; the nectar of hummingbird flowers had very high sucrose content; microchiroptera nectars showed hexose richness and microchiropteran fruits had a sucrose content similar to passerine fruits; and megachiroptera nectars and fruits were sucrose-rich (Baker et al., 1998). The dichotomy between sucrose-rich nectars in hummingbird-pollinated plants and predominantly hexose-rich nectars in sunbird-pollinated plants appears to have little to do with bird physiology and may rather reflect patterns of nectar secretion or plant physiology and opportunist nectar feeders (Nicolson and Fleming, 2003; Fleming et al., 2004).

The hummingbird-passerine dichotomy was strongly emphasized until the discovery of South African plants with sucrose-dominant nectars, which are pollinated by passerines that demonstrate sucrose digestion and preference (Lotz and Schondube, 2006). Flowers adapted for specialized passerine nectarivores have nectar similar to that of hummingbird flowers in terms of volume (approx. 10–30 mL), concentration (15–25% w/w) and sucrose content (40–60% of total sugar). In contrast, flowers adapted to generalized bird pollinators are characterized by large volumes (approximately 40–100 mL) of extremely dilute (8–12%) nectar with minimal sucrose (0–5%; Johnson and Nicolson, 2008).

Rufous hummingbirds (*Selasphorus rufus*) preferred 50% sucrose to higher and lower concentrations, and they could distinguish solutions differing by only 1% sucrose (Blem et al., 2000). Sucrase activity is 10 times higher in hummingbirds than in passerines (Schondube and Martinez del Rio, 2004). Neither sex nor temperature affected sugar preferences among green-backed firecrown hummingbirds (*Sephanoides sephanioides*; Chalcoff et al., 2008). Patterns of hummingbird sugar preference can be affected by different mechanisms, both pre- and post-ingestive. At low concentrations, gustatory thresholds may play an important role in sugar selection. At intermediate and high concentrations, however, sugar selection can be explained by sugar assimilation rates and velocity of food processing generated by osmotic constraints (Medina-Tapia et al., 2012).

Species belonging to the Sturnidae–Muscicapidae lineage do not express intestinal sucrase, despite having generalist diets comprising fruits with sugars of diverse kinds (Gatica et al., 2006). Members of the Sturnidae–Muscicapidae lineage are intolerant of solutions or fruit above 11–15% sucrose (Brown et al., 2012). Considering the phylogenetic constraint hypothesis for sucrose digestion in the Muscicapidae superfamily, the lack of sucrase activity is a shared, derived character only for the Cinclidae–Sturnidae–Turdinae lineage (Gatica et al., 2006).

Within an experimental meal with varying sucrose concentration, captive whitebellied sunbirds (*Cinnyris talatala*) demonstrated a measurable increase in feeding frequency and food intake within 10 min after a decrease in sucrose concentration (Köhler et al., 2008). Similarly, Knysna turacos (*Tauraco corythaix*) preferred an artificial sucrose diet to an equicaloric glucose diet at low concentrations, whereas purple-crested turacos (*Gallirex porphyreolophus*) showed no preference for either diet. Both turacos species preferred a sucrose diet to an equimolar glucose diet at low concentrations. At high concentrations, neither species showed a preference for either equicaloric or equimolar diets; thus, energy requirements influence food preferences more than sugar type and birds will select fruit that is higher in energy irrespective of sugar type (Wilson and Downs, 2011).

### 7.4.3 Response to Salt

A comparison of the sodium chloride rejection thresholds among 58 bird species illustrated rejection thresholds ranging from 0.35% NaCl in a parrot to 37.5% NaCl in the pine siskin (*Carduelis pinus*; Rensch and Neunzig, 1925). Red-winged blackbirds (*Agelaius phoeniceus*) and European starlings preferred 0.1–1% NaCl solutions (w/v) to distilled water (Espaillat and Mason, 1990). The salt detection threshold of cockatiels is 0.16M NaCl (Matson et al., 2000) and 0.16M potassium chloride (Matson et al., 2001). With regard to the mechanism of salt perception, sodium in the oral cavity can cross the taste sensory cell membrane through the epithelial Na<sup>+</sup> channel (ENaC), thus triggering an action potential (Roura et al., 2012). Pigeons (*C. livia domestica*) learned to discriminate a safe 0.06M NaCl solution and a toxic equimolar LiCl solution. Because the pigeons avoided the LiCl solution within a short presentation period of 5 minutes, it is unlikely that the birds were using an interoceptive stimulus of faint, postingestive malaise as a conditioned cue; thus, the pigeons' discrimination performance between the two chloride solutions was attributed to gustation (Nakajima and Onimaru, 2006).

### 7.4.4 Response to Sour

Sourness is related to the acidity of food, which is often caused by bacterial fermentation and typically evokes a

rejection response. With regard to the mechanism of sour perception, the receptors for sour taste are thought to be transmembrane channels that are selective for hydrogen ions (Roura et al., 2012). Red-winged blackbirds and female starlings preferred distilled water to 0.01–0.1 M citric acid solutions (Espaillat and Mason, 1990). For the purpose of investigating sour detection thresholds, Matson et al. (2000) defined sourness as a pH, and they achieved sourness by varying the pH of a 0.05 M citrate buffer system. The sour detection threshold of cockatiels is pH 5.5 citric acid.

### 7.4.5 Response to Bitter

Bitter taste perception likely evolved as a protective mechanism against the ingestion of harmful compounds in food (Davis et al., 2010). Red-winged blackbirds and European starlings preferred distilled water to 0.5–5% tannic acid solutions (w/v; Espaillat and Mason, 1990). The bitter detection thresholds of cockatiels is 100 μM quinine, 1000 μM gramine, 500 μM hydrolysable tannin and 10,000 μM condensed tannin (Matson et al., 2004). Compared with pigs, chickens showed a lower sensitivity to glucosinolates (i.e., bitter plant metabolites); compared to ruminants, however, chickens showed a higher aversion to glucosinolates (Roura et al., 2012).

Bitter detection thresholds indicate that a birds' rejection of quinine occurs at lower concentrations than phytophagic mammals (Matson et al., 2004). White Leghorn and Rhode Island Red chickens were able to detect 2.0 mM quinine hydrochloride; broiler chickens detected 0.5 mM quinine hydrochloride (Kudo et al., 2010). Domestic chicks (14 days old) can discriminate between an untreated diet and a diet treated with 0.2% quinine hydrochloride (Ueda and Kainou, 2005).

Johnson et al. (2006) explored the functional significance of the phenolic compounds that impart a dark brown color to the nectar of the South African succulent shrub, *Aloe vryheidensis*. Dark-capped bulbuls were more likely to probe model flowers containing dark nectar than those containing clear nectar, suggesting a potential signaling function of dark nectar. The main effect of the phenolics, however, appears to be repellency of “unwanted” nectarivores that find their bitter taste unpalatable. Nectar-feeding honey bees and sunbirds are morphologically mismatched for pollinating *A. vryheidensis* flowers and strongly reject its nectar. Thus, the dark phenolic component of the nectar appears to function as a floral filter by attracting some animals visually and deterring others by its taste (Johnson et al., 2006).

The taste receptor type 2 (Tas2r) gene family encodes the chemoreceptors that are directly responsible for the detection of bitter compounds. The Tas2r cluster encodes up to 18 functional bitter taste receptors in the white-throated sparrow (*Zonotrichia albicollis*; Davis et al., 2010). Although

the tens to hundreds of taste buds observed among birds pales in comparison to the hundreds to thousands of taste buds found in other vertebrates (Table 7.4), this relative deficit does not preclude birds from detecting bitter compounds as effectively as those species with more taste buds. Future biochemical and genetic studies will be needed to identify the natural ligands for avian Tas2r gene clusters, and the intra- and inter-specific differences in these genes with variation in bitter taste perception (Davis et al., 2010).

#### 7.4.6 Response to Umami

Male starlings preferred 0.7–1% L-alanine solutions to distilled water (Espaillat and Mason, 1990). The T1R1 umami receptor gene and the T1R3 sweet/umami receptor gene have been identified in chickens (Shi and Zhang, 2005). Moreover, the expression of T1R1 has been reported in hypothalamus, liver, and abdominal fat (Byerly et al., 2010). Thus, avian taste receptors and umami receptor genes may be involved in the orchestration of postingestive and metabolic events (Roura et al., 2012). Further research is needed to comparatively investigate avian feeding responses to umami tastants.

#### 7.4.7 Response to Calcium

Calcium-deprived chickens preferred calcium-rich diets when offered a choice (Wood-Gush and Kare, 1966; Hughes and Wood-Gush, 1971). Similarly, consumption of supplementary calcium was inversely related to chicken's dietary calcium content (Taher et al., 1984). Further research is needed to distinguish the behavioral responses of birds to calcium as a tastant (i.e., sensory cue) versus the pre- and postingestive attributes of calcium-rich supplements. Although it is clear that animals can detect calcium in micromolar or low millimolar concentrations, it is less clear what they detect or how they detect it (Tordoff, 2001). The notion that calcium is a distinct taste quality is an anathema to many psychophysicists, who argue that there are very few basic taste qualities (sweet, sour, salty, bitter, and umami). To them, calcium taste is a complex of basic tastes, such as bitterness, sourness, and saltiness (Tordoff, 2001).


Calcium taste varies with both the form and the concentration of salt tested, but it nearly always includes sour and bitter components (Tordoff, 2001). The extracellular calcium-sensing receptor (CaR) is a multimodal sensor for several key nutrients, notably  $\text{Ca}^{2+}$  and L-amino acids, and is expressed abundantly throughout the gastrointestinal tract in humans (Conigrave and Brown, 2006). Although the T1r3 receptor gene in mice (Tordoff et al., 2008) and the CaR have been identified as calcium sensors, it is yet uncertain if they mediate calcium appetite or taste (Roura et al., 2012) in birds.

#### 7.4.8 Taste Behavior and Applications

Deterrents based merely on offensive flavors are not likely to be effective in the absence of aversive postingestive effects (Provenza, 1995). In this context, flavor is the perceptual integration of chemesthetic, olfactory, and gustatory stimuli. Red-winged blackbirds conditioned with sodium chloride paired with an intraperitoneal injection of a gastrointestinal toxin (lithium chloride) or a free choice of a postingestive, cathartic purgative (anthraquinone) or a postingestive, cholinesterase inhibitor (methiocarb) subsequently avoided the flavor (NaCl; Figure 7.5) and color of food experienced during conditioning. In contrast, blackbirds conditioned with sodium chloride paired with an intraperitoneal injection of an opioid antagonist (i.e., chemesthetic; naloxone hydrochloride) or a free choice of a preingestive, trigeminal irritant (methyl anthranilate) subsequently avoided only the color (not flavor; Figure 7.5) of food experienced during conditioning. Thus, red-winged blackbirds reliably integrate gustatory (and visual) experience with postingestive consequences to procure nutrients and avoid toxins (Werner and Provenza, 2011).

Avian taste behavior has been investigated in context of agricultural production, chemical defenses of insects and plants, coevolution in predator-prey and pollination systems, chemical ecology, conservation biology, and comparative physiology and taxonomy. For example, although avian feeding responses to secondary metabolites are species-specific (Saxton et al., 2011; Rios et al., 2012), increased sugar concentrations (not decreasing acid concentrations) are a functional cue for the onset of bird damage to ripening grapes (Saxton et al., 2009). Although increased sucrose content may deter sucrose-deficient birds from damaging commercial fruit (Brugger and Nelms, 1991), increased sucrose may also lead to increased crop damage by other species obligated to consume more of the less-digestible fruit to meet their energy requirements (Lane, 1997). This compensatory feeding hypothesis notwithstanding, McWhorter and Martinez del Rio (2000) observed a physiological constraint on sugar consumption among nectarivorous hummingbirds; the rate of intestinal sucrose hydrolysis can limit sugar assimilation and reduce sucrose preference. Indeed, the intake responses of nectar-feeding birds manifest from the integration of a behavioral response with the physiological processes that shape it (Martinez del Rio et al., 2001).

Several tastants have been used to condition aversions among birds associated with agricultural production. The risk of accidental poisoning of birds may be reduced by adding an aversive tastant (e.g., D-pulegone, quinine hydrochloride) to granular pesticides (Mastrota and Mench, 1995; Clapperton et al., 2012). Garlic oil was identified as an effective chemical repellent for European starlings (Hile et al., 2004) and quinine sulfate (bitterant) was used to



**FIGURE 7.5** Mean consumption ( $\pm 2$  SEM) of sodium chloride and citric acid subsequent to NaCl conditioning with: an intraperitoneal injection of a gastrointestinal toxin (lithium chloride), or a free-choice of a postingestive, cathartic purgative (anthraquinone) or a postingestive, cholinesterase inhibitor (methiocarb); or an intraperitoneal injection of an opioid antagonist (naloxone hydrochloride) or a free-choice of a pre-ingestive, trigeminal irritant (methyl anthranilate) in red-winged blackbirds (*Agelaius phoeniceus*). From Werner and Provenza (2011), baseline preference data from Werner *et al.* (2008); with permission.

condition taste aversions and thus reduce destructive feather pecking among laying hens (Harlander-Matauschek *et al.*, 2009, 2010).

Relative preference for specific tastants has been used to enhance feeding for poultry production. The preference of chickens for oily diets (i.e., long-chain versus medium-chain triacylglycerol) is mediated by gustation (Furuse *et al.*, 1996; Mabayo *et al.*, 1996), not satiety (Vermant *et al.*, 1997). In contrast, the avoidance of a saponin-rich diet is not mediated by taste in domestic chicks (Ueda and Shigemizu, 2001); rather, crop distension causes decreased feed intake associated with tea saponin (Ueda *et al.*, 2002).

Domestic chicks can use unpalatable taste (e.g., quinine) to adapt their visual foraging decisions (Rowe and Skelhorn, 2005; Skelhorn *et al.*, 2008). Moreover, European starlings and domestic chicks can learn to use bitter taste cues to regulate consumption of toxic prey (Skelhorn and Rowe, 2010; Barnett *et al.*, 2011). Similarly, red-winged blackbirds use affective processes (flavor-feedback relationships) to shift preference for both novel and familiar flavors (Werner *et al.*, 2008).

#### 7.4.9 Summary

The conventional notion regarding the “limited ability of birds to taste” (Kassarov, 2001) was shaped by a historic paradigm of taste research (i.e., elementary structure and

function). Avian taste perception is currently investigated in context of ontogenetic and phylogenetic relationships within ever-changing environments. Birds use taste cues to select nutrients and avoid toxins; thereby, they affect the distribution, diversity, and coevolution of their prey. Thus, taste cues and postingestive consequences have behavioral, ecological, and evolutionary implications for domestic and wild birds. Future avian gustation research will develop our understanding of comparative biochemistry, molecular biology, and ethology—from an emphasis on anatomical structure to the physiological bases of behavior and performance.

#### REFERENCES

- Allison, A.C., 1953. The morphology of the olfactory system in the vertebrates. *Biol. Rev.* 28, 195–244.
- Amo, L., Caro, S.P., Visser, M.E., 2011a. Sleeping birds do not respond to predator odour. *PLoS One* 6, e27576.
- Amo, L., Visser, M.E., van Oers, K., 2011b. Smelling out predators is innate in birds. *Ardea* 99, 177–184.
- Amo, L., Galvan, I., Tomas, G., Sanz, J.J., 2008. Predator odour recognition and avoidance in a songbird. *Funct. Ecol.* 22, 289–293.
- Amo, L., Avilés, J.M., Parejo, D., Pena, A., Rodríguez, J., Tomas, G., 2012a. Sex recognition by odour and variation in the uropygial gland secretion in starlings. *J. Anim. Ecol.* 81, 605–613.
- Amo, L., López-Rull, I., Pagán, I., Garcia, C.M., 2012b. Male quality and conspecific scent preferences in the house finch, *Carpodacus mexicanus*. *Anim. Behav.* 84, 1483–1489.

- Andres, K.H., 1970. Anatomy and ultrastructure of the olfactory bulb in fish, amphibia, reptiles, birds and mammals. In: Wolstenhome, G.E.W., Knight, J. (Eds.), *Taste and Smell in Vertebrates*. J and A Churchill, London, pp. 177–196.
- Avery, M.L., Schreiber, C.L., Decker, D.G., 1999. Fruit sugar preferences of house finches. *Wilson Bull.* 111, 84–88.
- Baker, H.G., Baker, I., Hodges, S.A., 1998. Sugar composition of nectars and fruits consumed by birds and bats in the tropics and subtropics. *Biotropica* 30, 559–586.
- Balthazart, J., Schoffeniels, E., 1979. Pheromones are involved in the control of sexual behaviour in birds. *Naturwissenschaften* 66, 55–56.
- Balthazart, J., Taziaux, M., 2009. The underestimated role of olfaction in avian reproduction? *Behav. Brain Res.* 200, 248–259.
- Bang, B.G., 1960. Anatomical evidence for olfactory function in some species of birds. *Nature* 188, 547–549.
- Bang, B.G., 1961. The surface pattern of the nasal mucosa and its relation to mucous flow—a study of chicken and herring gull nasal mucosae. *J. Morphol.* 109, 57–71.
- Bang, B.G., 1963. Comparative studies of the nasal organs of birds: a study of 28 species of birds of West Bengal. *PAVO* 1, 79–89.
- Bang, B.G., 1964. The nasal organs of the black and Turkey vultures: a comparative study of the cathartid species *Coragyps atratus atratus* and *Canhartes aura septentrionalis* (with notes on *Cathartes aura falklandica*, *Pseudogyps bengalensis*, and *Neophron percnopterus*). *J. Morphol.* 115, 153–184.
- Bang, B.G., 1965. Anatomical adaptations for olfaction in the snow petrel. *Nature* 205, 513–515.
- Bang, B.G., 1966. The olfactory apparatus of tube-nosed birds (Procellariiformes). *Acta Anat.* 65, 391–415.
- Bang, B.G., 1971. Functional anatomy of the olfactory system in 23 orders of birds. *Acta Anat. (Suppl. 79)*, S1–S76.
- Bang, B.G., Cobb, S., 1968. The size of the olfactory bulb in 108 species of birds. *Auk* 85, 55–61.
- Bang, B., Wenzel, B.M., 1985. Nasal cavity and olfactory system. In: King, A.S., McLelland, J. (Eds.), *Form and Function in Birds III*. Academic Press, London, pp. 195–225.
- Barnett, C.A., Skelhorn, J., Bateson, M., Rowe, C., 2011. Educated predators make strategic decisions to eat defended prey according to their toxin content. *Behav. Ecol.* 23, 418–424.
- Bath, W., 1906. Die Geschmacksorgane der Vögel und Krokodile. *Arch. Biontol.* 1, 1–47.
- Berkhoudt, H., 1985. Structure and function of avian taste receptors. In: Levy, A.S., McLelland, J. (Eds.), *Form and Function in Birds III*. Academic Press, New York, pp. 463–496.
- Bertin, A., Calandreau, L., Arnould, C., Lèvy, F., 2012. The developmental stage of chicken embryos modulates impact of in ovo olfactory stimulation on food preferences. *Chem. Senses* 37, 253–261.
- Bertin, A., Calandreau, L., Arnould, C., Nowak, R., Lèvy, F., Noirot, V., Bouvarel, I., Leterrier, C., 2010. *In ovo* olfactory experience influences post-hatch feeding behaviour in young chickens. *Ethology* 116, 1027–1037.
- Bilko, A., Altbacker, V., Hudson, R., 1994. Transmission of food preference in the rabbit—the means of information-transfer. *Physiol. Behav.* 56, 907–912.
- Blem, C.R., Blem, L.B., Felix, J., van Gelder, J., 2000. Rufous hummingbird sucrose preference: precision of selection varies with concentration. *Condor* 102, 235–238.
- Bonadonna, F., Nevitt, G.A., 2004. Partner-specific odor recognition in an Antarctic seabird. *Science* 306, 835.
- Bonadonna, F., Cunningham, G.B., Jouventin, P., Hesters, F., Nevitt, G.A., 2003a. Evidence for nest-odour recognition in two species of diving petrel. *J. Exp. Biol.* 206, 3719–3722.
- Bonadonna, F., Hesters, F., Jouventin, P., 2003b. Scent of a nest: discrimination of own-nest odours in Antarctic prions, *Pachyptila desolata*. *Behav. Ecol. Sociobiol.* 54, 174–178.
- Bonadonna, F., Sanz-Aguilar, A., 2012. Kin recognition and inbreeding avoidance in wild birds: the first evidence for individual kin-related odour recognition. *Anim. Behav.* 84, 509–513.
- Brown, M., Downs, C.T., Johnson, S.D., 2008. Sugar preferences of nectar feeding birds – a comparison of experimental techniques. *J. Avian Biol.* 39, 479–483.
- Brown, M., Downs, C.T., Johnson, S.D., 2010a. Concentration-dependent sugar preferences of the Malachite sunbird (*Nectarinia famosa*). *Auk* 127, 151–155.
- Brown, M., Downs, C.T., Johnson, S.D., 2010b. Sugar preferences and digestive efficiency in an opportunistic avian nectarivore, the dark-capped bulbul *Pycnonotus tricolor*. *J. Ornithol.* 151, 637–643.
- Brown, M., Downs, C.T., Johnson, S.D., 2010c. Sugar preferences of a generalist nonpasserine flower visitor, the African speckled mousebird (*Colius striatus*). *Auk* 127, 781–786.
- Brown, M., Downs, C.T., Johnson, S.D., 2012. African red-winged starlings prefer hexose sugar solutions, but do not like them too sweet. *J. Ornithol.* 153, 265–272.
- Brugger, K.E., Nelms, C.O., 1991. Sucrose avoidance by American robins (*Turdus migratorius*): implications for control of bird damage in fruit crops. *Crop Prot.* 10, 455–460.
- Burne, T.H.J., Rogers, L.J., 1999. Changes in olfactory responsiveness by the domestic chick after early exposure to odorants. *Anim. Behav.* 58, 329–336.
- Butler, A.B., Cotterill, R.M.J., 2006. Mammalian and avian neuroanatomy and the question of consciousness in birds. *Biol. Bull.* 211, 106–127.
- Byerly, M.S., Simon, J., Cogburn, L.A., Bihan-Duval, E.L., Duclos, M.J., Aggrey, S.E., Porter, T.E., 2010. Transcriptional profiling of hypothalamus during development of adiposity in genetically selected fat and lean chickens. *Physiol. Genomics* 42, 157–167.
- Callanan, D., Dixon, M., Widdicombe, J.G., Wise, J.C.M., 1974. Responses of geese to inhalation of irritant gases and injection of phenyl diguanide. *Resp. Physiol.* 22, 157–166.
- Calvin, A.D., Williams, C.M., Westmoreland, N., 1957. Olfactory sensitivity in the domestic pigeon. *Am. J. Psychol.* 188, 255–256.
- Campagna, S., Mardon, J., Celerier, A., Bonadonna, F., 2011. Potential semiochemical molecules from birds: a practical and comprehensive compilation of the last 20 years studies. *Chem. Senses* 37, 3–25.
- Caro, S.P., Balthazart, J., 2010. Pheromones in birds: myth or reality? *J. Comp. Physiol. A* 196, 751–766.
- Caspers, B.A., Krause, E.T., 2010. Odour-based natal nest recognition in the zebra finch (*Taeniopygia guttata*), a colony-breeding songbird. *Biol. Lett.* 7, 184–186.
- Caterina, M.J., Schumacher, M.A., Tominaga, M., Rosen, T.A., Levine, J.D., Julius, D., 1997. The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature* 389, 816–824.
- Chalcoff, V.R., Aizen, M.A., Galetto, L., 2008. Sugar preferences of the green-backed firecrown hummingbird (*Sephanoides sephanioides*): a field experiment. *Auk* 125, 60–66.
- Chamorro, C.A., de Paz, P., Fernández, J.G., Anel, L., 1993. Fungiform papillae of the pig and the wild boar analysed by scanning electron microscopy. *Scanning Microsc.* 7, 313–322.

- Clapperton, B.K., Porter, R.E.R., Day, T.D., Waas, J.R., Matthews, L.R., 2012. Designer repellents: combining olfactory, visual or taste cues with a secondary repellent to deter free-ranging house sparrows from feeding. *Pest Manage. Sci.* 68, 870–877.
- Clark, L., 1991a. Odor detection thresholds in tree swallows and cedar waxwings. *Auk* 108, 177–180.
- Clark, L., 1991b. The nest protection hypothesis: the adaptive use of plant secondary compounds by European starlings. In: Loye, J.E., Zuk, M. (Eds.), *Bird-Parasite Interactions: Ecology Evolution and Behaviour*. Oxford University Press, Oxford, pp. 205–221.
- Clark, L., 1995. Modulation of avian responsiveness to chemical irritants: effects of prostaglandin E1 and analgesics. *J. Exp. Zool.* 271, 432–440.
- Clark, L., 1996. Trigeminal repellents do not promote conditioned odor avoidance in European starlings. *Wilson Bull.* 108, 36–52.
- Clark, L., 1997. A review of the bird repellent effects of 117 carbocyclic compounds. In: Mason, J.R. (Ed.), *Repellents in Wildlife Management*. National Wildlife Research Center, Fort Collins, CO, pp. 343–352.
- Clark, L., 1998. Physiological, ecological, and evolutionary bases for the avoidance of chemical irritants by birds. In: Nolan, V. Jr., Ketterson, E.D. (Eds.), *Current Ornithology*, vol. 14. Plenum Press, New York, pp. 1–37.
- Clark, L., Avery, M.L., 2013. Factors influencing the effectiveness of chemical repellents in management of birds on airports. In: DeVault, T.L., Blackwell, B.F., Belant, J.L. (Eds.), *Wildlife in Airport Environments*. Johns Hopkins University Press, Baltimore.
- Clark, L., Mason, J.R., 1985. Use of nest material as insecticidal and anti-pathogenic agents by the European starling. *Oecologia* 67, 169–176.
- Clark, L., Mason, J.R., 1987. Olfactory discrimination of plant volatiles by the European starling. *Anim. Behav.* 35, 227–235.
- Clark, L., Mason, J.R., 1988. Effect of biologically active plants used as nest material and the derived benefit to starling nestlings. *Oecologia* 77, 174–180.
- Clark, L., Mason, J.R., 1989. Sensitivity of brown-headed cowbirds to volatiles. *Condor* 91, 922–932.
- Clark, L., Mason, J.R., 1993. Interactions between sensory and post-ingestional repellents in starlings: methyl anthranilate and sucrose. *Ecol. Appl.* 3, 262–270.
- Clark, L., Shah, P.S., 1991. Nonlethal bird repellents: in search of a general model relating repellency and chemical structure. *J. Wildl. Manage.* 55, 538–545.
- Clark, L., Shah, P.S., 1992. Information content of prey odor plumes: what do foraging Leach's storm petrels know? In: Doty, R.L., Müller-Schwartz, D. (Eds.), *Chemical Signals in Vertebrates*. Plenum Press, New York, pp. 421–427.
- Clark, L., Shah, P.S., 1993. Chemical bird repellents: possible use in cyanide ponds. *J. Wildl. Manage.* 57, 657–664.
- Clark, L., Shah, P., 1994. Tests and refinements of a general structure-activity model for avian repellents. *J. Chem. Ecol.* 20, 321–339.
- Clark, L., Smeraski, C.A., 1990. Seasonal shifts in odor acuity by starlings. *J. Exp. Zool.* 255, 22–29.
- Clark, L., Avilova, K.V., Bean, N.J., 1993. Odor thresholds in passerines. *Comp. Biochem. Physiol. A* 104, 305–312.
- Clark, L., Shah, P.S., Mason, J.R., 1991. Chemical repellency in birds: relationship between chemical structure and avoidance response. *J. Exp. Zool.* 260, 310–322.
- Conigrave, A.D., Brown, E.M., 2006. Taste receptors in the gastrointestinal tract II. L-amino acid sensing by calcium-sensing receptors: implications for GI physiology. *Am. J. Physiol. Gastrointest. Liver Physiol.* 291, G753–G761.
- Cortright, D.N., Krause, J.E., Broom, D.C., 2007. TRP channels and pain. *Biochim. Biophys. Acta* 1772, 978–988.
- Crocker, D.R., Perry, S.M., 1990. Plant chemistry and bird repellents. *Ibis* 132, 300–308.
- Crocker, D.R., Perry, S.M., Wilson, M., Bishop, J.D., Scanlon, C.B., 1993. Repellency of cinnamic acid derivatives to captive rock doves. *J. Wildl. Manage.* 57, 113–122.
- Cunningham, G.B., Nevitt, G.A., 2011. Evidence for olfactory learning in procellariiform seabird chicks. *J. Avian Biol.* 42, 85–88.
- Davis, R.G., 1973. Olfactory psychophysical parameters in man, rat, dog and pigeon. *J. Comp. Physiol. Psychol.* 85, 221–232.
- Davis, J.K., Lowman, J.J., Thomas, P.J., ten Hallers, B.F.H., Koriabine, M., Huynh, L.Y., Maney, D.L., de Jong, P.J., Martin, C.L., Thomas, J.W., 2010. Evolution of a bitter taste receptor gene cluster in a new world sparrow. *Genome Biol. Evol.* 2, 358–370.
- De León, A., Mínguez, E., Belliure, B., 2003. Self-odour recognition in European storm-petrel chicks. *Behaviour* 140, 925–933.
- DeGroof, G., Gwinner, H., Steiger, S., Kempnaers, B., van der Linden, A., 2010. Neural correlates of behavioural olfactory sensitivity changes seasonally in European Starlings. *PLoS One* 5, e14337.
- Douglas III, H.D., 2008. Prenuptial perfume: alloanoointing in the social rituals of the crested auklet (*Aethia cristatella*) and the transfer of arthropod deterrents. *Naturwissenschaften* 95, 45–53.
- Douglas III, H.D., 2013. Colonial seabird's paralytic perfume slows lice down: an opportunity for parasite-mediated selection? *Int. J. Parasitol.* 43, 399–407.
- Downs, C.T., 1997. Sugar preference and apparent sugar assimilation in the red lory. *Austral. J. Zool.* 45, 613–619.
- Downs, C.T., Perrin, M.R., 1996. Sugar preferences of some southern African nectarivorous birds. *Ibis* 138, 455–459.
- Dubbeldam, J.L., Karten, H.J., 1978. The trigeminal system in the pigeon (*Columba livia*). I. Projections of the gasserian ganglion. *J. Comp. Neurol.* 180, 661–678.
- Dubbeldam, J.L., Veenman, C.L., 1978. Studies on the somatotopy of the trigeminal system in the mallard, *Anas platyrhynchos* L. I. The ganglion trigeminale. *Neth. J. Zool.* 28, 150–160.
- Dugas-Ford, J., Rowell, J.J., Ragsdale, C.W., 2012. Cell-type homologies and the origins of the neocortex. *PNAS* 109, 16974–16979. <http://dx.doi.org/10.1073/pnas.1204773109>.
- Duncan, C.J., 1960. The sense of taste in birds. *Ann. Appl. Biol.* 48, 409–414.
- Eicholz, M.W., Dassow, J.A., Stafford, J.D., Weatherhead, P.J., 2012. Experimental evidence that nesting ducks use mammalian urine to assess predator abundance. *Auk* 129, 638–644.
- Elliot, B., 1937. Total distribution of taste buds on the tongue of the kitten at birth. *J. Comp. Neurol.* 66, 361–373.
- Espallat, J.E., Mason, J.R., 1990. Differences in taste preference between red-winged blackbirds and European starlings. *Wilson Bull.* 102, 292–299.
- Fleming, P.A., Bakken, B.H., Lotz, C.N., Nicolson, S.W., 2004. Concentration and temperature effects on sugar intake and preferences in a sunbird and a hummingbird. *Funct. Ecol.* 18, 223–232.
- Fleming, P.A., Xie, S., Napier, K., McWhorter, T.J., Nicolson, S.W., 2008. Nectar concentration affects sugar preferences in two Australian honeyeaters and a lorikeet. *Funct. Ecol.* 22, 599–605.
- Forsman, J.T., Monkkonen, M., Korpimäki, E., Thomson, R.L., 2013. Mammalian nest predator feces as a cue in avian habitat selection decisions. *Behav. Ecol.* 24, 262–266.
- Furuse, M., Mabayo, R.T., Okumura, J.-I., 1996. The role of gustation in oil preference in the chicken. *Jpn. Poult. Sci.* 33, 256–260.

- Gagliardo, A., 2013. Forty years of olfactory navigation in birds. *J. Exp. Biol.* 216, 2165–2171.
- Gagliardo, A., Filannino, C., Ioalè, P., Pecchia, T., Wikelski, M., Vallortigara, G., 2011. Olfactory lateralization in homing pigeons: a GPS study on birds released with unilateral olfactory inputs. *J. Exp. Biol.* 214, 593–598.
- Gagliardo, A., Ioalè, P., Savini, M., Wild, J.M., 2006. Having the nerve to home: trigeminal magnetoreceptor versus olfactory mediation of homing in pigeons. *J. Exp. Biol.* 209, 2888–2892.
- Gagliardo, A., Ioalè, P., Savini, M., Wild, J.M., 2009. Navigational abilities of adult and experienced homing pigeons deprived of olfactory or trigeminally mediated magnetic information. *J. Exp. Biol.* 212, 3119–3124.
- Gagliardo, A., Pecchia, T., Savini, M., Odetti, F., Ioalè, P., Vallortigara, G., 2007. Olfactory lateralization in homing pigeons: initial orientation of birds receiving a unilateral olfactory input. *Eur. J. Neurosci.* 25, 1511–1516.
- Ganchrow, D., Ganchrow, J.R., 1985. Number and distribution of taste buds in the oral cavity of hatchling chicks. *Physiol. Behav.* 34, 889–894.
- Gatica, C.D.L., Gonzalez, S.P., Vasquez, R.A., Sabat, P., 2006. On the relationship between sugar digestion and diet preference in two Chilean avian species belonging to the Muscicapoidae superfamily. *Rev. Chilena Hist. Nat.* 79, 287–294.
- Gentle, M.J., 1975. Gustatory behavior of the chicken and other birds. In: Wright, P., Caryl, P.E., Vowles, D.M. (Eds.), *Neural and Endocrine Aspects of Behaviors in Birds*. Elsevier Scientific, Amsterdam, pp. 305–308.
- Gentle, M.J., 1989. Cutaneous sensory afferents recorded from the *nerve intramandibularis* of *Gallus gallus* var *domesticus*. *J. Comp. Physiol.* A 164, 763–774.
- Gentle, M.J., 2011. Pain issues in poultry. *Appl. Anim. Behav. Sci.* 135, 252–258.
- Gentle, M.J., Hill, F.L., 1987. Oral lesions in the chicken: behavioural responses following nociceptive stimulation. *Physiol. Behav.* 40, 781–783.
- Gentle, M.J., Hunter, L.N., 1993. Neurogenic inflammation in the chicken (*Gallus gallus* var *domesticus*). *Comp. Biochem. Physiol.* C 105, 459–462.
- Getty, R., 1975. *Sisson and Grossman's the Anatomy of the Domestic Animals*. W.B. Saunders Company, Philadelphia.
- Godard, R.D., Bowers, B.B., Wilson, C.M., 2007. Eastern bluebirds *Sialia sialis* do not avoid nest boxes with chemical cues from two common nest predators. *J. Avian Biol.* 38, 128–131.
- Gomez, G., Celli, A., 2008. The peripheral olfactory system of the domestic chicken: physiology and development. *Brain Res. Bull.* 76, 208–216.
- Grubb, T.C., 1974. Olfactory navigation to the nesting burrow in Leach's petrel (*Oceanodroma leucorhoa*). *Anim. Behav.* 22, 192–202.
- Gottschaldt, K.M., 1985. Structure and function of avian somatosensory receptors. In: King, A.S., McLelland, J. (Eds.), *Form and Function in Birds*, vol. 3. Academic Press, London, pp. 375–461.
- Güntürkün, O., 2012. The convergent evolution of neural substrates for cognition. *Psychol. Res.* 2, 212–219.
- Gwinner, H., 1997. The function of green plants in nests of European starlings (*Sturnis vulgaris*). *Behaviour* 134, 337–351.
- Gwinner, H., Oltrogge, M., Trost, L., Nienaber, U., 2000. Green plants in starling nests: effects on nestlings. *Anim. Behav.* 59, 301–309.
- Gwinner, H., Berger, S., 2005. European starlings: nestling condition, parasites and green nest material during the breeding season. *J. Ornithol.* 146, 365–371.
- Gwinner, H., Berger, S., 2008. Starling males select green nest material by olfaction using experience-independent and experience-dependent cues. *Anim. Behav.* 75, 971–976.
- Hagelin, J.C., 2004. Observations on the olfactory ability of the Kakapo *Strigops habroptilus*, the critically endangered parrot of New Zealand. *Ibis* 146, 161–164.
- Hagelin, J.C., 2007a. Odors and chemical signaling. In: Jamieson, B.G.M. (Ed.), *Reproductive Biology and Phylogeny of Birds*, vol. 6B. Science Publishers, Enfield, NH, pp. 75–119.
- Hagelin, J.C., 2007b. The citrus-like scent of crested auklets: reviewing the evidence for an avian olfactory ornament. *J. Ornithol.* 148, S195–S201.
- Hagelin, J.C., Jones, I.L., 2007. Bird odors and other chemical substances: a defense mechanism or overlooked mode of intraspecific communication? *Auk* 124, 741–761.
- Hagelin, J.C., Jones, I.L., Rasmussen, L.E.L., 2003. A tangerine-scented social odor in a monogamous seabird. *Proc. R. Soc. Ser. B* 270, 1323–1329.
- Hagelin, J.C., Simonet, J.C., Lyson, T.R., 2013. Embryonic domestic chickens can detect compounds in an avian chemosignal before breathing air. In: East, M.L., Dehnhard, M. (Eds.), *Chemical Signals in Vertebrates*, vol. 12. Springer, New York, pp. 363–377.
- Harlander-Matauschek, A., Beck, P., Piepho, H.-P., 2009. Taste aversion learning to eliminate feather pecking in laying hens, *Gallus gallus domesticus*. *Anim. Behav.* 78, 485–490.
- Harlander-Matauschek, A., Beck, P., Rodenburg, T.B., 2010. Effect of an early bitter taste experience on subsequent feather-pecking behaviour in laying hens. *Appl. Anim. Behav. Sci.* 127, 108–114.
- Henton, W.W., 1969. Conditioned suppression to odorous stimuli in pigeons. *J. Exp. Anal. Behav.* 12, 175–185.
- Henton, W.W., Smith, J.C., Tucker, D., 1966. Odor discrimination in pigeons. *Science* 153, 1138–1139.
- Hepper, P.G., 1988. Adaptive fetal learning: prenatal exposure to garlic affects postnatal preferences. *Anim. Behav.* 36, 935–936.
- Hiestand, W.A., Randall, W.C., 1941. Species differentiation in the respiration of birds following carbon dioxide administration and the location of inhibitory receptors in the upper respiratory tract. *J. Cell. Comp. Physiol.* 17, 333–340.
- Hile, A.G., Shan, Z., Zhang, S.-Z., Block, E., 2004. Aversion of European starlings (*Sturnis vulgaris*) to garlic oil treated granules: garlic oil as an avian repellent. Garlic oil analysis by nuclear magnetic resonance spectroscopy. *J. Agr. Food Chem.* 52, 2192–2196.
- Hinman, A., Chuang, H., Bautista, D.M., Julius, D., 2006. TRP channel activation by reversible covalent modification. *PNAS* 103, 19564–19568.
- Holland, R.A., Thorup, K., Gagliardo, A., Bisson, I.-A., Knecht, E., Mizrahi, D., Wikelski, M., 2009. Testing the role of sensory systems in the migratory heading of a songbird. *J. Exp. Biol.* 212, 4065–4071.
- Holzer, P., 2011. Transient receptor potential (TRP) channels as drug targets for diseases of the digestive system. *Pharmacol. Ther.* 131, 142–170.
- Houston, D.C., 1986. Scavenging efficiency of turkey vultures in tropical forests. *Condor* 88, 318–323.
- Hu, H.J., Bhawe, G., Gereau, R.W., 2002. Prostaglandin and protein kinase A-dependent modulation of vanilloid receptor function by metabotropic glutamate receptor 5: potential mechanism for thermal hyperalgesia. *J. Neurosci.* 22, 7444–7452.
- Hughes, B.O., Wood-Gush, D.G.M., 1971. A specific appetite for calcium in domestic chickens. *Anim. Behav.* 19, 490–499.

- Hutchison, L.V., Wenzel, B.M., 1980. Olfactory guidance in foraging by procellariiforms. *Condor* 82, 314–319.
- Hwang, S.W., Cho, H., Kwak, J., Lee, S.-Y., Kang, C.-J., Jung, J., Cho, S., Min, K.H., Suh, Y.-G., Kim, D., Oh, U., 2000. Direct activation of capsaicin receptors by products of lipoxygenases: endogenous capsaicin-like substances. *PNAS* 97, 6155–6160.
- Hyman, L.H., 1942. *Comparative Vertebrate Anatomy*, second ed. University of Chicago Press, Chicago.
- Jackson, S., Nicolson, S.W., Lotz, C.N., 1998a. Sugar preferences and “side bias” in cape sugarbirds and lesser double-collared sunbirds. *Auk* 115, 156–165.
- Jackson, S., Nicolson, S.W., van Wyk, B.-E., 1998b. Apparent absorption efficiencies of nectar sugars in the cape sugarbird, with a comparison of methods. *Physiol. Zool.* 71, 106–115.
- Johnson, L.S., Murphy, S.M., Parrish, G.W., 2011. Lack of predator-odor detection and avoidance in a songbird, the House Wren. *J. Field Ornithol.* 82, 150–157.
- Johnson, S.D., Hargreaves, A.L., Brown, M., 2006. Dark, bitter-tasting nectar function as a filter of flower visitors in a bird-pollinated plant. *Ecology* 87, 2709–2716.
- Johnson, S.D., Nicolson, S.W., 2008. Evolutionary associations between nectar properties and specificity in bird pollination systems. *Biol. Lett.* 4, 49–52.
- Jones, I.L., Hagelin, J.C., Major, H.L., Rasmussen, L.E.L., 2004. An experimental field study of the function of crested auklet feather odor. *Condor* 106, 71–78.
- Jordt, S.-E., Julius, D., 2002. Molecular basis for species-specific sensitivity to “hot” chili peppers. *Cell* 108, 421–430.
- Jung, Y., Wirkus, D., Amendola, D., Gomez, G., 2005. Characteristics of odorant elicited calcium fluxes in acutely-isolated chick olfactory neurons. *J. Comp. Physiol. A* 191, 511–520.
- Kare, M.R., 1961. Comparative aspects of taste. In: Kare, M.R., Halpern, B.P. (Eds.), *Physiological and Behavioral Aspects of Taste*. University of Chicago Press, Chicago, pp. 13–23.
- Kare, M.R., Mason, J.R., 1986. The chemical senses in birds. In: Sturkie, P.D. (Ed.), *Avian Physiology*. Springer-Verlag, New York.
- Karlsson, A.C., Jensen, P., Elgland, M., Laur, K., Fymer, T., Konradsson, P., Laska, M., 2010. Red junglefowl have individual body odors. *J. Exp. Biol.* 213, 1619–1624.
- Kassarov, L., 2001. Do cyanogenic glycosides and pyrrolizidine alkaloids provide some butterflies with a chemical defense against their bird predators? A different point of view. *Behaviour* 138, 45–67.
- Keller, M., Baum, M.J., Brock, O., Brennan, P.A., Bakker, J., 2009. The main and the accessory olfactory systems interact in the control of mate recognition and sexual behavior. *Behav. Brain Res.* 200, 268–276.
- Keverne, E.B., Murphy, C.L., Silver, W.L., Wysocki, C.J., Meredith, M., 1986. Non-olfactory chemoreceptors of the nose: recent advances in understanding the vomeronasal and trigeminal systems. *Chem. Senses* 11, 119–133.
- Kirifides, M.L., Kurnellas, M.P., Clark, L., Bryant, B.P., 2004. Calcium responses of chicken trigeminal ganglion neurons to methyl anthranilate and capsaicin. *J. Exp. Biol.* 207, 715–722.
- Ko, I.W.P., 1996. Sugar preferences of fruit-eating birds in Hong Kong. *Hong Kong Bird Rep.*, 165–169.
- Koda, H., Minagawa, M., Si-Hong, L., Mizumura, K., Kumazawa, T., 1996. H1-receptor-mediated excitation and facilitation of the heat response by histamine in canine visceral polymodal receptors studied in vitro. *J. Neurophysiol.* 76, 1396–1404.
- Köhler, A., Leseigneur, C.D.C., Verburt, L., Nicolson, S.W., 2010. Dilute bird nectars: viscosity constrains food intake by licking in a sunbird. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 299, R1068–R1074.
- Köhler, A., Verburt, L., Fleming, P.A., Nicolson, S.W., 2008. Changes in nectar concentration: how quickly do whitebellied sunbirds (*Cinnyris talatala*) adjust feeding patterns and food intake? *J. Comp. Physiol. B* 178, 785–793.
- Kolattukudy, P.E., Bohnet, S., Rogers, L., 1987. Diesters of 3-hydroxy fatty acids produced by the uropygial glands of female mallards uniquely during the mating season. *J. Lipid Res.* 28, 582–588.
- Krause, E.T., Caspers, B.A., 2012. Are olfactory cues involved in nest recognition in two social species of estrildid finches? *PLoS One* 7, e36615.
- Kudo, K.-i., Shiraishi, J.-i., Nishimura, S., Bungo, T., Tabata, S., 2010. The number of taste buds is related to bitter taste sensitivity in layer and broiler chickens. *Anim. Sci. J.* 81, 240–244.
- Kuenzel, W.J., 2007. Neurobiological basis of sensory perception: welfare implications of beak trimming. *Poult. Sci.* 86, 1273–1282.
- Lalloué, F.L., Ayer-Le-Lièvre, C.S., Sicard, G., 2003. Analysis of the functional maturation of olfactory neurons in chicks before and after birth. *Chem. Senses* 28, 729–737.
- Lane, S.J., 1997. Preferences and apparent digestibilities of sugars by fruit damaging birds in Japan. *Ann. Appl. Biol.* 130, 361–370.
- Leseigneur, C.D.C., Nicolson, S.W., 2009. Nectar concentration preferences and sugar intake in the white-bellied sunbird, *Cinnyris talatala* (Nectariniidae). *J. Comp. Physiol. B* 179, 673–679.
- Lindenmaier, P., Kare, M.R., 1959. The taste end-organs of the chicken. *Poult. Sci.* 38, 545–550.
- Lotz, C.N., Nicolson, S.W., 1996. Sugar preferences of a nectarivorous passerine bird, the lesser double-collared sunbird (*Nectarinia chalybea*). *Funct. Ecol.* 10, 360–365.
- Lotz, C.N., Schondube, J.E., 2006. Sugar preferences in nectar- and fruit-eating birds: behavioral patterns and physiological causes. *Biotropica* 38, 3–15.
- Mabayo, R.T., Okumura, J.-I., Furuse, M., 1996. Dietary flavor modifies oil preferences in the chicken. *Appl. Anim. Behav. Sci.* 49, 213–221.
- Macadar, A.W., Rausch, L.J., Wenzel, B.M., Hutchison, L.V., 1980. Electrophysiology of the olfactory pathway in the pigeon. *J. Comp. Physiol.* 137, 39–46.
- Macpherson, L.J., Bailong, X., Kwan, K.Y., Petrus, M.J., Dubin, A.E., Hwang, S., Cravatt, B., Corey, D.P., and Patapoutian, A., 2007. An ion channel essential for sensing chemical damage. *J. Neurosci.* 27, 11412–11415.
- Mardon, J., Bonadonna, F., 2009. Atypical homing or self-odour avoidance? Blue petrels (*Halobaena caerulea*) are attracted to their mate’s odour but avoid their own. *Behav. Ecol. Sociobiol.* 63, 537–542.
- Mardon, J., Saunders, S.M., Bonadonna, F., 2011. From preen secretions to plumage: the chemical trajectory of blue petrels’ *Halobaena caerulea* social scent. *J. Avian Biol.* 42, 29–38.
- Mardon, J., Saunders, S.M., Anderson, M.J., Couchoux, C., Bonadonna, F., 2010. Species, gender, and identity: cracking petrels’ sociochemical code. *Chem. Senses* 35, 309–321.
- Markman, S., Tadmor-Melamed, H., Arieli, A., Izhaki, I., 2006. Sex differences in food intake and digestive constraints in a nectarivorous bird. *J. Exp. Biol.* 209, 1058–1063.
- Martinez del Rio, C., Schondube, J.E., McWhorter, T.J., Herrera, L.G., 2001. Intake responses in nectar feeding birds: digestive and metabolic causes, osmoregulatory consequences, and coevolutionary effects. *Am. Zool.* 41, 902–915.

- Mason, J.R., Clark, L., 1995. Capsaicin detection in trained European starlings: the importance of olfaction and trigeminal chemoreception. *Wilson Bull.* 107, 165–169.
- Mason, J.R., Clark, L., 1997. Avian repellents: options, modes of action, and economic considerations. In: Mason, J.R. (Ed.), *Repellents in Wildlife Management*. USDA, National Wildlife Research Center, Fort Collins, CO, pp. 371–391.
- Mason, J.R., Clark, L., 2000. The chemical senses in birds. In: Whittow, G.C. (Ed.), *Sturkie's Avian Physiology*, fifth ed. Academic Press, San Diego, pp. 39–56.
- Mason, J.R., Maruniak, J.A., 1983. Behavioral and physiological effects of capsaicin in red-winged blackbirds. *Pharmacol. Biochem. Behav.* 19, 857–862.
- Mason, J.R., Adams, M.A., Clark, L., 1989. Anthranilate repellency to starlings: chemical correlates and sensory perception. *J. Wildl. Manage.* 53, 55–64.
- Mason, J.R., Bean, N.J., Shah, P.S., Clark, L., 1991. Taxon-specific differences in responsiveness to capsaicin and several analogues: correlates between chemical structure and behavioral aversiveness. *J. Chem. Ecol.* 17, 2539–2551.
- Mastrota, F.N., Mench, J.A., 1995. Evaluation of taste repellents with northern bobwhites for deterring ingestion of granular pesticides. *Environ. Toxicol. Chem.* 14, 631–638.
- Mata, A., Bosque, C., 2004. Sugar preferences, absorption efficiency and water influx in a neotropical nectarivorous passerine, the bananaquit (*Coereba flaveola*). *Comp. Biochem. Physiol. A* 139, 395–404.
- Mathis, A., Ferrari, M.C.O., Windel, N., Messier, F., Chivers, D.P., 2008. Learning by embryos and the ghost of predation future. *Proc. R. Soc. Ser. B* 275, 2603–2607.
- Matson, K.D., Millam, J.R., Klasing, K.C., 2000. Taste threshold determination and side-preference in captive cockatiels (*Nymphicus hollandicus*). *Appl. Anim. Behav. Sci.* 69, 313–326.
- Matson, K.D., Millam, J.R., Klasing, K.C., 2001. Thresholds for sweet, salt, and sour taste stimuli in cockatiels (*Nymphicus hollandicus*). *Zoo Biol.* 20, 1–13.
- Matson, K.D., Millam, J.R., Klasing, K.C., 2004. Cockatiels (*Nymphicus hollandicus*) reject very low levels of plant secondary compounds. *Appl. Anim. Behav. Sci.* 85, 141–156.
- Mateo, J.M., Johnston, R.E., 2000. Kin recognition and the 'armpit effect': evidence of self-referent phenotype matching. *Proc. R. Soc. Ser. B* 267, 695–700.
- Matthes, E., 1934. Geruchsorgan, Lubosch Handbuch der vergleichenden Anatomie der Wirbeltiere, Gropfert, Kallius, vol. 11, Urban and Schwarzenbeig, Berlin.
- McKeegan, D.E.F., 2002. Spontaneous and odour evoked activity in single avian olfactory bulb neurones. *Brain Res.* 929, 48–58.
- McKeegan, D.E.F., 2009. Avian chemoreception: an electrophysiological approach. international symposium on olfaction and taste. *Ann. N. Y. Acad. Sci.* 1170, 438–441.
- McKeegan, D.E.F., Lippens, N., 2003. Adaptation responses of single avian olfactory bulb neurones. *Neurosci. Lett.* 344, 83–86.
- McKeegan, D.E.F., Demmers, T.G.M., Wathes, C.M., Jones, R.B., Gentle, M.J., 2002. Stimulus-response functions of single avian olfactory bulb neurones. *Brain Res.* 953, 101–111.
- McKeegan, D.E.F., Smith, F.S., Demmers, T.G.M., Wathes, C.M., Jones, R.B., 2005. Behavioral correlates of olfactory and trigeminal gaseous stimulation in chickens, *Gallus domesticus*. *Physiol. Behav.* 84, 761–768.
- McWhorter, T.J., Martinez del Rio, C., 2000. Does gut function limit hummingbird food intake? *Physiol. Biochem. Zool.* 73, 313–324.
- Medina-Tapia, N., Ayala-Berdon, J., Morales-Pérez, L., Melo, L.M., Schondube, J.E., 2012. Do hummingbirds have a sweet-tooth? Gustatory sugar thresholds and sugar selection in the broad-billed hummingbird *Cyananthus latirostris*. *Comp. Biochem. Physiol. A* 161, 307–314.
- Meisami, E., 1991. Chemoreception. In: Prosser, C.L. (Ed.), *Neural and Integrative Animal Physiology*. John Wiley and Sons, New York, NY, pp. 335–362.
- Mennella, J.A., Jagnow, C.P., Beauchamp, G.K., 2001. Prenatal and postnatal flavor learning by human infants. *Pediatrics* 107, e88. <http://dx.doi.org/10.1542/peds.107.6.e88>.
- Michelsen, W.J., 1959. Procedure for studying olfactory discrimination in pigeons. *Science* 130, 630–631.
- Miller, L., 1942. Some tagging experiments with black-footed albatrosses. *Condor* 44, 3–9.
- Miller, Jr., I.J., Reedy, Jr., F.E., 1990. Variations in human taste bud density and taste intensity perception. *Physiol. Behav.* 47, 1213–1219.
- Mínguez, E., 1997. Olfactory nest recognition by British storm-petrel chicks. *Anim. Behav.* 53, 701–707.
- Moncrieff, R.W., 1946. *The Chemical Senses*. Wiley, New York.
- Moore, C.A., Elliot, R., 1946. Numerical and regional distribution of taste buds on the tongue of the bird. *J. Comp. Neurol.* 84, 119–131.
- Nakajima, S., Onimaru, S., 2006. Salt discrimination in domestic pigeons (*Columba livia domestica*): poisonous LiCl solution versus equimolar safe NaCl solution. *J. Ethol.* 24, 59–65.
- Nevitt, G.A., 2011. The neuroecology of dimethyl sulfide: a global-climate regulator turned marine infochemical. *Integr. Comp. Biol.* 51, 819–825.
- Nevitt, G.A., Veit, R.R., Kareiva, P., 1995. Dimethyl sulphide as a foraging cue for Antarctic procellariiform seabirds. *Nature* 376, 680–682.
- Necker, R., 1974. Dependence of mechanoreceptor activity on skin temperature in sauropsids. II. Pigeon and duck. *J. Comp. Physiol.* 92, 75–83.
- Necker, R., 2000. The somatosensory system. In: Whittow, G.C. (Ed.), *Sturkie's Avian Physiology*, fifth ed., pp. 57–69.
- Nicolson, S.W., Fleming, P.A., 2003. Nectar as food for birds: the physiological consequences of drinking dilute sugar solutions. *Plants Syst. Evol.* 238, 139–153.
- Norman, D.L., Mason, J.R., Clark, L., 1992. Capsaicin effects on consumption of food by cedar waxwings and house finches. *Wilson Bull.* 104, 549–551.
- Odendaal, T.C., Brown, M., Downs, C.T., Johnson, S.D., 2010. Sugar preferences and digestive efficiency of the village weaver: a generalist avian pollinator of African plants. *J. Exp. Biol.* 213, 2531–2535.
- Ohta, T., Ikemi, Y., Murakami, M., Imagawa, T., Otsuguro, K., Ito, S., 2006. Potentiation of transient receptor potential V1 functions by the activation of metabotropic 5-HT receptors in rat primary sensory neurons. *J. Physiol.* 576, 809–822.
- Papi, F., Casini, G., 1990. Pigeons with ablated pyriform cortex home from familiar but not from unfamiliar sites. *PNAS* 87, 3783–3787.
- Papi, F., Fiore, L., Fiaschi, V., Benvenuti, S., 1971. The influence of olfactory nerve section on the homing capacity of carrier pigeons. *Monit. Zool. Ital.* 5, 265–267.
- Patapoutian, A., Tate, S., Woolf, C.J., 2009. Transient receptor potential channels: targeting pain at the source. *Nat. Rev. Drug Discov.* 8, 55–68.
- Patzke, N., Manns, M., Güntürkün, O., Ioalè, P., Gagliardo, A., 2010. Navigation-induced ZENK expression in the olfactory system of pigeons (*Columba livia*). *Eur. J. Neurosci.* 31, 2062–2072.

- Petit, C., Hossaert-McKey, M., Perret, P., Blondel, J., Lambrechts, M.M., 2002. Blue tits use selected plants and olfaction to maintain an aromatic environment for nestlings. *Ecol. Lett.* 5, 585–589.
- Pierau, F.-K., Sann, H., Harti, G., 1986. Resistance of birds to capsaicin and differences in their substance P (SP) system. *Proc. Int. Union Physiol. Sci.* 16, 207–211.
- Porter, R.H., Picard, M., 1998. Effects of early odor exposure in domestic chicks. *Reprod. Nutr. Dev.* 38, 441–448.
- Provenza, F.D., 1995. Origins of food preference in herbivores. In: Mason, J.R. (Ed.), *Repellents in Wildlife Management*. National Wildlife Research Center, Fort Collins, CO, pp. 81–90.
- Rahn, H., Ar, A., Paganelli, C.V., 1979. How bird eggs breathe. *Sci. Am.* 240, 46–55.
- Rastogi, A., Kumari, Y., Rani, S., Kumar, V., 2011. Phase inversion of neural activity in the olfactory and visual systems of a night-migratory bird during migration. *Eur. J. Neurosci.* 34, 99–109.
- Rehkämper, G., Frahm, H.D., Cnotka, J., 2008. Mosaic evolution and adaptive brain component alteration under domestication seen on the background of evolutionary theory. *Brain Behav. Evol.* 71, 115–126.
- Rehkämper, G., Haase, E., Frahm, H.D., 1988. Allometric comparison of brain weight and brain structure volumes in different breeds of the domestic pigeon, *Columba livia* f.d. (fantails, homing pigeons, strassers). *Brain Behav. Evol.* 31, 141–149.
- Rensch, B., Neunzig, R., 1925. Experimentelle untersuchungen über den geschmackssinn der vogel II. *J. Ornithol.* 73, 633–646.
- Rescorla, R.A., Solomon, R.L., 1967. Two-process learning theory: relationships between Pavlovian conditioning and instrumental learning. *Psychol. Rev.* 74, 151–182.
- Restrepo, D., Zviman, M.M., Rawson, N.E., 1995. Imaging of intracellular calcium in chemosensory receptor cells. In: Spielman, A.I., Brand, J.G. (Eds.), *Experimental Cell Biology of Taste and Olfaction*. CRC Press, Boca Raton, pp. 387–398.
- Rieke, G.K., Wenzel, B.M., 1974. The ipsilateral and olfactory projection field in the pigeon. In: Denton, D.A., Coghlan, J.P. (Eds.), *Olfaction and Taste*, vol. 5. Academic Press, New York, pp. 361–368.
- Rieke, G.K., Wenzel, B.M., 1978. Forebrain projections of the pigeon olfactory bulb. *J. Morph.* 158, 41–55.
- Rios, J.M., Mangione, A.M., Marone, L., 2012. Tolerance to dietary phenolics and diet breadth in three seed-eating birds: implications for graminivory. *J. Exp. Zool. A* 317, 425–433.
- Robinson, P.P., Winkles, P.A., 1990. Quantitative study of fungiform papillae and taste buds on the cat's tongue. *Anat. Rec.* 226, 108–111.
- Roper, T.J., 1999. Olfaction in birds. In: In: Snowden, C.T., Roper, T.J. (Eds.), *Advances in the Study of Behaviour*, vol. 28. Academic Press, Boston, MA, pp. 247–332.
- Roth, T.C., Cox, J.G., Lima, S.L., 2008. Can foraging birds assess predation risk by scent? *Anim. Behav.* 76, 2021–2027.
- Roura, E., Baldwin, M.W., Klasing, K.C., 2012. The Avian Taste System: an Update Proceedings of the Australian Poultry Science Symposium, Sydney, New South Wales, Australia 23, 97–104.
- Rowe, C., Skelhorn, J., 2005. Colour biases are a question of taste. *Anim. Behav.* 69, 587–594.
- Saito, I., 1966. Comparative anatomical studies of the oral organs of the poultry. V. Structures and distribution of taste buds of the fowl. *Bull. Fac. Agric. Univ. Miyazaki* 13, 95–102.
- Saito, S., Shingai, R., 2006. Evolution of thermo TRP ion channel homologs in vertebrates. *Physiol. Genomics* 27, 219–230.
- Saito, S., Fukuta, N., Shingai, R., Tominaga, M., 2011. Evolution of vertebrate transient receptor potential vanilloid 3 channels: opposite temperature sensitivity between mammals and western clawed frogs. *PLoS Genet.* 7, e1002041.
- Sann, H., Harti, G., Pierau, F.-K., Simon, E., 1987. Effect of capsaicin upon afferent and efferent mechanisms of nociception and temperature regulation in birds. *Can. J. Physiol. Pharmacol.* 65, 1347–1354.
- Saxton, V.P., Creasy, G.L., Paterson, A.M., Trought, M.C.T., 2009. Behavioral responses of European blackbirds and Australasian silvereyes to varying acid and sugar levels in artificial grapes. *Am. J. Enol. Vit.* 60, 82–86.
- Saxton, V.P., Mulder, I., Creasy, G.L., Paterson, A.M., Ross, J.G., Trought, M.C.T., 2011. Comparative behavioural responses of silvereyes (*Zosterops lateralis*) and European blackbirds (*Turdus merula*) to secondary metabolites in grapes. *Austral. Ecol.* 36, 233–239.
- Schaal, B., Marlier, L., Soussignan, R., 2000. Human fetuses learn odours from their pregnant mother's diet. *Chem. Senses* 25, 729–737.
- Schondube, J.E., Martinez del Rio, C., 2003. Concentration-dependent sugar preferences in nectar-feeding birds: mechanisms and consequences. *Funct. Ecol.* 17, 445–453.
- Schondube, J.E., Martinez del Rio, C., 2004. Sugar and protein digestion in flowerpiercers and hummingbirds: a comparative test of adaptive convergence. *J. Comp. Physiol. B* 174, 263–273.
- Schrader, E., 1970. Die Topographie der Kopfnerven vom Huhn (Ph.D. Dissertation). Freie Univ., Berlin.
- Schwenk, K., 1985. Occurrence, distribution and functional significance of taste buds in lizards. *Copeia* 1, 91–101.
- Shah, P., Clark, L., Mason, J.R., 1991. Prediction of avian repellency from chemical structure: the aversiveness of vanillin, vanillyl alcohol and veratryl alcohol. *Pestic. Biochem. Physiol.* 40, 169–175.
- Shaw, C.L., Rutter, J.E., Austin, A.L., Garvin, M.C., Whelan, R.J., 2011. Volatile and semivolatile compounds in gray catbird uropygial secretions vary with age and between breeding and wintering grounds. *J. Chem. Ecol.* 37, 329–339.
- Shi, P., Zhang, J., 2005. Contrasting modes of evolution between vertebrate sweet/umami receptor genes and bitter receptor genes. *Mol. Biol. Evol.* 23 (2), 292–300.
- Shibuya, T., Tucker, D., 1967. Single unit responses of olfactory receptors in vultures. In: Hayashi, T. (Ed.), *Olfaction-and Taste*. Pergamon Press, Oxford, pp. 219–220.
- Shibuya, T., Tonosaki, K., 1972. Electrical responses of single olfactory receptor cells in some vertebrates. In: Schneider, D. (Ed.), *Olfaction and Taste*. Wissenschaftliche Verlagsgesellschaft MBH, Stuttgart, pp. 102–108.
- Silver, W.L., Maruniak, J.A., 1981. Trigeminal chemoreception in the nasal and oral cavities. *Chem. Senses* 6, 295–305.
- Skelhorn, J., Griksaitis, D., Rowe, C., 2008. Colour biases are more than a question of taste. *Anim. Behav.* 75, 827–835.
- Skelhorn, J., Rowe, C., 2010. Birds learn to use distastefulness as a signal of toxicity. *Proc. R. Soc. B* 277, 1729–1734.
- Sneddon, H., Hadden, R., Hepper, P.G., 1998. Chemosensory learning in the chicken embryo. *Phys. Behav.* 64, 133–139.
- Snyder, G.K., Peterson, T.T., 1979. Olfactory sensitivity in the black-billed magpie and in the pigeon. *Comp. Biochem. Physiol. A* 62, 921–925.
- Stager, K.E., 1964. The role of olfaction in food location by the Turkey vulture (*Cathartes aura*). *Los Ang. Cnty. Mus. Contrib. Sci.* 81, 1–63.
- Stager, K.E., 1967. Avian olfaction. *Am. Zool.* 7, 415–420.
- Stattelman, A.J., Talbot, R.B., Coulter, D.B., 1975. Olfactory thresholds of pigeons (*Columba livia*), quail (*Colinus virginianus*) and chickens (*Gallus gallus*). *Comp. Biochem. Physiol. A* 50, 807–809.
- Steiger, S.S., Fidler, A.E., Valcu, M., Kempenaers, B., 2008. Avian olfactory receptor gene repertoires: evidence for a well-developed sense of smell in birds? *Proc. R. Soc. Ser. B* 275, 2309–2317.

- Stucky, C.L., Dubin, A.E., Jeske, N.A., Malin, S.A., McKerny, D.D., Story, G.M., 2009. Roles of transient receptor potential channels in pain. *Brain Res. Rev.* 60, 2–23.
- Szolcsanyi, J., 1982. Capsaicin type pungent agents producing pyrexia. In: Milton, A.S. (Ed.), *Pyretics and Antipyretics*. Springer-Berlin Heidelberg, pp. 437–478.
- Szolcsanyi, J., Sann, H., Pierau, F.-K., 1986. Nociception in pigeons is not impaired by capsaicin. *Pain* 27, 247–260.
- Taher, A.I., Gleaves, E.W., Beck, M., 1984. Special calcium appetite in laying hens. *Poult. Sci.* 63, 2261–2267.
- Tewksbury, J.J., Nabhan, G.P., 2001. Seed dispersal. Directed deterrence by capsaicin in chilies. *Nature* 412, 403–404.
- Todrank, J., Heth, G., Restrepo, D., 2011. Effects of *in utero* odorant exposure on neuroanatomical development of the olfactory bulb and odour preferences. *Proc. R. Soc. Ser. B* 278, 1949–1955.
- Tolhurst, B.E., Vince, M.A., 1976. Sensitivity to odours in the embryo of the domestic fowl. *Anim. Behav.* 24, 772–779.
- Tordoff, M.G., 2001. Calcium: taste, intake, and appetite. *Physiol. Rev.* 81, 1567–1597.
- Tordoff, M.G., Shao, H., Alarcón, L.K., Margolskee, R.F., Mosinger, B., Bachmanov, A.A., Reed, D.R., McCaughey, S., 2008. Involvement of TIR3 in calcium–magnesium taste. *Physiol. Genomics* 34, 338–348.
- Tucker, D., 1963. Olfactory, vomeronasal and trigeminal receptor responses to odorants. In: Zotterman, Y. (Ed.), *Olfaction and Taste*. Pergamon, New York, pp. 45–69.
- Tucker, D., 1965. Electrophysiological evidence for olfactory function in birds. *Nature* 207, 34–36.
- Tucker, D., 1971. Nonolfactory responses from the nasal cavity: Jacobson's organ and the trigeminal system. In: Beidler, L.M. (Ed.), *Handbook of Sensory Physiology IV: Chemical Senses Olfaction*. Springer-Verlag, Berlin, pp. 151–181.
- Tucker, D., Graziadei, P.C., Smith, J.C., 1974. Recovery of olfactory function in pigeons after bilateral transection of the olfactory nerves. In: Denton, D.A., Coghlan, J.P. (Eds.), *Olfaction and Taste*, vol. 5. Academic Press, New York, pp. 369–374.
- Ueda, H., Kainou, S., 2005. Aversion to quinine is associated with taste sensation in chicks. *J. Poult. Sci.* 42, 254–262.
- Ueda, H., Shigemizu, G., 2001. Feeding response to tea saponin in chicks given diet selection. *J. Poult. Sci.* 38, 333–342.
- Ueda, H., Takagi, A., Katou, K., Matsumoto, S., 2002. Feeding behavior in chicks fed tea saponin and quinine sulfate. *J. Poult. Sci.* 39, 34–41.
- Vermaut, S., De Coninck, K., Flo, G., Cokelaere, M., Onagbesan, M., Decuyper, E., 1997. Effect of deoiled jojoba meal on feed intake in chickens: satiating or taste effect? *J. Agr. Food Chem.* 45, 3158–3163.
- Walker, J.C., Tucker, D., Smith, J.C., 1979. Odor sensitivity mediated by trigeminal nerve in the pigeon. *Chem. Senses* 4, 107–116.
- Walker, J.C., Walker, D.B., Tambiah, C.R., Gilmore, K.S., 1986. Olfactory and nonolfactory odor detection in pigeons: elucidation by a cardiac acceleration paradigm. *Physiol. Behav.* 38, 575–580.
- Wallraff, H.G., 1988. Olfactory deprivation in pigeons: examination of methods applied in homing experiments. *Comp. Biochem. Physiol. A Comp. Physiol.* 89, 621–629.
- Wallraff, H., 2005. Beyond familiar landmarks and integrated routes: goal-oriented navigation by birds. *Connect. Sci.* 17, 91–106.
- Warner, R.L., McFarland, L.Z., Wilson, W.O., 1967. Microanatomy of the upper digestive tract of the Japanese quail. *Am. J. Vet. Res.* 28, 1537–1548.
- Watkins, R.W., Lumley, J.A., Gill, E.L., Bishop, J.D., Langton, S.D., Mac-Nicoll, A.D., Price, N.R., Drew, M.G.B., 1999. Quantitative structure-activity relationships (QSAR) of cinnamic acid bird repellents. *J. Chem. Ecol.* 25, 2825–2845.
- Weldon, P.J., Rappole, J.H., 1997. *J. Chem. Ecol.* 23, 2609–2632.
- Wellmann, A.E., Downs, C.T., 2009. Sugar preferences and digestion by cape white-eyes, *Zosterops virens*, fed artificial fruit diets. *Afr. Zool.* 44, 106–116.
- Wenzel, B.M., 1973. Chemoreception. In: Farner, D.S., King, J.R., Parkes, K.C. (Eds.), *Avian Biology*, vol. 3. Academic Press, New York, pp. 389–415.
- Wenzel, B.M., Sieck, M.H., 1972. Olfactory perception and bulbar electrical activity in several avian species. *Physiol. Behav.* 9, 287–293.
- Werner, S.J., Kimball, B.A., Provenza, F.D., 2008. Food color, flavor, and conditioned avoidance among red-winged blackbirds. *Physiol. Behav.* 93, 110–117.
- Werner, S.J., Provenza, F.D., 2011. Reconciling sensory cues and varied consequences of avian repellents. *Physiol. Behav.* 102, 158–163.
- Whelan, R.J., Levin, T.C., Owen, J.C., Garvin, M.C., 2010. Short-chain carboxylic acids from gray catbird (*Dumetella carolinensis*) uropygial secretions vary with testosterone levels and photoperiod. *Comp. Biochem. Physiol. B* 156, 183–188.
- Whittaker, D.J., Soini, H.A., Atwell, J.W., Hollars, C., Novotny, M.V., Ketterson, E.D., 2010. Songbird chemosignals: volatile compounds in preen gland secretions vary among individuals, sexes and populations. *Behav. Ecol.* 21, 608–614.
- Whittaker, D.J., Richmond, K.M., Miller, A.K., Kiley, R., Burns, C.B., Atwell, J.W., Ketterson, E.D., 2011. Intraspecific preen oil preferences in dark-eyed juncos (*Junco hyemalis*). *Behav. Ecol.* 22, 1256–1263.
- Wild, J.M., 1985. The avian somatosensory system. I. Primary spinal afferent input to the spinal cord and brainstem in the pigeon (*Columba livia*). *J. Comp. Neurol.* 240, 377–395.
- Wilson, A.L., Downs, C.T., 2011. Food preferences of Knysna and purple-crested turacos fed varying concentrations of equicaloric and equimolar artificial fruit. *J. Exp. Biol.* 214, 607–612.
- Wolf, C.J., Ma, Q., 2007. Nociceptors-noxious stimulus detectors. *Neuron* 55, 353–364.
- Wood-Gush, D.G.M., Kare, M.R., 1966. The behaviour of calcium-deficient chickens. *Brit. Poult. Sci.* 7, 285–290.
- Zelenitsky, D.K., Therrien, F.O., Ridgely, R.C., McGee, A.R., Witmer, L.M., 2011. Evolution of olfaction in non-avian theropod dinosaurs and birds. *Proc. R. Soc. Ser. B* 278, 3625–3634.
- Zidar, J., Løvlie, H., 2012. Scent of the enemy: behavioral responses to predator faecal odour in the fowl. *Anim. Behav.* 84, 547–554.
- Zhao, H., Zhang, J., 2012. Mismatches between feeding ecology and taste receptor evolution: an inconvenient truth. *PNAS* 109, E1464.
- Zygmunt, P.M., Petersson, J., Andersson, D.A., Chuang, H., Sörgård, M., Di Marzo, V., Julius, D., Högestätt, E.D., 1999. Vanilloid receptors on sensory nerves mediate the vasodilator action of anandamide. *Nature* 400, 452–457.

This page intentionally left blank