

December 12, 2013

# Ice Crystal Formation and Evolution in Five Campaigns: START08, HIPPO Global, DC3, PREDICT and TORERO

Minghui Diao, *National Center of Atmospheric Research*

Joshua DiGangi, *Princeton University*

Anthony O'Brien, *Princeton University*


Mark A. Zondlo, *Princeton University*



This work is licensed under a [Creative Commons CC BY-NC International License](https://creativecommons.org/licenses/by-nc/4.0/).



# Ice crystal formation and evolution in five campaign: START08, HIPPO Global, DC3, PREDICT and TORERO



NCAR ASP postdoctoral fellowship; NASA NESSF Fellowship: NASA NNX09AO51H; Walbridge Graduate Award; National Science Foundation (Grant No. EEC-0540832); NSF ATM-0840732 for VCSEL (Princeton); START08, HIPPO Global, DC3, PREDICT, TORERO Science Team


## Cirrus cloud and ice supersaturation

Cirrus clouds have large but highly uncertain impacts on Earth's climate [Chen et al. 2000]. Cirrus cloud formation requires supersaturation of the relative humidity with respect to ice (RHi). However, it has not clear how ice crystal regions initiate from ice supersaturated regions (ISSRs, regions where RHi > 100%), grow in size and eventually dissipate. Here we show the time evolution of cirrus clouds by analyzing the relationship between ice crystal regions and ISSRs at temperature (T) ≤ -40 °C.

## VCSEL hygrometer on the NSF Gulfstream V research plane



1. Stratosphere-Troposphere Analyses of Regional Transport campaign (2008)
2. HIAPER Pole-to-Pole Observations (HIPPO) Global campaign (2009-2011)
3. PRE-Depression Investigation of Cloud-systems in the Tropics (PREDICT) campaign (2010)
4. Deep Convective Clouds & Chemistry (DC3) campaign (2012)
5. Tropical Ocean Troposphere Exchange of Reactive halogen species and Oxygenated VOC (TORERO) campaign (2012)



GV instruments	Principal Investigator	Measurement	Accuracy	Precision
VCSEL hygrometer	Mark Zondlo, Princeton U.	Water vapor (Zondlo et al., 2010)	6%	≤ 1%
2-DC	Dave Rogers, NCAR	Ice particle number density (Nc) and mean diameter (Dc)	Measurement range: 25-800 μm (Korolev et al., 2011)	
SID-2H instrument	Andrew Heymsfield, NCAR	Ice particle Nc and Dc	Measurement range: 1-50 μm (Cotton et al., 2010)	

## Definitions of ice crystal regions (ICRs) and ice supersaturated regions (ISSRs)

**ISSRs:** regions with spatially continuous ISS.

**ICRs:** regions with spatially continuous ice crystal distribution.

"With ice crystals" as where the ice crystals are observed during the 1 Hz measurements, while the remaining regions are considered to be clear-sky regions.

**One ISSR+ICR sample:** a set of spatially continuous ISSRs and ICRs.

### References:

Chen, T., W.B. Rossow and Y.C. Zhang (2000), Radiative effects of cloud-type variations. *J. Clim.*, 13, 264-286.  
Diao, M., Zondlo, M. A., Heymsfield, A. J., Beaton, S. P. and Rogers, D. C.: Evolution of ice crystal regions on the microscale based on in situ observations, *Geophysical Research Letters*, 40, 3473-3478, doi:10.1002/grl.50665, 2013.  
M. Diao, M. A. Zondlo, A. J. Heymsfield, L. M. Avallone, M. E. Paige, S. P. Beaton, T. Campos, and D. C. Rogers. Cloud-scale ice supersaturated regions spatially correlate with high water vapor heterogeneities. *Atmos. Chem. Phys. Discuss.*, 13, 22249-22296, 2013.

Minghui Diao<sup>1</sup>, Joshua DiGangi<sup>2</sup>, Anthony O'Brien<sup>2</sup>, Mark A. Zondlo<sup>2</sup>

1. National Center of Atmospheric Research/Advanced Study Program (ASP)

2. Civil and Environmental Engr, Princeton University, NJ

Contact: [diao@ucar.edu](mailto:diao@ucar.edu)

## Five phases of ice crystal region evolution


Phase 1: Clear-sky ISSRs

Phase 2: Ice crystal nucleation


Phase 3: Ice crystal early growth

Phase 4: Ice crystal later growth

Phase 5: Sedimentation/evaporation




one ISSR + ICR sample




Phase	Description	Spatial ratio M = $\frac{\sum(L_{ICR})}{L_{ISSR+ICR}}$	Spatial ratio N = $\frac{\sum(L_{ISSR})}{L_{ISSR+ICR}}$
1	Clear-sky ISSRs	0	1
2	Nucleation	$0 < M < 1$	1
3	Early growth of ice crystals	$0 < M < 1$	$0 < N < 1$
4	Later growth of ice crystals	1	$0 < N \leq 1$
5	Evaporation/sedimentation	1	0

## RHi mean and standard deviation evolution




## Ice crystal number density (Nc) and mean diameter (Dc) evolution

Mean diameter Dc



Number density logNc



Growth rate of a single ice crystal


$$\frac{dD_{ice}}{dt} = \frac{1}{D_{ice}} (S_{v,out} - S_{v,eq}) * G_i(T, P)$$

Rogers and Yau, 1989 and Straka, 2009

[1] Mean diameter of ice crystals merges into a constant value as ice crystals grow, which agrees with the theory of ice crystal growth rate.

[2] Number density of ice crystals continues to increase throughout the ICR evolution. The increasing Nc agrees with previous simulations, where new ice crystals continue to form as the air parcel continues to be uplifted [Spichtinger and Gierens, 2009].

## Ice crystal region lifetime phases and horizontal spatial expansion evolution



[1] **DC3 shows more Phase 4** (ICRs with ISS buried inside) than other campaigns, suggesting that either the ICRs do not consume H<sub>2</sub>O efficiently, or continued strong uplift maintained ISS inside ICRs in DC3.

[2] **PREDICT shows less Phase 2** (ISSRs with ICRs buried inside), suggesting that ICRs expand immediately once they are formed in ISSRs.

[3] The peak of ICR/ISSR ratio happens at higher value in DC3 than the other two campaigns, suggesting that **ICRs expand relatively fast in DC3** so that most ICRs are larger than ISSRs. This finding further indicates that the dominance of Phase 4 in DC3 is contributed by strong uplift instead of inefficient depositional growth.

## Implications to understanding ice crystal evolution

1. We provided a new and simple method to distinguish three phases of ice cloud evolution by using in-situ and quasi-Eulerian sampling of two common parameters: 1) RHi 2) presence/absence of ice crystals.

2. We demonstrated the importance of separating out various evolution phases of ice crystals in Eulerian view observation, since they have different properties of RHi, Nc and Dc.

3. We compared ice crystal evolution between various geographical location and meteorological background. Our finding improves the understanding of the relative lifetime of ice crystal evolution.

4. Our result facilitates the comparison between in situ Eulerian observation and Lagrangian view cloud simulation/ parameterization.