

University of Massachusetts Amherst

From the Selected Works of Robert Hallock

1979

Behavior of Persistent Currents Under Conditions of Strong Decay

Robert Hallock, *University of Massachusetts - Amherst*
D. T Ekholm



Available at: https://works.bepress.com/robert_hallock/40/

be injected into a plasma from large limiter plates with good secondary electron yields to keep the sheath potential low.

This work was supported by the U. S. Department of Energy under Contract No. EY-76-C-03-0010 PA 26.

¹Lena Oren and R. J. Taylor, Nucl. Fusion **17**, 1143 (1977).

²P. E. Stott, C. C. Daughney, and R. A. Ellis, Nucl. Fusion **15**, 431 (1977).

³J. W. M. Paul *et al.*, in *Proceedings of the Sixth International Conference on Plasma Physics and Controlled Nuclear Fusion Research, Berchtesgaden, 1976*

(International Atomic Energy Agency, Vienna, 1977), Vol. 2, p. 269.

⁴S. Suckewer and R. J. Hawryluk, Phys. Rev. Lett. **40**, 1649 (1978).

⁵Lena Oren and R. J. Taylor, in *Proceedings of the Third International Conference on Plasma Surface Interactions in Controlled Fusion Devices, Abington, United Kingdom, March 1977 (to be published), paper 8.*

⁶K. Ohasa *et al.*, in *Proceedings of the Third International Conference on Plasma Surface Interactions in Controlled Fusion Devices, Abington, United Kingdom, March 1977 (to be published), paper 47.*

⁷P. J. Harbour and M. F. A. Harrison, in *Proceedings of the Third International Conference on Plasma Surface Interactions in Controlled Fusion Devices, Abington, United Kingdom, March 1977 (to be published), paper 25.*

Behavior of Persistent Currents under Conditions of Strong Decay

D. T. Ekholm and R. B. Hallock

Department of Physics and Astronomy, University of Massachusetts, Amherst, Massachusetts 01003
(Received 27 November 1978)

We report observations of the decay of persistent currents of thin superfluid ⁴He films for which the superfluid velocity deviates strongly from the previously observed $\log t$ behavior. We present data which document the evolution of this behavior as a function of film thickness. The Iordanskii-Langer-Fisher theory does not accurately describe our observations.

Superfluid ⁴He has the remarkable ability to attain a state of macroscopic persistent flow. It is generally assumed that such flow is, in fact, only metastable. In some cases the instability is weak enough to render the flow persistent over laboratory lifetimes. In other cases the metastability becomes clearly apparent. Previous to this report¹ all studies of the degradation in time of these persistent currents yielded results which could be adequately described by

$$v(t) = a - b \log t, \quad (1)$$

where $v(t)$ is the macroscopic velocity of the flow at time t and a and b are empirical parameters. In particular, Kojima *et al.*,² Kukich, Henkel, and Reppy,³ and Langer and Reppy⁴ have observed that the flow of bulk ⁴He in restricted geometries (compressed powders, Vycor, etc.) obeys Eq. (1). The Iordanskii-Langer-Fisher^{5,6} (ILF) model of persistent-current decay generally predicts Eq. (1) under conditions for which the superfluid velocity makes small deviations from a critical velocity, v_c .

Telschow and Hallock⁷ have also reported general agreement with Eq. (1) in studies of persistent flow in thin ⁴He films, but they observed this

behavior to be obeyed even for changes in the velocity of as much as 60% away from the initial⁸ velocity, v_0 . We present results here of studies of persistent currents in ⁴He films for which the decays in some cases appear to cause the complete extinction of the flow. Our results as a function of film thickness show clearly for the first time the evolution of deviations from Eq. (1).

The apparatus used for the present work is an improved version of that used^{7,9} earlier in this laboratory. Persistent currents are generated on a Pyrex annular ring without rotation using a thermal technique.⁷ The speed of the persistent flow is measured using the well-known techniques of Doppler-shifted third sound.¹⁰ We have observed that the strength and nature of the velocity decays does not depend on the presence of the third sound necessary to make the measurements. Film thickness values are determined¹¹ from the temperature and vapor pressure and are consistent to within a few percent with thickness values deduced¹¹ from the observed third-sound velocity. Changes in the film thickness due to the flow of the film are observable¹² but are too small to be relevant here.

Data on the decay of the persistent currents are

obtained from measurements of the flow velocity as a function of time. In each of the cases presented in Fig. 1 we have plotted v/v_0 to facilitate the comparison although v_0 was within 20% of 25 cm/sec for each case. It is observed that even for substantial changes in the film velocity the persistent-current decays are remarkably well represented by Eq. (1). Our present results are in qualitative agreement¹³ with earlier work⁷ in the limited region of overlap of the data. A clear evolution of deviations from Eq. (1) is observed as the film thickness is reduced below about eight layers.

It is clear that one must search beyond Eq. (1) to describe our observations. A simple empirical expression fits the data rather well over the full range of film thickness values studies. We find

$$v(t) = A(1 + Bt)^{-n}. \quad (2)$$

This is consistent with a deceleration of the form

$$dv/dt = -cv^\alpha, \quad (3)$$

where $\alpha = n^{-1} + 1$. The observed dependence of α on film thickness is shown in Fig. 2.

The form Eq. (3) can be recovered from the ILF theory if the full expression for the activation energy, E_a , of a vortex pair oriented perpendicular

to the substrate is used, i.e.,

$$E_a = (d\langle\rho_s\rangle K^2/2\pi)\{\ln(K/2\pi va) - 1\}.$$

Here K is the quantum of circulation and a is the vortex core radius. With this $\alpha = d\langle\rho_s\rangle K^2/2\pi k_B T$ and this¹⁴ is shown on Fig. 2 as the solid curve. We conclude that the ILF theory is only in qualitative agreement with the data. The observed dependence of α on d is more closely represented by the form $\alpha = \alpha_0 + (d - d^*)^4$ over the relatively limited range of film thickness values for which persistent-current decays can be accurately measured. We find $\alpha_0 \approx 2$. For a true two-dimensional superfluid we would expect^{15,16} Eq. (3) with $\alpha \approx 3$.

An objection to the use of the ILF theory^{5,6} has been advanced by Donnelly and Roberts.¹⁷ They have argued that the velocities needed to provide an appropriate energy barrier are much larger than the macroscopic currents¹⁸ typically present in a restricted-geometry flow apparatus. Donnelly and Roberts propose that a finite barrier exists even in the absence of superflow due to the presence of boundaries in the real system. The competition between the barriers for forward and backward nucleation yields a prediction




FIG. 1. Velocity of the metastable film currents as a function of time, at $T = 1.45$ K. The ratio v/v_0 has been used to facilitate comparison among the various film thickness values. The numbers in the upper right are the measured film thickness values in atomic layers.




FIG. 2. The exponent α as a function of film thickness. The solid curve is the value of $\alpha = d\langle\rho_s\rangle K^2/2\pi k_B T$ at $T = 1.45$ K.

for the decay of persistent flow of the form given by Donnelly, Hills, and Roberts¹⁹ as

$$v = F \ln \left(\frac{1 + G \exp(-\nu_0 t)}{1 - G \exp(-\nu_0 t)} \right). \quad (4)$$

Here $F = k_B T / p_c$, where k_B is the Boltzman constant, T the temperature, and p_c the momentum of the critical excitation. G is a number which depends on the initial velocity of the persistent flow and

$$\nu_0 = (2nKfp_c/k_B T) \exp(-\Delta E/k_B T),$$

where n is the linear density of candidates for nucleation, f is an attempt frequency, and ΔE the barrier height for $v=0$. In general Eq. (4) represents the data reasonably well although for times late in the evolution of the persistent flow the data indicate that the velocity decreases more slowly than predicted.¹⁹ The quantity ν_0 as deduced from fits to the data of Fig. 1 is shown in Fig. 3 versus the thickness of the helium film. The solid lines are the predictions¹⁹ for two arbitrary values of f .

We have also observed that it is possible to change the rate of decay of a given metastable flow merely by changing the temperature. An example of this is shown in Fig. 4. The solid curve represents the time evolution of the velocity of a film at $T=1.50$ K at a thickness of eight atomic layers. During a subsequent decay from




FIG. 3. Nucleation rate ν_0 as a function of film thickness at 1.45 K. The circles represent the observed values of ν_0 and the solid lines are obtained from Ref. 19.

the same initial velocity, represented by the solid circles, the temperature was changed for a time to a lower value of 1.40 K. Upon returning the temperature to its original value the decay rate increased. The crosses plotted here represent the coordinates $(v, t - t_0^*)$ where t is the clock time and t_0^* is the interval of time during which the film was at the lower temperature. One interpretation, among others, of this observation is that the vorticity present in the film is frozen in by a lowering of the temperature. Under this point of view what matters is the thermally activated *motion* (hopping) of the vortices rather than the thermal nucleation of the vortices themselves.

In conclusion, our observations of the complete decay of metastable superfluid ⁴He film currents has revealed substantial deviations from the decay rule encountered in all previous experiments. Predictions based on the ILF theory^{5,6} are in qualitative but not quantitative agreement with our results and hence we conclude that that theory is incomplete. The predictions of Donnelly,




FIG. 4. Decay of a metastable current subject to a change in the temperature. The solid curve shows the decay at the constant temperature of 1.50 K. The symbols display the observed decay as discussed in the text. The transition to lower temperatures was made during the interval between the arrows. The later warmup interval is compressed by the time scale. When the persistent current is warmed it continues its decay—it does not reproduce a virgin decay with a lower initial velocity. t_0^* was defined as the interval of time between the midpoints of the cooldown and warmup periods. At large values of the time several of the data represented by the dots were essentially superimposed; this results in the appearance of more x's than dots for these data.

Hills, and Roberts¹⁹ are in qualitative agreement with the data except perhaps at times late in the evolution of the current decays. A quantitative test awaits theoretical estimates for the parameters which are inherent in the theory.

We are pleased to acknowledge helpful and productive conversations with V. Ambegaokar, S. Doniach, R. Donnelly, and J. Reppy. One of us (R.B.H.) is grateful for the hospitality of the Aspen Center for Physics. This work was supported by the National Science Foundation through Grants No. DMR 76-08260 and DMR 78-07762.

¹We have previously commented briefly on preliminary data: D. T. Ekholm and R. B. Hallock, *J. Phys. (Paris), Colloq.* **39**, C6-306 (1978).

²H. Kojima, W. Vieth, E. Guyon, and I. Rudnick, in *Low Temperature Physics, LT-13*, edited by W. J. O'Sullivan, K. D. Timmerhaus, and E. F. Hommel (Plenum, New York, 1974), Vol. 1, p. 239.

³G. Kukich, R. P. Henkel, and J. D. Reppy, *Phys. Rev. Lett.* **21**, 197 (1968).

⁴J. S. Langer and J. D. Reppy, in *Progress in Low Temperature Physics*, edited by C. J. Gorter (North-Holland, Amsterdam, 1970), Vol. 6, p. 1.

⁵S. V. Iordanskii, *Zh. Eksp. Teor. Fiz.* **48**, 708 (1965) [*Sov. Phys. JETP* **21**, 467 (1965)].

⁶J. S. Langer and M. E. Fisher, *Phys. Rev. Lett.* **19**, 560 (1967).

⁷K. L. Telschow and R. B. Hallock, *Phys. Rev. Lett.* **37**, 1484 (1976).

⁸We define the velocity v_0 to be the velocity of the flow at $t = 15$ sec after the creation of the persistent current.

⁹K. L. Telschow, D. T. Ekholm, and R. B. Hallock, in *Quantum Fluids and Solids*, edited by S. B. Trickey *et al.* (Plenum, New York, 1977), p. 421.

¹⁰See, for example, Ref. 7 and K. R. Atkins and I. Rudnick, in *Progress in Low Temperature Physics*, edited by C. J. Gorter (North-Holland, Amsterdam, 1970), Vol. 6, Chap. 2.

¹¹We have used $d^3 = 27[T \ln(P_0/P)]^{-1}$, where d is the film thickness in atomic layers, P_0 the saturated vapor pressure at temperature T , and P the pressure in the container which houses the persistent-current flow path. These measurements are consistent to within a few percent with thickness values deduced from $C_3^2 = 3 \langle \rho_s/\rho \rangle (\beta/d^3) [1 + TS/L]^2$, where C_3 is the third-sound velocity, $\langle \rho_s/\rho \rangle$ the reduced superfluid fraction in the film, β the van der Waals constant, S the entropy, and L the latent heat.

¹²D. T. Ekholm and R. B. Hallock, *Phys. Rev. B* (to be published).

¹³One quantitative difference between this work and that reported in Ref. 7 is that a given decay here appears to occur in a somewhat thicker film than in the work reported in Ref. 7. A difference in the substrate could be the cause. The present work utilized a precision capacitive pressure gauge not available earlier and it is possible that our earlier thickness values were in error by a few tenths of an atomic layer. The relative comparisons presented here are of course unaffected.

¹⁴To calculate $\langle \rho_s \rangle$ we have used $\rho_s(1 - D/d)$, where ρ_s is the bulk superfluid density and $D = 0.5 + 2.45(\rho T / \rho_s T \lambda)$.

¹⁵B. A. Huberman, R. J. Meyerson, and S. Doniach, *Phys. Rev. Lett.* **40**, 780 (1978).

¹⁶V. Ambegaokar, B. I. Halperin, D. R. Nelson, and E. D. Siggia, *Phys. Rev. Lett.* **40**, 783 (1978).

¹⁷R. J. Donnelly and P. H. Roberts, *Philos. Trans. Roy. Soc. London* **271**, 41 (1971).

¹⁸This ignores the possibility that higher-velocity, local, microscopic currents of velocity proportional to the macroscopic velocity might exist.

¹⁹R. J. Donnelly, R. N. Hills, and P. H. Roberts, to be published.

Ultrasonic Attenuation near the 208-K Phase Transition of RbAg_4I_5

K. M. Leung and D. L. Huber

Department of Physics, University of Wisconsin, Madison, Wisconsin 53706

(Received 21 August 1978)

Using general expressions for the velocity and ultrasonic attenuation coefficient we have analyzed ultrasonic data for the 208-K phase transition in RbAg_4I_5 . From fitting the velocity in the high-temperature phase we obtain values for the strain and pseudospin coupling constants. We extract the intrinsic pseudospin relaxation rate from an analysis of the attenuation. The Jahn-Teller model is found to provide a good description of the ultrasonic behavior associated with the phase transition.

The 208-K phase transition in the superionic conductor RbAg_4I_5 has been of considerable interest recently.¹⁻¹³ The nature of this transition has been established to be three-dimensional Ising-

like.⁴ This conclusion is based on a study of the specific heat, which yields Ising-like exponents and amplitudes, and on a study of the birefringence in the ordered phase. The birefringence