

2001

Numerical Solution of Fuzzy Differential Equation by Runge-Kutta Method

S. Abbasbandy

T. Allah Viranloo

Numerical Solution of Fuzzy Differential Equation by Runge-Kutta Method

S. Abbasbandy* T. Allah viranloo[†]

April 13, 2002

Abstract

In this paper numerical algorithms for solving 'fuzzy ordinary differential equations' are considered. A scheme based on the 4th Runge-Kutta method in detail is discussed and this is followed by a complete error analysis. The algorithm is illustrated by solving some linear and nonlinear fuzzy Cauchy problems.

Keywords: Fuzzy differential equation, 4th Runge-Kutta method, Fuzzy Cauchy problem.

AMS subject classification: 34A12, 65L05.

1 Introduction

The concept of fuzzy derivative was first introduced by S.L. Chang, L.A. Zadeh in [1] it was followed up by D. Dubois, H. Prade in [2], who defined and used the extension principle. The fuzzy differential equation and the initial value problem were regularly treated by O. Kaleva in [5],[6] and by S. Seikkala in [7], The numerical method for solving fuzzy differential equations is introduced by M. Ma, M. Friedman, A. Kandel in [10] by the standard Euler method. paper organize as follows:

In section 2 some basic results on fuzzy numbers and definition of a fuzzy derivative, which have been discussed by S. Seikkala in [10] are given. In section 3 we define the problem, this is a fuzzy Cauchy problem whose numerical solution is the main interest of this work. The numerically solving fuzzy differential equation by 4th Runge-kutta method is discussed in section 4. The proposed algorithm is illustrated by solving some examples in section 5 and conclusion is in section 6.

*Department of Mathematics, Imam Khomeini International University, Qazvin, Iran.(abbasbandy@yahoo.com)

[†]Department of Mathematics Science and Research Branch, Islamic Azad University, Tehran, Iran.

2 Preliminaries

Consider the initial value problem

$$\begin{cases} y'(t) = f(t, y(t)); & a \leq t \leq b, \\ y(a) = \alpha, \end{cases} \quad (1)$$

The basis of all Runge-Kutta method is to express the difference between the value of y at t_{n+1} and t_n as

$$y_{n+1} - y_n = h \sum_{i=1}^m w_i k_i. \quad (2)$$

where for $i = 1, 2, \dots, m$, the w_i 's are constants and

$$k_i = f(t_n + \alpha_i h, y_n + h \sum_{j=1}^{i-1} \beta_{ij} k_j). \quad (3)$$

Equation (2) is to be exact for powers of h through h^m . Therefore, the truncation error T_m , can be written as

$$T_m = \gamma_m h^{m+1} + O(h^{m+2}),$$

The true magnitude of γ_m will generally be much less than the bound. Thus, if the $O(h^{m+2})$ term is small compared with $\gamma_m h^{m+1}$, as we expect it will be if h is small, then the bound on $\gamma_m h^{m+1}$ will usually be a bound on the error as a whole. The famous nonzero constants α_i , β_{ij} in 4th Runge-Kutta method are $m = 4$ and

$$\alpha_1 = 0 \quad , \quad \alpha_2 = \alpha_3 = 1/2 \quad , \quad \alpha_4 = 1 \quad , \quad \beta_{21} = 1/2 \quad , \quad \beta_{32} = 1/2 \quad , \quad \beta_{43} = 1,$$

and we have, see [9]

$$\begin{aligned} y_0 &= \alpha, \\ k_1 &= f(t_i, y_i), \\ k_2 &= f(t_i + \frac{h}{2}, y_i + \frac{h}{2}k_1), \\ k_3 &= f(t_i + \frac{h}{2}, y_i + \frac{h}{2}k_2), \\ k_4 &= f(t_i + h, y_i + hk_3), \\ y_{i+1} &= y_i + \frac{h}{6}(k_1 + 2k_2 + 2k_3 + k_4), \end{aligned} \quad (4)$$

where

$$a = t_0 \leq t_1 \leq \dots \leq t_N = b \text{ and } h = \frac{(b-a)}{N} = t_{i+1} - t_i. \quad (5)$$

Theorem 1 Let $f(t, y)$ belong to $C^4[a, b]$ and let its partial derivatives are bounded and assume there exists, P, M , positive numbers, where

$$|f(t, y)| < M, \quad \left| \frac{\partial^{i+j} f}{\partial t^i \partial y^j} \right| < \frac{P^{i+j}}{M^{j-1}}, \quad i + j \leq m$$

then in the 4th Runge-Kutta method $y(t_{i+1}) - y_{i+1} \approx \frac{73}{720} h^5 MP^4 + O(h^6)$.

Proof see [9].

A triangular fuzzy number v is defined by three numbers $a_1 < a_2 < a_3$ where the graph of $v(x)$, the membership function of the fuzzy number v , is a triangle with base on the interval $[a_1, a_3]$ and vertex at $x = a_2$.

We specify v as $(a_1/a_2/a_3)$. We will write: (1) $v > 0$ if $a_1 > 0$; (2) $v \geq 0$ if $a_1 \geq 0$; (3) $v < 0$ if $a_3 < 0$; and (4) $v \leq 0$ if $a_3 \leq 0$.

Let E be the set of all upper semicontinuous normal convex fuzzy numbers with bounded r -level intervals. It means that if $v \in E$ then the r -level set

$$[v]_r = \{s \mid v(s) \geq r\}, \quad 0 < r \leq 1,$$

is a closed bounded interval which is denoted by

$$[v]_r = [v_1(r), v_2(r)].$$

Let I be a real interval. A mapping $x : I \rightarrow E$ is called a fuzzy process and its r -level set is denoted by

$$[x(t)]_r = [x_1(t; r), x_2(t; r)], \quad t \in I, \quad r \in (0, 1].$$

Lemma 1 Let $v, w \in E$ and s scalar, then for $r \in (0, 1]$

$$[v + w]_r = [v_1(r) + w_1(r), v_2(r) + w_2(r)],$$

$$[v - w]_r = [v_1(r) - w_2(r), v_2(r) - w_1(r)],$$

$$[v \cdot w]_r = [\min\{v_1(r) \cdot w_1(r), v_1(r) \cdot w_2(r), v_2(r) \cdot w_1(r), v_2(r) \cdot w_2(r)\},$$

$$\max\{v_1(r) \cdot w_1(r), v_1(r) \cdot w_2(r), v_2(r) \cdot w_1(r), v_2(r) \cdot w_2(r)\}],$$

$$[sv]_r = s[v]_r.$$

3 A fuzzy cauchy problem

Consider the fuzzy initial value problem

$$\begin{cases} y'(t) = f(t, y(t)), & t \in I = [0, T], \\ y(0) = y_0, \end{cases} \quad (6)$$

where f is a continuous mapping from $R_+ \times R$ into R and $y_0 \in E$ with r -level intervals

$$[y_0]_r = [y_1(0; r), y_2(0; r)], \quad r \in (0, 1].$$

The extension principle of Zadeh leads to the following definition of $f(t, y)$ when $y = y(t)$ is a fuzzy number

$$f(t, y)(s) = \sup\{y(\tau) \mid s = f(t, \tau)\}, \quad s \in R.$$

It follows that

$$[f(t, y)]_r = [f_1(t, y; r), f_2(t, y; r)], \quad r \in (0, 1].$$

where

$$\begin{aligned} f_1(t, y; r) &= \min\{f(t, u) \mid u \in [y_1(r), y_2(r)]\}, \\ f_2(t, y; r) &= \max\{f(t, u) \mid u \in [y_1(r), y_2(r)]\}. \end{aligned} \quad (7)$$

Theorem 2 *Let f satisfy*

$$|f(t, v) - f(t, \bar{v})| \leq g(t, |v - \bar{v}|), \quad t \geq 0, \quad v, \bar{v} \in R,$$

where $g : R_+ \times R_+$ is a continuous mapping such that $r \rightarrow g(t, r)$ is nondecreasing, the initial value problem

$$u'(t) = g(t, u(t)), \quad u(0) = u_0, \quad (8)$$

has a solution on R_+ for $u_0 > 0$ and that $u(t) = 0$ is the only solution of (8) for $u_0 = 0$. Then the fuzzy initial value problem (6) has a unique fuzzy solution.

Proof see [7].

In this paper we suppose (6) satisfies the hypothesis of theorem 2, also.

4 4th Runge-Kutta method

Let the exact solution $[Y(t)]_r = [Y_1(t; r), Y_2(t; r)]$ is approximated by some $[y(t)]_r = [y_1(t; r), y_2(t; r)]$. From (2),(3) we define

$$\begin{aligned} y_1(t_{n+1}; r) - y_1(t_n; r) &= h \sum_{i=1}^4 w_i k_{i,1}(t_n, y(t_n; r)), \\ y_2(t_{n+1}; r) - y_2(t_n; r) &= h \sum_{i=1}^4 w_i k_{i,2}(t_n, y(t_n; r)), \end{aligned} \quad (9)$$

where the w_i s are constants and

$$\begin{aligned} [k_i(t, y(t; r))]_r &= [k_{i,1}(t, y(t; r)), k_{i,2}(t, y(t; r))], \quad i = 1, 2, 3, 4 \\ k_{i,1}(t_n, y(t_n; r)) &= f(t_n + \alpha_i h, y_1(t_n) + h \sum_{j=1}^{i-1} \beta_{ij} k_{j,1}(t_n, y(t_n; r))), \\ k_{i,2}(t_n, y(t_n; r)) &= f(t_n + \alpha_i h, y_2(t_n) + h \sum_{j=1}^{i-1} \beta_{ij} k_{j,2}(t_n, y(t_n; r))), \end{aligned} \quad (10)$$

and

$$\begin{aligned}
k_{1,1}(t, y(t; r)) &= \min\{f(t, u) | u \in [y_1(t; r), y_2(t; r)]\}, \\
k_{1,2}(t, y(t; r)) &= \max\{f(t, u) | u \in [y_1(t; r), y_2(t; r)]\}, \\
k_{2,1}(t, y(t; r)) &= \min\{f(t + \frac{h}{2}, u) | u \in [z_{1,1}(t, y(t; r)), z_{1,2}(t, y(t; r))]\}, \\
k_{2,2}(t, y(t; r)) &= \max\{f(t + \frac{h}{2}, u) | u \in [z_{1,1}(t, y(t; r)), z_{1,2}(t, y(t; r))]\}, \\
k_{3,1}(t, y(t; r)) &= \min\{f(t + \frac{h}{2}, u) | u \in [z_{2,1}(t, y(t; r)), z_{2,2}(t, y(t; r))]\}, \\
k_{3,2}(t, y(t; r)) &= \max\{f(t + \frac{h}{2}, u) | u \in [z_{2,1}(t, y(t; r)), z_{2,2}(t, y(t; r))]\}, \\
k_{4,1}(t, y(t; r)) &= \min\{f(t + h, u) | u \in [z_{3,1}(t, y(t; r)), z_{3,2}(t, y(t; r))]\}, \\
k_{4,2}(t, y(t; r)) &= \max\{f(t + h, u) | u \in [z_{3,1}(t, y(t; r)), z_{3,2}(t, y(t; r))]\}.
\end{aligned} \tag{11}$$

Where in the 4th Runge-Kutta method,

$$\begin{aligned}
z_{1,1}(t, y(t; r)) &= y_1(t; r) + \frac{h}{2}k_{1,1}(t, y(t; r)), & z_{1,2}(t, y(t; r)) &= y_2(t; r) + \frac{h}{2}k_{1,2}(t, y(t; r)), \\
z_{2,1}(t, y(t; r)) &= y_1(t; r) + \frac{h}{2}k_{2,1}(t, y(t; r)), & z_{2,2}(t, y(t; r)) &= y_2(t; r) + \frac{h}{2}k_{2,2}(t, y(t; r)), \\
z_{3,1}(t, y(t; r)) &= y_1(t; r) + hk_{3,1}(t, y(t; r)), & z_{3,2}(t, y(t; r)) &= y_2(t; r) + hk_{3,2}(t, y(t; r)).
\end{aligned} \tag{12}$$

Define,

$$\begin{aligned}
F[t, y(t; r)] &= k_{1,1}(t, y(t; r)) + 2k_{2,1}(t, y(t; r)) + 2k_{3,1}(t, y(t; r)) + k_{4,1}(t, y(t; r)), \\
G[t, y(t; r)] &= k_{1,2}(t, y(t; r)) + 2k_{2,2}(t, y(t; r)) + 2k_{3,2}(t, y(t; r)) + k_{4,2}(t, y(t; r)).
\end{aligned} \tag{13}$$

The exact and approximate solutions at $t_n, 0 \leq n \leq N$ are denoted by $[Y(t_n)]_r = [Y_1(t_n; r), Y_2(t_n; r)]$ and $[y(t_n)]_r = [y_1(t_n; r), y_2(t_n; r)]$ respectively. The solution is calculated by grid points at (5). By (9),(13) we have

$$\begin{aligned}
Y_1(t_{n+1}; r) &\approx Y_1(t_n; r) + \frac{h}{6}F[t_n, Y(t_n; r)], \\
Y_2(t_{n+1}; r) &\approx Y_2(t_n; r) + \frac{h}{6}G[t_n, Y(t_n; r)].
\end{aligned} \tag{14}$$

We define

$$\begin{aligned}
y_1(t_{n+1}; r) &= y_1(t_n; r) + \frac{h}{6}F[t_n, y(t_n; r)], \\
y_2(t_{n+1}; r) &= y_2(t_n; r) + \frac{h}{6}G[t_n, y(t_n; r)].
\end{aligned} \tag{15}$$

The following lemmas will be applied to show convergence of these approximates i.e.,

$$\begin{aligned}
\lim_{h \rightarrow 0} y_1(t; r) &= Y_1(t; r), \\
\lim_{h \rightarrow 0} y_2(t; r) &= Y_2(t; r).
\end{aligned}$$

Lemma 2 *Let a sequence of numbers $\{W_n\}_{n=0}^N$ satisfy*

$$|W_{n+1}| \leq A|W_n| + B, \quad 0 \leq n \leq N-1,$$

for some given positive constants A and B . Then

$$|W_n| \leq A^n|W_0| + B\frac{A^n - 1}{A - 1}, \quad 0 \leq n \leq N.$$

Proof see [10].

Lemma 3 *Let a sequence of numbers $\{W_n\}_{n=0}^N$, $\{V_n\}_{n=0}^N$ satisfy*

$$|W_{n+1}| \leq |W_n| + A.\max\{|W_n|, |V_n|\} + B,$$

$$|V_{n+1}| \leq |V_n| + A.\max\{|W_n|, |V_n|\} + B,$$

for some given positive constants A and B , and denote

$$U_n = |W_n| + |V_n|, \quad 0 \leq n \leq N.$$

Then

$$U_n \leq \bar{A}^n U_0 + \bar{B} \frac{\bar{A}^n - 1}{\bar{A} - 1}, \quad 0 \leq n \leq N,$$

where $\bar{A} = 1 + 2A$ and $\bar{B} = 2B$.

Proof see [10].

Let $F(t, u, v)$ and $G(t, u, v)$ are obtained by substituting $[y(t)]_r = [u, v]$ in (13),

$$F(t, u, v) = k_{1,1}(t, u, v) + 2k_{2,1}(t, u, v) + 2k_{3,1}(t, u, v) + k_{4,1}(t, u, v),$$

$$G(t, u, v) = k_{1,2}(t, u, v) + 2k_{2,2}(t, u, v) + 2k_{3,2}(t, u, v) + k_{4,2}(t, u, v).$$

The domain where F and G are defined is therefore

$$K = \{(t, u, v) | 0 \leq t \leq T, \quad -\infty < v < \infty, \quad -\infty < u \leq v\}.$$

Theorem 3 *Let $F(t, u, v)$ and $G(t, u, v)$ belong to $C^4(K)$ and let the partial derivatives of F and G be bounded over K . Then, for arbitrary fixed $r, 0 \leq r \leq 1$, the approximately solutions (14) converge to the exact solutions $Y_1(t; r)$ and $Y_2(t; r)$ uniformly in t .*

Proof : It is sufficient to show

$$\begin{aligned} \lim_{h \rightarrow 0} y_1(t_N; r) &= Y_1(t_N; r), \\ \lim_{h \rightarrow 0} y_2(t_N; r) &= Y_2(t_N; r), \end{aligned}$$

where $t_N = T$. For $n = 0, 1, \dots, N-1$, by using Taylor theorem we get

$$\begin{aligned} Y_1(t_{n+1}; r) &\approx Y_1(t_n; r) + \frac{h}{6}F[t_n, Y_1(t_n; r), Y_2(t_n; r)] + \frac{73}{720}h^5 MP^4 + O(h^6), \\ Y_2(t_{n+1}; r) &\approx Y_2(t_n; r) + \frac{h}{6}G[t_n, Y_1(t_n; r), Y_2(t_n; r)] + \frac{73}{720}h^5 MP^4 + O(h^6), \end{aligned} \quad (16)$$

denote

$$\begin{aligned} W_n &= Y_1(t_n; r) - y_1(t_n; r), \\ V_n &= Y_2(t_n; r) - y_2(t_n; r), \end{aligned}$$

we have from (14) and (15)

$$\begin{aligned} W_{n+1} &\approx W_n + \frac{h}{6}\{F[t_n, Y_1(t_n; r), Y_2(t_n; r)] - F[t_n, y_1(t_n; r), y_2(t_n; r)]\} + \frac{73}{720}h^5 MP^4 + O(h^6), \\ V_{n+1} &\approx V_n + \frac{h}{6}\{G[t_n, Y_1(t_n; r), Y_2(t_n; r)] - G[t_n, y_1(t_n; r), y_2(t_n; r)]\} + \frac{73}{720}h^5 MP^4 + O(h^6). \end{aligned}$$

Then

$$\begin{aligned} |W_{n+1}| &\leq |W_n| + \frac{1}{3}Lh \cdot \max\{|W_n|, |V_n|\} + \frac{73}{720}h^5 MP^4 + O(h^6), \\ |V_{n+1}| &\leq |V_n| + \frac{1}{3}Lh \cdot \max\{|W_n|, |V_n|\} + \frac{73}{720}h^5 MP^4 + O(h^6), \end{aligned}$$

for $t \in [0, T]$ and $L > 0$ is a bound for the partial derivatives of F and G . Thus by lemma 3

$$\begin{aligned} |W_n| &\leq (1 + \frac{2}{3}Lh)^n |U_0| + (\frac{73}{360}h^5 MP^4 + O(h^6)) \frac{(1 + \frac{2}{3}Lh)^n - 1}{\frac{2}{3}Lh}, \\ |V_n| &\leq (1 + \frac{2}{3}Lh)^n |U_0| + (\frac{73}{360}h^5 MP^4 + O(h^6)) \frac{(1 + \frac{2}{3}Lh)^n - 1}{\frac{2}{3}Lh}, \end{aligned}$$

where $|U_0| = |W_0| + |V_0|$. In particular

$$\begin{aligned} |W_N| &\leq (1 + \frac{2}{3}Lh)^N |U_0| + (\frac{73}{240}h^4 MP^4 + O(h^5)) \frac{(1 + \frac{2}{3}Lh)^{\frac{T}{h}} - 1}{L}, \\ |V_N| &\leq (1 + \frac{2}{3}Lh)^N |U_0| + (\frac{73}{240}h^4 MP^4 + O(h^5)) \frac{(1 + \frac{2}{3}Lh)^{\frac{T}{h}} - 1}{L}. \end{aligned}$$

Since $W_0 = V_0 = 0$, we obtain

$$\begin{aligned} |W_N| &\leq (\frac{73}{240}MP^4) \frac{e^{\frac{2}{3}LT} - 1}{L} h^4 + O(h^5), \\ |V_N| &\leq (\frac{73}{240}MP^4) \frac{e^{\frac{2}{3}LT} - 1}{L} h^4 + O(h^5), \end{aligned}$$

and if $h \rightarrow 0$ we get $W_N \rightarrow 0$ and $V_N \rightarrow 0$ which completes the proof.

5 Examples

Example 1 Consider the fuzzy initial value problem, [10],

$$\begin{cases} y'(t) = y(t), & t \in I = [0, 1], \\ y(0) = (.75 + .25r, 1.125 - .125r), & 0 < r \leq 1. \end{cases}$$

By using 4th Runge-Kutta method we have

$$y_1(t_{n+1}; r) = y_1(t_n; r) \left[1 + h + \frac{h^2}{2} + \frac{h^3}{6} + \frac{h^4}{24} \right],$$

$$y_2(t_{n+1}; r) = y_2(t_n; r) \left[1 + h + \frac{h^2}{2} + \frac{h^3}{6} + \frac{h^4}{24} \right].$$

The exact solution is given by

$$Y_1(t; r) = y_1(0; r)e^t, \quad Y_2(t; r) = y_2(0; r)e^t.$$

which at $t = 1$,

$$Y(1; r) = [(.75 + .25r)e, (1.125 - .125r)e], \quad 0 < r \leq 1.$$

The exact and approximate solutions we obtained by Euler method [10], and 4th Runge-Kutta method are compared and plotted at $t = 1$ in Fig.(5.1).




Fig.(5.1), ($h = .5$)

Example 2 Consider the fuzzy initial value problem, [10],

$$\begin{cases} y'(t) &= ty(t), \quad [a, b] = [-1, 1], \\ y(-1) &= (\sqrt{e} - .5(1-r), \sqrt{e} + .5(1-r)), \quad 0 < r \leq 1. \end{cases}$$

We separate between two steps.

(a) $t < 0$: The parametric form in this case is

$$y'_1(t; r) = ty_2(t; r), y'_2(t; r) = ty_1(t; r),$$

with the initial conditions given. The unique exact solution is

$$Y_2(t; r) = \frac{A+B}{2}y_2(0; r) + \frac{A-B}{2}y_1(0; r),$$

$$Y_1(t; r) = \frac{A-B}{2}y_2(0; r) + \frac{A+B}{2}y_1(0; r),$$

where $A = e^{\frac{(t^2-a^2)}{2}}$, $B = \frac{1}{A}$.

(b) $t \geq 0$: The parametric equations are

$$y'_1(t; r) = ty_1(t; r), y'_2(t; r) = ty_2(t; r),$$

with the initial conditions given. The unique exact solution at $t > 0$ is

$$Y_1(t; r) = y_1(0; r)e^{\frac{t^2}{2}}, Y_2(t; r) = y_2(0; r)e^{\frac{t^2}{2}}.$$

By using 4th Runge-Kutta method at $t_n, 0 \leq n \leq N$ we have

$$k_{1,1}(t_n; r) = \min\{t.u | u \in [y_1(t_n; r), y_2(t_n; r)]\},$$

$$k_{1,2}(t_n; r) = \max\{t.u | u \in [y_1(t_n; r), y_2(t_n; r)]\},$$

$$k_{2,1}(t_n; r) = \min\{(t + \frac{h}{2}).u | u \in [z_{1,1}(t_n; r), z_{1,2}(t_n; r)]\},$$

$$k_{2,2}(t_n; r) = \max\{(t + \frac{h}{2}).u | u \in [z_{1,1}(t_n; r), z_{1,2}(t_n; r)]\},$$

$$k_{3,1}(t_n; r) = \min\{(t + \frac{h}{2}).u | u \in [z_{2,1}(t_n; r), z_{2,2}(t_n; r)]\},$$

$$k_{3,2}(t_n; r) = \max\{(t + \frac{h}{2}).u | u \in [z_{2,1}(t_n; r), z_{2,2}(t_n; r)]\},$$

$$k_{4,1}(t_n; r) = \min\{(t + h).u | u \in [z_{3,1}(t_n; r), z_{3,2}(t_n; r)]\},$$

$$k_{4,2}(t_n; r) = \max\{(t + h).u | u \in [z_{3,1}(t_n; r), z_{3,2}(t_n; r)]\}.$$

Where

$$z_{1,1}(t_n; r) = y_1(t_n; r) + \frac{h}{2}k_{1,1}(t_n; r), \quad z_{1,2}(t_n; r) = y_2(t_n; r) + \frac{h}{2}k_{1,2}(t_n; r),$$

$$z_{2,1}(t_n; r) = y_1(t_n; r) + \frac{h}{2}k_{2,1}(t_n; r), \quad z_{2,2}(t_n; r) = y_2(t_n; r) + \frac{h}{2}k_{2,2}(t_n; r),$$

$$z_{3,1}(t_n; r) = y_1(t_n; r) + hk_{3,1}(t_n; r), \quad z_{3,2}(t_n; r) = y_2(t_n; r) + hk_{3,2}(t_n; r).$$

By considering $t > 0$ and $t < 0$, the above minimizing and maximizing problems can be solving by GAMS software. The exact and approximate solutions are compared and plotted in Fig.(5.2).




Fig.(5.2), ($h = .4$)

Example 3 Consider the fuzzy initial value problem

$$y'(t) = c_1 y^2(t) + c_2, \quad y(0) = 0,$$

where $c_i > 0$, for $i = 1, 2$ are triangular fuzzy numbers, [11].

The exact solution is given by

$$Y_1(t; r) = l_1(r) \tan(w_1(r)t),$$

$$Y_2(t; r) = l_2(r) \tan(w_2(r)t),$$

with

$$l_1(r) = \sqrt{c_{2,1}(r)/c_{1,1}(r)}, \quad l_2(r) = \sqrt{c_{2,2}(r)/c_{1,2}(r)},$$

$$w_1(r) = \sqrt{c_{1,1}(r) \cdot c_{2,1}(r)}, w_2(r) = \sqrt{c_{1,2}(r) \cdot c_{2,2}(r)},$$

where

$$[c_1]_r = [c_{1,1}(r), c_{1,2}(r)] \text{ and } [c_2]_r = [c_{2,1}(r), c_{2,2}(r)].$$

The r-level sets of $y'(t)$ are

$$Y'_1(t; r) = c_{2,1}(r) \sec^2(w_1(r)t),$$

$$Y'_2(t; r) = c_{2,2}(r) \sec^2(w_2(r)t),$$

which defines a fuzzy number. We have

$$f_1(t, y; r) = \min\{c_1 \cdot u^2 + c_2 | u \in [y_1(t; r), y_2(t; r)], c_1 \in [c_{1,1}(r), c_{1,2}(r)], c_2 \in [c_{2,1}(r), c_{2,2}(r)]\},$$

$$f_2(t, y; r) = \max\{c_1 \cdot u^2 + c_2 | u \in [y_1(t; r), y_2(t; r)], c_1 \in [c_{1,1}(r), c_{1,2}(r)], c_2 \in [c_{2,1}(r), c_{2,2}(r)]\},$$

By using Rung-Kutta method at $t_n, 0 \leq n \leq N$

$$k_{1,1}(t_n; r) = (c_{1,1}(r) \cdot y_1^2(t_n; r) + c_{2,1}(r))$$

$$k_{1,2}(t_n; r) = (c_{1,2}(r) \cdot y_2^2(t_n; r) + c_{2,2}(r))$$

$$k_{2,1}(t_n; r) = (c_{1,1}(r) \cdot z_{1,1}^2(t_n; r) + c_{2,1}(r))$$

$$k_{2,2}(t_n; r) = (c_{1,2}(r) \cdot z_{1,2}^2(t_n; r) + c_{2,2}(r))$$

$$k_{3,1}(t_n; r) = (c_{1,1}(r) \cdot z_{2,1}^2(t_n; r) + c_{2,1}(r))$$

$$k_{3,2}(t_n; r) = (c_{1,2}(r) \cdot z_{2,2}^2(t_n; r) + c_{2,2}(r))$$

$$k_{4,1}(t_n; r) = (c_{1,1}(r) \cdot z_{3,1}^2(t_n; r) + c_{2,1}(r))$$

$$k_{4,2}(t_n; r) = (c_{1,2}(r) \cdot z_{3,2}^2(t_n; r) + c_{2,2}(r))$$

Where

$$z_{1,1}(t_n; r) = y_1(t_n; r) + \frac{h}{2}k_{1,1}(t_n; r), \quad z_{1,2}(t_n; r) = y_2(t_n; r) + \frac{h}{2}k_{1,2}(t_n; r),$$

$$z_{2,1}(t_n; r) = y_1(t_n; r) + \frac{h}{2}k_{2,1}(t_n; r), \quad z_{2,2}(t_n; r) = y_2(t_n; r) + \frac{h}{2}k_{2,2}(t_n; r),$$

$$z_{3,1}(t_n; r) = y_1(t_n; r) + hk_{3,1}(t_n; r), \quad z_{3,2}(t_n; r) = y_2(t_n; r) + hk_{3,2}(t_n; r).$$

There are several nonlinear programings and can be solving by GAMS software. Thus the suggested 4th Rung-Kutta method in this paper can be used. The exact and approximate solutions are shown in Fig.(5.3) at $t = 1$.




Fig.(5.3), ($h = .5$)

6 Conclusion

We note that the convergence order of the Euler method in [10] is $O(h)$. It is shown that in proposed method, the convergence order is $O(h^4)$ and the comparison of solutions of example (1), (2) in this paper and [10] shows that this solutions are better for these examples.

References

- [1] S.L. Chang, L.A. Zadeh, On fuzzy mapping and control, *IEEE Trans, Systems Man Cybernet.* 2 (1972),30–34.
- [2] D.Dubois, H. Prade, Towards fuzzy differential calculus: Part 3, differentiation, *Fuzzy Sets and Systems* 8 (1982),225–233.
- [3] M.L Puri, D.A. Ralescu, Differentials of fuzzy functions, *J. Math. Anal. Appl.* 91 (1983),321–325.
- [4] R. Goetschel, W. Voxman, Elementary fuzzy calculus, *Fuzzy Sets and Systems* 18 (1986),31–43.
- [5] O. Kaleva, Fuzzy differential equations, *Fuzzy Sets and Systems* 24 (1987),301–317.

- [6] O. Kaleva, The Cauchy problem for Fuzzy differential equations, *Fuzzy Sets and Systems* 35 (1990), 389–396.
- [7] S. Seikkala, On the fuzzy initial value problem, *Fuzzy Sets and Systems* 24 (1987) 319–330.
- [8] R. L. Burden, J. D. Faires, Numerical Analysis, 1997.
- [9] A. Ralston, P. Rabinowitz, First Course In Numerical Analysis, 1978.
- [10] M. Ma, M. Friedman, A. Kandel, Numerical Solutions of fuzzy differential equations, *Fuzzy Sets and Systems* 105 (1999), 133–138.
- [11] James J. Buckley, Thomas Feuring, Fuzzy differential equations, *Fuzzy Sets and Systems* 110 (2000), 43–54.