

University of Texas at El Paso

From the SelectedWorks of Theodore V. Cooper

2009

Obesity-relevant behaviors: Patterns and correlates in a Hispanic college sample.

D. Hu

T. J. Taylor

Theodore V. Cooper, *University of Texas at El Paso*



SELECTEDWORKS™

Available at: http://works.bepress.com/theodore_v_cooper/14/

Running Head: OBESITY-RELEVANT BEHAVIORS

5

10 Obesity-Relevant Behaviors: Patterns and Correlates in a Hispanic College Sample

Dixie Hu, B.S., Thom Taylor, B.A.S., & Theodore V. Cooper, Ph.D.

The University of Texas at El Paso

El Paso, TX 79968 USA

15

Manuscript submitted for publication

20

Corresponding Author:

Theodore V. Cooper, Ph.D.
Department of Psychology
University of Texas at El Paso
500 West University Avenue
El Paso, TX 79968
(915) 747-6270
tvcooper@utep.edu

25

30

35

Abstract*Objective*

Obesity rates in Hispanic groups and the role of lifestyle factors warrant the investigation of multiple health behaviors for prevention and intervention aims. Patterns and correlates of exercise and dietary behaviors were examined in Hispanic college students.

40 *Participants and Methods*

Data were collected from 693 Hispanic undergraduate students during fall 2006 and spring 2007 semesters. Participants completed questionnaires for three health behaviors (exercise, dietary fat, and fruit/vegetable stages of change) along with demographic, psychosocial, and acculturation measures.

45 *Results*

Less than 1% of students had 0 obesity-relevant risks, while 68% indicated 2 or more risks. Only 2% of the sample met fruit and vegetable guidelines. Lower income was associated with greater obesity-relevant risks, while stress coping ability was associated with fewer such risks.

Conclusions

50 These findings indicate high levels of specific obesity risk behaviors in Hispanic college students and suggest specific demographic and psychosocial targets for prevention and intervention.

Keywords: Obesity, Hispanic, multiple health behaviors, exercise, diet

55

Obesity-Relevant Behaviors: Patterns and Correlates in a Hispanic College Sample

Over two-thirds of Americans (70.6%) are either overweight or obese, while one in three adolescents are overweight or at risk of overweight.¹ Clinical guidelines define adult overweight and obesity as a body mass index (BMI) of 25–29.9 kg/m² and ≥ 30 kg/m², respectively.² Obesity is a serious health concern as it is associated with a number of preventable or reversible medical complications, including type 2 diabetes,³ hypertension,⁴ elevated cholesterol levels,⁵ and coronary heart disease.^{6,7}

While obesity rates are increasing across all ethnic groups,⁸ certain groups are at greater risk.¹ Hispanics have demonstrated the greatest growth in obesity prevalence in the United States—an 80% increase between 1991 and 1998.⁸ In particular, Mexican American adolescents are at greater risk of obesity than non-Hispanic White and non-Hispanic Black adolescents.⁹ Data from the National Health and Nutrition Examination Survey from 1992-2002 indicate that over 40% of Mexican American adolescents are overweight, and over 22% are obese.¹⁰ Studies assessing the correlates of obesity in Hispanic subgroups have implicated low physical activity,¹¹ poor diet,^{12,13} acculturation,^{14,15} and income and education levels.^{10,11,16} Perceived stress and current smoking have also been associated with unhealthy eating behaviors in Hispanics¹⁷ and college students,¹⁸ meriting the investigation of these variables as potential correlates of obesity-promoting behaviors in Hispanic groups.

Given that the highest increases in overweight and obesity occur between 18 and 29 years of age,⁸ an examination of activity and eating behaviors is warranted for the development of prevention and intervention efforts. College is a salient transition period for health behavior change in adolescents and young adults.^{19,20} In fact, only 21.4% of Hispanic college students

nationwide regularly engage in moderate physical activity, while less than one-third (31.5%) consume at least five fruits and vegetables daily.²¹

80 Emergent research suggests that these unhealthy behaviors frequently co-occur, with potentially multiplicative health consequences.²²⁻²⁵ Berrigan and colleagues²² note that the most common pattern of health behaviors in U.S. adults involves a lack of adherence to exercise, dietary fat, and fruit and vegetable recommendations. To that end, the rationale underlying research on multiple health behavior change (MHBC) as opposed to single behavior foci is one
85 of maximizing health benefits and reducing healthcare costs.²⁶

The transtheoretical model (TTM) is one avenue for tailoring MHBC interventions, as it addresses individual psychosocial and motivational mediators of behavior change.²⁷ The TTM conceptualizes temporal behavior change along a cyclical continuum of readiness divided into five stages: precontemplation, contemplation, preparation, action, and maintenance.²⁸ In
90 *precontemplation*, individuals are not considering behavior change within the next six months, while individuals in the *contemplation* stage plan to alter behavior within six months; *preparation* individuals have either taken moderate action or intend to act within one month. Adequate behavior modifications from zero to six months comprise the *action* stage, while the *maintenance* stage entails consistent changed behavior for more than six months.²⁸

95 TTM constructs have been successfully applied to a number of single health behaviors, including smoking,^{29, 30} physical activity,^{23, 31, 32} and diet.³³ The model's efficacy has been demonstrated in two MHBC interventions on smoking, dietary fat, and sun exposure,^{34,35} as well as interventions on physical activity and dietary fat reduction.^{36, 37} Yet, the paucity of TTM literature addressing MHBC for ethnic college groups suggests the need for further empirical
100 study.

Given the emerging field of MHBC, evidence of multiple behavior intervention efficacy in college students is limited. One pilot study demonstrated significant increases in physical activity and consumption of healthy fats in college students after a brief image-based intervention,³⁸ though similar studies in Hispanic college students have yet to be performed.

105 As such, examining patterns and correlates of physical activity and dietary behaviors using TTM constructs in this cohort serves to critically inform the design and planning of future MHBC interventions in the context of obesity risk. These comprise current study aims. Specifically, this study examined the extent of risk in lack of adherence to physical activity, dietary fat, and 5-A-Day fruit and vegetable recommendations, as well as correlates of risk in
110 Hispanic college students. Relevant correlates include sociodemographic (age, gender, income, acculturation), health-related (smoking status, BMI), and psychosocial (perceived stress, coping ability) constructs. Individuals in this study were hypothesized to report being in earlier stages of change (i.e. not meeting recommendations) for the three behaviors; risk was hypothesized to correlate with high acculturation, low income, high BMI, smoking, and greater perceived stress.

115 **Methods**

Participants

Data were collected in 2006-2007 from 693 Hispanic undergraduate students at least 18 years of age in the U.S.–México border region. University undergraduate Hispanic enrollment at the time of data collection (76.1%) nearly perfectly mirrored the Hispanic population in the region
120 (76.6%).³⁹ Students were 59% female, with an average age of 19.6 years ($SD=3.33$ years).

Design and Procedure

Cross-sectional data were collected after Institutional Review Board approval. Participants provided informed consent and completed seven paper-and-pencil questionnaires assessing

readiness outcomes on three obesity-relevant behaviors, as well as demographic, health related,
125 and stress and coping variables. Students were debriefed and given course credit for voluntary
participation.

Measures

Potential Correlates

Age, gender, annual household income, household size (number of household members), and
130 self-reported height and weight (for BMI calculation) were obtained. Annual household income
was divided into four distinct groups: Less than US \$15,000, between \$15,000 and \$30,000,
between \$30,000 and \$50,000, and more than \$50,000. Household size was included to account
for low income scores from single-living students.

The *Rhode Island Stress and Coping Inventory* (RISCI) evaluates perceived stress and coping
135 ability using 12 items.⁴⁰ Respondents rate the frequency of items (e.g. “I had no time to relax”)
within the past month from 1 (Never) to 5 (Frequently). Adequate reliability, internal
consistency, and content validity have been demonstrated, as well as high correlations between
the RISCI subscales and the five-item Mental Health Inventory.⁴⁰ Scores for the stress subscale
range from 7 to 35 (higher scores indicate greater perceived stress), while the coping ability
140 subscale ranges from 5 to 25 (higher scores indicate greater coping ability). Adequate reliability
was observed in both the 7 item stress subscale ($\alpha = .82$) and the 5 item coping ability subscale
($\alpha = .81$).

The *Short Acculturation Scale for Hispanics* (SASH)⁴¹ assesses the degree to which
Hispanics have adopted American culture by measuring three aspects of acculturation: language,
145 media, and ethnic social relations. Acculturation scores range from 1 through 5, with scores
averaging above 2.99 indicative of more acculturated respondents. Adequate psychometric

properties have been demonstrated for this 12 item measure,⁴¹ and internal consistency was demonstrated in this sample ($\alpha = .91$).

Smoking status was assessed using the *Stage of Change for Smoking Cessation (Short Form)*, based on a six-item algorithm measuring readiness to quit smoking along five stages—precontemplation, contemplation, preparation, action, and maintenance.²⁹

Obesity-Relevant Outcome Variables

Health behaviors associated with obesity risk were examined. Specifically, recommendations for regular exercise, dietary fat reduction, and fruit and vegetable consumption were examined using a staging algorithm, with low stage of change (SOC) scores suggestive of failure to meet recommendations.

The *Stage of Change for Exercise (Short Form)* places respondents into one of five stages of readiness to exercise based on a single item response to a specific definition: “Regular Exercise is any *planned* physical activity (e.g., brisk walking, aerobics, jogging, bicycling, swimming, rowing, etc.) performed to increase physical fitness. Such activity should be performed 3 to 5 times per week for 20-60 minutes per session. Exercise does not have to be painful to be effective but should be done at a level that increases your breathing rate and causes you to break a sweat. Do you exercise regularly according to that definition?”⁴²

The *Stage of Change (Dietary Fat)* evaluates intentions to reduce dietary fat using three steps: the first assesses stage classification based on respondents’ intentions or actions taken to “consistently avoid eating high fat foods.”⁴³ The second step confirms respondents’ perceptions of dietary fat avoidance with actual behavior in 5 items (e.g. “Do you often use light, fat free, or no salad dressing?”), while the final step assigns participants whose perceptions do not match behavior into precontemplation, contemplation, or preparation.⁴³

170 The *Stage of Change (5 A Day)* applies two items to classify stage of change for fruit and
vegetable consumption: the first item assesses the number of fruit and vegetable servings
consumed per day.⁴⁴ The second item evaluates stage of change, in which a response of fewer
than five servings is assigned to precontemplation, contemplation, or preparation. Responses of
five or more servings are assigned to action or maintenance on a temporal basis.⁴⁴

175 *Statistical Analyses*

Stage distributions were reported for each behavior to provide initial insight into TTM
motivational constructs. Above all, there were two primary aims for analyses: to examine the
extent of obesity risk in lack of adherence to multiple health behavior recommendations, and to
identify demographic and other correlates of this increased risk. For the first objective, SOC for
180 each behavior was dichotomized from a 5-stage algorithm to represent obesity risk. Those who
reported precontemplation, contemplation, or preparation stages failed to perform or to meet
recommendations for the desired behavior and were categorized as being “at risk” (1), while
individuals reporting action or maintenance stages were categorized as “not at risk” (0). These
variables were then summed to create a composite score ranging from 0 to 3, with higher scores
185 representing increased risk for obesity from dietary and exercise behaviors.

SOC was dichotomized because the proportional odds assumption in the 5 stages could not
be met for ordinal regression approaches. In the second aim for analyses, correlates of increased
risk (represented by the composite score) were examined using ordinal logistic regression.⁴⁵
Predictor variables of interest were age, gender, household income, SASH acculturation score,
190 BMI, smoking status, as well as the RISC stress and coping scores. All analyses were conducted
using STATA 10.1.⁴⁶

Results

Participant characteristics and stage distributions are reported in Table 1 and Figure 1, respectively. Roughly 32% of participants were either overweight or obese. Examining behaviors singly, regular exercise was the only behavior for which more than half of the sample reported no risk. Otherwise, the majority of participants were at risk for lack of adherence to dietary fat and fruit and vegetable recommendations. Notably, only 2% of the sample reported consuming at least 5 fruits and vegetables daily. Consequently, only 1% of this Hispanic sample had 0 obesity related behavioral risks, while most participants (68% of the sample) reported 2 or more risks (see Table 1).

Because of the limited number of individuals reporting 0 health risk behaviors, these individuals were removed from inferential analysis for model stability. The ordinal regression met the proportional odds assumption, $\chi^2(10) = 13.17, p > .22$; it revealed a significant model with three significant predictors of increased risk, $\chi^2(10) = 27.39, p < .01$. Odds ratios (*ORs*) and confidence intervals (*CI*s) are reported in Table 2. Holding other factors constant, those who reported household incomes between \$30k-\$50k relative to those with household incomes greater than \$50k were marginally more likely to report an increase in risk factors ($OR = 1.48, p = .07$). However, relative to those who reported annual household incomes above \$50k, those with incomes of \$15k-\$30k were significantly more likely to report more behavioral risks for obesity ($OR = 1.93, p < .01$). While the log odds coefficients for those in the \$30k-\$50k and \$15k-\$30k income groups did not statistically differ from one another, $\chi^2(1) = 1.58, p > .21$, the observed larger *ORs* and associated coefficients for the lowest household income categories assessed in this study likely indicates a monotonic negative association between income and obesity risk for Hispanic college students. Notably, increased coping ability was associated with

fewer risk behaviors reported ($OR = .95, p < .05$). Finally, age was marginally associated with an increased odds of reporting an additional risk factor for obesity ($OR = 1.05, p = .06$; see Table 2).

Comment

Multiple obesity-relevant behaviors (exercise, dietary fat consumption, and fruit/vegetable
220 intake) and correlates were examined in a Hispanic college student sample. Virtually none of the
sample adhered to recommendations for all three behaviors, while the majority of students failed
to follow guidelines for two or three behaviors concurrently. Results are consistent with findings
across adult population groups that suggest the frequent occurrence of multiple risk
behaviors.^{22,24} Lack of sufficient fruit and vegetable intake comprised the most common
225 behavioral risk, in which 98% of the sample failed to consume at least five fruits and vegetables
daily. This finding is higher than the 73% of college students nationwide who do not meet this
dietary recommendation.²¹ Higher fruit and vegetable consumption among college students has
been associated with greater levels of physical activity⁴⁷ and inversely correlated with high-fat
fried food and fast food intake⁴⁸—salient associations in the context of obesity risk and nontrivial
230 weight gain during the first two years of college.⁴⁸ Health education programs and interventions
are thus encouraged to incorporate nutritional components to promote fruit and vegetable intake
in Hispanic students.

Over half of students in this study did not make efforts to reduce dietary fat, while 42% of
individuals reported being in action or maintenance stages for this behavior. Evidence from the
235 only published national college health risk behavior survey suggests that a majority of students
already consume low levels of dietary fat (i.e., fewer than 2 servings of high fat foods daily).²¹
Thus, it may be that students in this sample were in fact eating or perceiving low levels of fat
intake, with little or no intention for reduction. Alternatively, activity and fruit and vegetable

recommendations may be relatively well known to students, while dietary fat is a more recently
240 emerging weight related variable. College students with increased knowledge of dietary
guidelines are more likely to adopt healthy eating patterns.⁴⁹ As such, dietary fat may be an ideal
target for obesity prevention and intervention psychoeducation efforts.

Contrary to predictions, an encouraging majority in this sample reported exercising regularly.
These findings are in line with results from one study in which 75% of US college students
245 reported engaging in some form of physical activity.⁵⁰ In one cluster analysis, Reedy and
colleagues⁵¹ noted that individuals in a “Physically Active” cluster were more likely to increase
fruit and vegetable consumption as a result of a tailored intervention. This implicates exercise as
a gateway behavior for dietary change.⁵² Given that the majority of students in our sample
reported action or maintenance stages for exercise, as well as contemplation or preparation stages
250 for fruit and vegetable intake, MHBC interventions that combine activity and dietary based
components—particularly as they address self-efficacy and other TTM constructs associated
with these stages—may promote healthy eating and exercise behaviors in Hispanic college
students.

Correlates of Multiple Risks

255 Lower socioeconomic status has been widely implicated with multiple risk behaviors in adult
populations,^{22, 24, 25, 53, 54} and with overweight, obesity, and class II obesity in college students.⁵⁰
In one study, Hispanic females with low education were more likely to have a desired BMI in the
overweight category,⁵⁵ while a separate study found that adolescents with higher socioeconomic
status had greater weight awareness and were more likely to adopt healthy weight management
260 methods.⁵⁶ Though education level in this study was uniform and high, lower income was
nonetheless associated with obesity-relevant risks. Prevention and intervention strategies are thus

encouraged to address healthy weight attitudes, awareness, and management in Hispanic college students with lower income or other socioeconomic indicators. In addition, food environments play a role in students' behavioral and financial choices.^{57, 58} As population and policy efforts have targeted children's and adolescents' dietary intake in schools,^{59, 60} similar efforts that address availability and cost-containment of healthy foods are encouraged in college settings.

Contrary to hypotheses, perceived stress was not associated with multiple behavioral risks in this study. Higher levels of perceived stress have been associated with weight changes during the first year in college,⁶¹ as well as poor health outcomes in Hispanics.⁶² While stress scores were moderate in this cohort, results suggest that Hispanic students confronted with stressors may resort less to unhealthy behaviors (e.g. consumption of calorie-rich foods) than to other outlet sources, such as exercise or family support.⁶³

Additionally, students who reported higher stress coping ability reported fewer risk behaviors. An increased ability to cope with daily life stressors (using a cognitive/emotional management coping style) has been associated with better physical functioning in Mexican Americans and Mexican immigrants.⁶² Further, Chiang and colleagues⁶³ identified exercise, hobbies, and activities with family as the most frequently cited coping practices among Latino college students. Current and prior findings suggest the examination of specific coping styles (e.g. positive reframing, social support, and acceptance) and their impact on health outcomes to complement other multiple behavior change intervention components.

Greater cigarette consumption has been associated with increased behavioral risks in adult populations.⁶⁴ Among college students, current smoking is correlated with more frequent consumption of highly caloric meals at restaurants, less frequent use of exercise facilities, and eating in front of a television.⁶⁵ In contrast, smoking in this study was not significantly

285 associated with multiple risk behaviors. This may be related to the limited number of smokers in
the study, and thus reduced power, or the predominantly light smoking nature of our sample and
Hispanic college students more generally.⁶⁶ That is, low levels of smoking may be less clearly
associated with other deleterious health risks.

With regard to Hispanic acculturation and health risks, evidence is mixed in the broad
290 literature. On the one hand, higher acculturation has been implicated with obesity^{13, 14, 16} and
increased risks such as physical inactivity,¹⁵ poor diet,¹² and poor self-rated health.⁶⁷ Still, a
number of studies generate contrary or no evidence of the impact of acculturation.^{11, 68} The
current results are consistent with the latter interpretation in that acculturation may be less
strongly related to multiple obesity relevant behaviors than to other salient influences.

295 Socioeconomic indicators, for instance, may be more useful in understanding health behavior
change,⁶⁹ and may be more fitting in the context of intervention design and planning.

Though roughly one third of the sample was either overweight or obese, BMI was not
associated with increased behavioral risks. Similarly, a longitudinal study of first and second
year college students found no associations between changes in BMI and changes in exercise and
300 dietary behaviors.⁴⁸ Three potential justifications emerge for the current finding. First, young age
and other sample characteristics may underlie the lack of association for BMI and multiple risk
behaviors. However, that high BMI is evidenced in younger age groups weakens this
explanation. Second, the energy balance equation of caloric intake versus expenditure may
account for this finding. That is, the majority of individuals met exercise recommendations;
305 however, amounts and intensities of exercise were not assessed. It is possible that students
performed exercise at levels that offset the energy balance far in the desired direction, buffering
against dietary risks and contributing to observed BMI. Finally, the behavioral measures may not

have captured culturally relevant or detailed estimates of dietary intake or physical activity. For instance, individuals with no intention to reduce dietary fat may have answered “No” to the question, “Do you often eat bread, rolls, or muffins without butter or margarine?” for the reason that these items were not staples in their diets. Carrera and colleagues⁷⁰ have identified a “traditional Mexican diet” cluster in which individuals consumed more fruits, vegetables, and fiber (likely from tortillas and legumes) as well as cholesterol and total energy. To that end, more precise estimates of dietary intake may be achieved through food frequency questionnaires, though these were beyond the aims and scope of this study.

Limitations

Results of the study warrant careful interpretation for several reasons. A reliance on self-report may lead to under- or overestimations of BMI or socially desirable health behaviors; however, self-reported height and weight correlate highly with measured BMI in young adults.⁷¹ Obesity-promoting behaviors such as soft-drink consumption or television viewing were not examined and may further inform MHBC intervention planning in Hispanic college groups. As noted, detailed estimates of exercise and dietary behaviors were not assessed, such as percentage of dietary fat from daily caloric intake, types of fats consumed, or types and duration of exercise performed.

Several strengths of the study merit mentioning. Few studies have examined multiple health behaviors in Hispanic college students, one of the fastest growing ethnic minorities entering college.⁷² Further, the use of a large, fairly homogenous sample of individuals of Mexican descent enhances generalizability for use by researchers and healthcare providers, perhaps particularly those working within the growing number of Hispanic Serving Institutions.

Conclusions

This study provides an examination of multiple obesity-relevant behaviors in Hispanic college students. Reports of regular exercise and poor nutrition with income and stress coping associations highlight unique patterns and correlates of behaviors contributing to obesity risk.

335 Maximizing health benefits and reducing healthcare costs are two of the aims that drive multiple health behavior change research and are particularly relevant for college health professionals implementing prevention and intervention programs. Future directions include longitudinal studies of multiple health behavior change and the development, implementation, and assessment of interventions tailored to underserved ethnocultural groups.

340

Acknowledgements

This research was supported by NIMH-COR Grant: T34 MH019978-05 and portions of the work were funded by A Smoke Free Paso del Norte, an initiative of the Paso del Norte Health
345 Foundation.

The authors would like to acknowledge the work and support of members of the Prevention and Treatment in Clinical Health research laboratory.

350

References

1. Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of overweight and obesity in the United States, 1999-2004. *JAMA*. 2006;295:1549-1555.
2. National Heart, Lung, and Blood Institute. Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults: The evidence report. NIH Publication No. 98-4083;1998.
3. Kopelman, P. Health risks associated with overweight and obesity. *Obes Rev*. 2007;8(Suppl 1):13-17.
4. Huang Z, Willett WC, Manson JE, et al. Body weight, weight change, and risk for hypertension in women. *Ann Intern Med*. 1998;128:81-88.
- 360 5. Paramsothy P, Knopp R, Bertoni AG, Tsai MY, Rue T, Heckbert SR. Combined hyperlipidemia in relation to race/ethnicity, obesity, and insulin resistance in the Multi-Ethnic Study of Atherosclerosis. *Metabolism*. 2009;58:2121-219.
6. Yusuf S, Hawken S, Ôunpuu S, et al. Obesity and the risk of myocardial infarction in 27000 participants from 52 countries. *Lancet*. 2005;366:1640-1649.
- 365 7. Gray RS, Fabsitz RR, Cowan LD, et al. Relation of generalized and central obesity to cardiovascular risk factors and prevalent coronary heart disease in a sample of American Indians: The Strong Heart Study. *Int J Obes*. 2000;24:849-860.
8. Mokdad AH, Serdula MK, Dietz WH, Bowman BA, Marks JS, Koplan JP. The spread of the obesity epidemic in the United States, 1991-1998. *JAMA*. 1999;282:1519-1522.
- 370 9. Hedley AA, Ogden CL, Johnson CL, Carroll MD, Curtin LR, Flegal KM. Prevalence of overweight and obesity among US children, adolescents, and adults, 1999-2002. *JAMA*. 2004;291:2847-2850.

10. Forrest KY, Leeds MJ. Prevalence and associated factors of overweight among Mexican-American adolescents. *J Am Diet Assoc.* 2007;107:1797-1800.
- 375 11. Slattery ML, Sweeney C, Edwards S, et al. Physical activity patterns and obesity in Hispanic and non-Hispanic White women. *Med Sci Sports Exerc.* 2006;38:33-41.
12. Allen ML., Elliott MN, Morales LS, Diamant, AL, Hambarsoomian, K, Schuster, MA. Adolescent participation in preventive health behaviors, physical activity, and nutrition: Differences across immigrant generations for Asians and Latinos compared with Whites. *Am J*
- 380 *Public Health.* 2007;97:337-343.
13. Lin H, Bermudez OI, Tucker KL. Dietary patterns of Hispanic elders are associated with acculturation and obesity. *J Nutr.* 2003;133:3651-3657.
14. Hubert HB, Snider J, Winkleby, MA. Health status, health behaviors, and acculturation factors associated with overweight and obesity in Latinos from a community and agricultural
- 385 labor camp survey. *Prev Med.* 2005;40:642-651.
15. Unger JB, Reynolds K, Shakib S, Spruijitt-Metz D, Sun P, Johnson A. Acculturation, physical activity, and fast-food consumption among Asian-American and Hispanic adolescents. *J Community Health.* 2004;29:467-481.
16. Bowie JV, Juon HS, Cho J, Rodriguez EM. Factors associated with overweight and obesity
- 390 among Mexican Americans and Central Americans: Results from the 2001 California Health Interview Survey. *Prev Chronic Dis.* 2007;4:A10.
17. Jenkins SK, Rew L, Sternglanz RW. Eating behaviors among school-age children associated with perceptions of stress. *Issues Compr Pediatr Nurs.* 2005;28:175-191.
18. Carroll SL, Lee RE, Kaur H, Harris KJ, Strother ML, Huang TTK. Smoking, weight loss
- 395 intention and obesity-promoting behaviors in college students. *J Am Coll Nutr.* 2006;25:348-353.

19. Baronowski T, Cullen KW, Basen-Engquist K, et al. Transitions out of high school: Time of increased cancer risk? *Prev Med.* 1997;26:694-703.
20. Wong Y, Huang YC, Chen SL, Yamamoto S. Is the college environment adequate for accessing to nutrition education: A study in Taiwan. *Nutr Res.* 1999;19:1327-1337.
- 400 21. Centers for Disease Control and Prevention. Youth risk behavior surveillance: National college health risk behavior survey—United States, 1995. *MMWR CDC Surveill Summ.* 1997;46:1-54.
22. Berrigan D, Dodd K, Troiano RP, Krebs-Smith SM, Barbash RB. Patterns of health behavior in U.S. adults. *Prev Med.* 2003;36:615-623.
- 405 23. Emmons KM, Marcus BH, Linnan L, Rossi JS, Abrams DB. Mechanisms in multiple risk factor interventions: Smoking, physical activity, and dietary fat intake among manufacturing workers. *Prev Med.* 1994;23:481-489.
24. Fine LJ, Philogene S, Gramling R., Coups, EJ, Sinha S. Prevalence of multiple chronic disease risk factors: 2001 National Health Interview Survey. *Am J Prev Med.* 2004; 27:18-24.
- 410 25. Pronk NP, Anderson LH, Crain AL, et al. Meeting recommendations for multiple healthy lifestyle factors: Prevalence, clustering, and predictors among adolescent, adult, and senior health plan members. *Am J Prev Med.* 2004;27:25-33.
26. Prochaska JJ, Spring B, Nigg CR. Multiple health behavior change research: An introduction and overview. *Prev Med.* 2008;46:181-188.
- 415 27. Prochaska JO, DiClemente CC. Stages and processes of self-change of smoking: Toward an integrative model of change. *J Consult Clin Psychol.* 1983;51:390-395.

28. Velicer WF, Prochaska JO, Fava JL, Norman GJ, Redding CA. Smoking cessation and stress management: Applications of the transtheoretical model of behavior change. *Homeost Health Dis.* 1998;38:216-233.
- 420 29. DiClemente CC, Prochaska JO, Fairhurst SK, Velicer WF, Velasquez MM, Rossi JS. The process of smoking cessation: An analysis of precontemplation, contemplation, and preparation stages of change. *J Consult Clin Psychol.* 1991;59:295-304.
30. Prochaska JO, Velicer WF, Rossi JS, et al. Stages of change and decisional balance for 12 problem behaviors. *Health Psychol.* 1994;13:39-46.
- 425 31. King TK, Marcus BH, Pinto BM, Emmons KM, Abrams DB. Cognitive-behavioral mediators of changing multiple behaviors: Smoking and sedentary lifestyle. *Prev Med.* 1996;25:684-691.
32. Marshall SJ, Biddle SJH. The transtheoretical model of behavior change: A meta-analysis of applications to physical activity and exercise. *Ann Behav Med.* 2001;23:229-246.
- 430 33. Park A, Nitzke S, Kritsch K, et al. Internet-based interventions have potential to affect short-term mediators and indicators of dietary behavior of young adults. *J Nutr Educ Behav.* 2008;40:288-297.
34. Prochaska JO, Velicer WF, Rossi JS, et al. Multiple risk expert systems interventions: impact of simultaneous stage-matched expert system interventions for smoking, high-fat diet, and sun
435 exposure in a population of parents. *Health Psychol.* 2004;23:503-516.
35. Prochaska JO, Velicer WF, Redding C, et al. Stage-based expert systems to guide a population of primary care patients to quit smoking, eat healthier, prevent skin cancer, and receive regular mammograms. *Prev Med.* 2005;41:406-416.

36. Clark M, Hampson SE, Avery L, Simpson R. Effects of a tailored lifestyle self-management
440 intervention in patients with type 2 diabetes. *Br J Health Psychol.* 2004;9:365-379.
37. Johnson SS, Paiva AL, Cummins CO, et al. Transtheoretical model-based multiple behavior
intervention for weight management: effectiveness on a population basis. *Prev Med.*
2008;46:238-246.
38. Werch CE, Bian H, Moore MJ, Ames S, DiClemente CC, Weiler RM. Brief multiple
445 behavior interventions in a college student health care clinic. *J Adolesc Health.* 2007;41:577-
585.
39. US Census Bureau. The Hispanic population: Census 2000 brief. Available at:
<http://factfinder.census.gov>. Accessed January 7, 2009.
40. Fava JL, Ruggiero L, Grimley DM. The development and structural confirmation of the
450 Rhode Island Stress and Coping Inventory. *J Behav Med.* 1998;21:601-611.
41. Marín G, Sabogal F, Marín BV, Otero-Sabogal R, Perez-Stable EJ. Development of a short
acculturation scale for Hispanics. *Hispanic J Behav Sci.* 1987;9:183-205.
42. Marcus BH, Selby VC, Niaura RS, Rossi JS. Self-efficacy and the stages of exercise
behavior change. *Res Q Exerc Sport.* 1992;63:60-66.
- 455 43. Greene GW, Rossi SR, Rossi JS, Velicer WF, Fava JL, Prochaska JO. Dietary applications of
the stages of change model. *J Am Diet Assoc.* 1999;99:673-678.
44. Vallis M, Ruggiero L, Greene G, et al. Stages of change for healthy eating in diabetes:
Relation to demographic, eating-related, health care utilization, and psychosocial factors.
Diabetes Care. 2003;26:1468-1474.
- 460 45. Hamilton LS. *Statistics with STATA*. Belmont, CA: Thomson Higher Education; 2006.
46. StataCorp. *STATA 10 for Windows* [Computer Software]. College Station, TX: Author; 2007.

47. Adams TB, Colner W. The association of multiple risk factors with fruit and vegetable intake among a nationwide sample of college students. *J Am Coll Health*. 2008;56:455-461.
48. Racette SB, Deusinger SS, Strube MJ, Highstein GR, Deusinger RH. Weight changes, exercise, and dietary patterns during freshman and sophomore years of college. *J Am Coll Health*. 2005;53:245-251.
49. Kolodinsky J, Harvey-Berino JR, Berlin L, Johnson RK, Reynolds TW. Knowledge of current dietary guidelines and food choice by college students: Better eaters have higher knowledge of dietary guidance. *J Am Diet Assoc*. 2007;107:1409-1413.
- 470 50. Nelson TF, Gortmaker SL, Subramanian SV, Cheung L, Wechsler H. Disparities in overweight and obesity among US college students. *Am J Health Behavior*. 2007;31:363-373.
51. Reedy J, Haines PS, Campbell MK. The influence of health behavior clusters on dietary change. *Prev Med*. 2005;41:268-275.
52. Tucker M, Reicks M. Exercise as a gateway behavior for healthful eating among older adults: An exploratory study. *J Nutr Educ Behav*. 2002;34 Suppl. 1: S14-S19.
- 475 53. Poortinga W. The prevalence and clustering of four major lifestyle risk factors in an adult English population. *Prev Med*. 2007;44:124-128.
54. Galán I, Rodríguez-Artalejo F, Tobías A, Díez-Gañán L, Gandarillas A, Zorrilla B. Agregación de factores de riesgo ligados al comportamiento y su relación con la salud subjetiva. [Clustering of behavioral risk factors and their association with subjective health]. *Gac Sanit*. 2005;19:370-378.
- 480 55. Winkleby MA, Gardner CD, Taylor CB. The influence of gender and socioeconomic factors on Hispanic/White differences in body mass index. *Prev Med*. 1996;25:203-211.

- 485 56. Wardle J, Robb KA, Johnson F, et al. Socioeconomic variation in attitudes to eating and weight in female adolescents. *Health Psychol.* 2004;23:275-282.
57. Lieux EM, Manning CK. Evening meals selected by college students: Impact of the foodservice system. *J Am Diet Assoc.* 1992;92:560-566.
58. Beerman KA, Jennings G, Crawford S. The effect of student residence on food choice. *J Am Coll Health.* 1990;28:250-220.
- 490 59. Cullen KW, Hartstein J, Reynolds KD, et al. Improving the school food environment: Results from a pilot study in middle schools. *J Am Diet Assoc.* 2007;107:484-489.
60. Cullen KW, Watson K, Zakeri I. Improvements in middle school student dietary intake after implementation of the Texas Public School Nutrition Policy. *Am J Public Health.* 2008;98:111-117.
- 495 61. Serlachius A, Hamer M, Wardle J. Stress and weight change in university students in the United Kingdom. *Physiol Behav.* 2007;92:548-553.
62. Farley T, Galves A, Dickinson LM, Diaz Perez MJ. Stress, coping, and health: A comparison of Mexican immigrants, Mexican Americans, and non-Hispanic whites. *J Immigr Health.* 2005;7:213-220.
- 500 63. Chiang L, Hunter CD, Yeh CJ. Coping attitudes, sources, and practices among Black and Latino college students. *Adolescence.* 2004;39:793-815.
64. Chiolero A, Wietlisbach V, Ruffieux C, Paccaud F, Cornuz J. Clustering of risk behaviors with cigarette consumption: A population-based survey. *Prev Med.* 2006;42:348-353.
- 505 65. Carroll SL, Lee RE, Kaur H, Harris KJ, Strother ML, Huang TT-K. Smoking, weight loss intention and obesity-promoting behaviors in college students. *J Am Coll Nutr.* 2006;25:348-353.

66. Taylor T, Murray AA, Hu D, Salgado F, Blow J, Cooper TV. *Light smoking patterns among students enrolled in a brief smoking cessation intervention*. Poster presented at the annual meeting of the Association of Behavioral and Cognitive Therapies 2007; Philadelphia, Pennsylvania.
- 510 67. Phillips LJ, Hammock RL., Blanton JM. Predictors of self-rated health status among Texas residents. *Prev Chronic Dis* [serial online]. 2005;2(4):1-10. Available at http://www.cdc.gov/pcd/issues/2005/oct/04_0147.htm. Accessed July 15, 2006.
68. Surís AM, Trapp MC, DiClemente CC, Cousins J. Application of the Transtheoretical Model of behavior change for obesity in Mexican American women. *Addict Behav*. 1998;23:655-668.
- 515 69. Neff JA, Hoppe SK. Acculturation and drinking patterns among U.S. Anglos, Blacks, and Mexican Americans. *Alcohol Alcohol*. 1992;27:293-308.
70. Carrera PM, Gao X, Tucker KL. A study of dietary patterns in the Mexican-American population and their association with obesity. *J Am Diet Assoc*. 2007;107:1735-1742.
71. Kuczmarski MF, Kuczmarski RJ, Najjar M. Effects of age on validity of self-reported height, 520 weight, and body mass index: Findings from the third National Health and Nutrition Examination Survey, 1988-1994. *J Am Diet Assoc*. 2001;101:28-34.
72. National Center for Education Statistics. Table 192: College enrollment and enrollment rates of recent high school completers, by race/ethnicity: 1960 through 2006. *Digest of Education Statistics: 2007*. Available at http://nces.ed.gov/programs/digest/d07/tables/dt07_192.asp. 525 Accessed May 20, 2009.

Table 1
Participant Characteristics ($n = 693$)

Categorical Demographics	%	Continuous Variables	<i>M</i>	<i>SD</i>
Gender		Age	19.6	3.33
Male	41	BMI	23.9	4.91
Female	59	Acculturation	2.97	0.74
BMI Categories		RISCI Coping	23.2	5.75
Underweight	7	RISCI Stress	17.9	3.61
Normal	61			
Overweight	21.5			
Obese	10.5			
Hispanic Identification		Risk Variables	%	
Mexican National	14	Exercise Risk		
Mexican American	72	No Regular Exercise	37	
Other Hispanic/Latin	14	Regular Exercise	63	
Household Income		5 a Day Risk		
>\$50k per year	28	No Regular Consumption	98	
\$30k to \$50k per year	27	Regular Consumption	2	
\$15k to \$30k per year	33	Dietary Fat Risk		
Unknown Income	12	No Regular Low Fat Diet	58	
		Low Fat Diet	42	
Reported Smoking Status		Total Risk		
At Least Daily Smoking	6	0 Risks	1	
At Least Weekly Smoking	5	1 Risk	31	
At Least Monthly Smoking	4	2 Risks	43	
Less Than Monthly Smoking	6	3 Risks	25	
Non-Smoker	79			

Table 2
Ordinal Logistic Regression of Obesity Relevant Health Behaviors

	OR	<i>p</i>	95% CI
Age (from 18)	1.05	0.06	1.00 - 1.11
Reported Household Income ¹			
\$30k to \$50k Income	1.48	0.07	0.97 - 2.27
\$15k to \$30k Income	1.93	<.01	1.28 - 2.92
Unknown Income	0.89	0.69	0.51 - 1.57
Female Gender (vs. Male)	0.92	0.65	0.66 - 1.29
Acculturation	0.96	0.70	0.76 - 1.20
BMI	0.99	0.57	0.96 - 1.02
Smoking Status ²	1.31	0.23	0.84 - 2.04
RISCI Stress	1.02	0.11	1.00 - 1.05
RISCI Coping	0.95	0.03	0.91 - 0.99

¹Income reference group is >\$50k per year Reported Household Income.

²Smoking Status reference group is Non-Smokers.

550


555

560


Figure Caption

Figure 1. Stage distributions for exercise, 5 A Day, and dietary fat recommendations


Exercise (n = 692)



5 A Day (n = 651)



Dietary Fat (n = 692)



Note. PC=Precontemplation, C=Contemplation, P=Preparation, A=Action, M=Maintenance