

Trinity University

From the Selected Works of William F. Trench

1989

Existence of solutions of functional-differential systems with prescribed asymptotic behavior

William F. Trench, *Trinity University*



Available at: https://works.bepress.com/william_trench/76/

EXISTENCE OF SOLUTIONS OF FUNCTIONAL DIFFERENTIAL SYSTEMS WITH PRESCRIBED ASYMPTOTIC BEHAVIOR

WILLIAM F. TRENCH

Proc. Fourth International Conference on Differential Equations, pp. 148-158 (Rousse, Bulgaria, 13-19 August, 1989); edited by P. Popivanov and S. Tersian.

We consider the $n \times n$ ($n \geq 1$) system of functional differential equations

$$(1) \quad X' = FX, \quad t > t_0.$$

For now we make no specific assumptions on the form of the functional F . For example, (1) may be a system of ordinary differential equations, an integro-differential system, a system with one or more deviating arguments, or a combination of these. To allow for the possibility that the values of $(FX)(t)$ for $t \geq t_0$ may depend on the values of $X(\tau)$ for some $\tau < t_0$ (as in the case of a delay equation, for example), we make the following definition.

Definition 1. *If $-\infty < t_0 < \infty$, then $\mathcal{C}_n(t_0)$ is the space of continuous n -vector functions $X = (x_1, \dots, x_n)$ on $(-\infty, \infty)$ which are constant on $(-\infty, t_0]$, with the topology induced by the following definition of convergence:*

$$X_j \rightarrow X \quad \text{as} \quad j \rightarrow \infty$$

if

$$\lim_{j \rightarrow \infty} \left[\sup_{-\infty < t \leq T} \|X_j(t) - X(t)\| \right] = 0$$

for every T in $(-\infty, \infty)$. (Here $\|\cdot\|$ is any convenient vector norm.)

Notice that $\mathcal{C}_n(t_1) \subset \mathcal{C}_n(t_0)$ if $t_0 \leq t_1$. We will say X is a *solution of (1) on $[t_0, \infty)$* if $X \in \mathcal{C}_n(a)$ for some $a \leq x_0$ and X satisfies (1) for $t \geq t_0$ (derivative from the right at t_0). We are interested in giving conditions on the functional F which imply that (1) has a solution \hat{X} such that $\lim_{t \rightarrow \infty} \hat{X}(t) = C$, where C is a given constant vector.

The Schauder–Tychonoff theorem has proved to be a powerful tool for establishing existence theorems of the kind that interest us here. More precisely, the following special case of this theorem, which is essentially the form stated by Coppel [1] has yielded many useful results.

Lemma 1. *Let \mathcal{S} be a closed convex subset of $C_n(t_0)$, and suppose that \mathcal{T} is a transformation of \mathcal{S} such that (a) $\mathcal{T}(\mathcal{S}) \subset \mathcal{S}$; (b) \mathcal{T} is continuous (i.e., if $\{X_j\} \subset \mathcal{S}$ and $X_j \rightarrow X$, then $\mathcal{T}X_j \rightarrow \mathcal{T}X$); and (c) the family of functions $\mathcal{T}(\mathcal{S})$ is uniformly bounded and equicontinuous on every compact subinterval of $[t_0, \infty)$. Then there is an \hat{X} in \mathcal{S} such that $\mathcal{T}\hat{X} = \hat{X}$.*

The following theorem illustrates one way in which Lemma 1 can be applied to our problem. We omit the proof, since this theorem follows from Theorem 3, below.

Theorem 1. *Suppose that there are constants a and M ($M > 0$) and a continuous function $w: [a, \infty) \rightarrow (0, \infty)$ such that $FX \in C_n[a, \infty)$ and $\|(FX)(t)\| \leq w(t)$ for $t \geq a$ whenever*

$$(2) \quad X \in \mathcal{C}_n(a) \text{ and } \|X(t)\| \leq M, \quad t \geq a.$$

Suppose further that

$$\int_a^\infty w(s) ds < \infty,$$

and that $\lim_{j \rightarrow \infty} (FX_j)(t) = (FX)(t)$ (pointwise) if each X_j satisfies (2) and $X_j \rightarrow X$. Let C be a given constant, with $\|C\| < M$. Then the system (1) has a solution \hat{X} on some interval $[t_0, \infty)$, such that $\lim_{t \rightarrow \infty} \hat{X}(t) = C$.

Although useful results can be obtained from this theorem, it is clear that the integrability condition on the functional F is very strong, since it implies that the integrals

$$(3) \quad \int_t^\infty \|(FX)(s)\| ds, \quad X \in \mathcal{S},$$

all converge, and even uniformly for all X in \mathcal{S} (i.e. $\int_t^\infty \|FX\| ds \leq \int_t^\infty w(s) ds$). It is quite possible to obtain useful results without requiring that the integrals (3) converge at all, so long as the integrals $\int^\infty (FX)(s) ds$ ($X \in \mathcal{S}$) converge in the ordinary (i.e., perhaps conditional) sense, and satisfy a uniform estimate of the form

$$(4) \quad \left\| \int_t^\infty (FX)(s) ds \right\| \leq \rho(t), \quad X \in \mathcal{S},$$

for some function ρ such that $\lim_{t \rightarrow \infty} \rho(t) = 0$. Moreover, it is important to exploit not just the assumption that the integrals in (4) converge, but also their rate of convergence. *Whenever possible, we should integrate before taking absolute values.* This point is often missed.

The author has pursued this theme in several papers (see, e.g., [2]–[5]). The results given here have been extended in [5].

Consider the following classical result for the linear system

$$(5) \quad \begin{bmatrix} x'_1 \\ x'_2 \\ \vdots \\ x'_n \end{bmatrix} = \begin{bmatrix} a_{11}(t) & a_{12}(t) & \dots & a_{1n}(t) \\ a_{21}(t) & a_{22}(t) & \dots & a_{2n}(t) \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1}(t) & a_{n2}(t) & \dots & a_{nn}(t) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix},$$

which follows easily from Theorem 1.

Theorem 2. *Suppose that $\{a_{ij}\}$ are continuous on $[a, \infty)$ and $\int_a^\infty |a_{ij}(t)| dt < \infty$ for $1 \leq i, j \leq n$. Let $C = (c_1, c_2, \dots, c_n)$ be a given constant vector. Then the system (5) has a solution \hat{X} such that $\lim_{t \rightarrow \infty} \hat{X} = C$.*

Example 1. Consider the system

$$(6) \quad \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \frac{\sin t}{t^\alpha} \begin{bmatrix} a_1 t^{-1} & b_1 \\ a_2 t^{-2} & b_2 t^{-1} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}, \quad t \geq a > 0,$$

where $b_1, b_2 \neq 0$ and $\alpha > 0$. Since

$$\int^\infty t^{-\alpha} |\sin t| dt \begin{cases} = \infty & \text{if } \alpha \leq 1, \\ < \infty & \text{if } \alpha > 1, \end{cases}$$

Theorem 2 does not apply to this system if $0 \leq \alpha \leq 1$; if $\alpha > 1$, then Theorem 2 implies that if c_1 and c_2 are given constants, then (6) has a solution $\hat{X} = (\hat{x}_1, \hat{x}_2)$ such that

$$\lim_{t \rightarrow \infty} x_i(t) = c_i, \quad i = 1, 2.$$

Theorem 2 provides no estimate of the *order* of convergence here, but it is straightforward to show that if $\alpha > 1$, then

$$x_1(t) = c_1 + O(t^{-\alpha+1}) \text{ and } x_2(t) = c_2 + O(t^{-\alpha}).$$

However, a more efficient use of integrability conditions for problems like this will show later that the true situation is as follows:

Suppose that $\alpha > 0$. Then:

(i) If c_1 is arbitrary and $c_2 \neq 0$, then (6) has a solution \hat{X} such that

$$x_1(t) = c_1 + O(t^{-\alpha}) \text{ and } x_2(t) = c_2 + O(t^{-\alpha-1}).$$

(ii) If c_1 is arbitrary and $c_2 = 0$, then (6) has a solution \hat{X} such that

$$x_1(t) = c_1 + O(t^{-\alpha-1}) \text{ and } x_2(t) = O(t^{-\alpha-2}).$$

The following theorem makes more efficient use of the Schauder–Tychonoff theorem (Lemma 1). Here it is convenient to rewrite (1) in component form as

$$x'_i = f_i X, \quad 1 \leq i \leq n, \quad t > t_0.$$

Theorem 3. *Let $C = (c_1, c_2, \dots, c_n)$ be a given constant vector. Let $\gamma_1, \dots, \gamma_n$ be continuous, positive and nonincreasing on $[t_0, \infty)$ and let M_1, \dots, M_n be positive constants. Let \mathcal{S} be the set of functions $X = (x_1, \dots, x_n)$ in $C_n(t_0)$ such that*

$$|x_i(t) - c_i| \leq M_i \gamma_i(t), \quad t \geq t_0, \quad 1 \leq i \leq n.$$

Suppose that F satisfies the following assumptions:

(i) $FX \in C_n[t_0, \infty)$ if $X \in \mathcal{S}$.

(ii) The family of functions $\mathcal{F} = \{FX \mid X \in \mathcal{S}\}$ is uniformly bounded on each subinterval of $[t_0, \infty)$.

(iii) If $\{X_j\} \subset \mathcal{S}$ and $X_j \rightarrow X$ (uniform convergence on every interval $(-\infty, T]$), then

$$\lim_{j \rightarrow \infty} (FX_j)(t) = (FX)(t) \text{ (pointwise), } t \geq t_0.$$

(iv) The integrals $\int^\infty (FX)(s) ds$ ($X \in \mathcal{S}$), converge, perhaps conditionally, and there are nonincreasing functions $\rho_1, \rho_2, \dots, \rho_n$ such that

$$(7) \quad 0 < \rho_i(t) \leq M_i \gamma_i(t), \quad 1 \leq i \leq n,$$

$$\lim_{t \rightarrow \infty} \rho_i(t) = 0, \quad 1 \leq i \leq n,$$

and, if $X \in \mathcal{S}$ and $t \geq t_0$,

$$(8) \quad \left| \int_t^\infty f_i X \, ds \right| \leq \rho_i(t), \quad 1 \leq i \leq n.$$

Then (1) has a solution \hat{X} on $[t_0, \infty)$ such that

$$|\hat{x}_i(t) - c_i| \leq \rho_i(t), \quad t \geq t_0, \quad 1 \leq i \leq n.$$

Proof. We define the transformation $Y = \mathcal{T}X$ in terms of components as

$$(9) \quad y_i(t) = \begin{cases} c_i - \int_t^\infty (f_i X)(s) \, ds, & t \geq t_0, \\ c_i - \int_{t_0}^\infty (f_i X)(s) \, ds, & t < t_0. \end{cases} \quad 1 \leq i \leq n.$$

Therefore, from (7), (8) and (9),

$$|y_i(t) - c_i| \leq \rho_i(t) \leq M_i \gamma_i(t);$$

hence, $\mathcal{T}(\mathcal{S}) \subset \mathcal{S}$, and $\mathcal{T}(\mathcal{S})$ is uniformly bounded on $[t_0, \infty)$, since \mathcal{S} is. Differentiating (9) shows that $y'_i(t) = (f_i X)(t)$ if $t \geq t_0$ and $y'_i(t) = 0$ if $t < t_0$; hence, the mean value theorem and assumption (iii) imply that the family $\mathcal{T}(\mathcal{S})$ is equicontinuous on every interval $(-\infty, T]$. The proof that \mathcal{T} is continuous is somewhat more delicate than in Theorem 1, since the integrals in question may converge conditionally. Suppose that $\{X_j\} \subset \mathcal{S}$ and $X_j \rightarrow X = (x_1, x_2, \dots, x_n)$ as $j \rightarrow \infty$. Denote $X_j = (x_{1j}, x_{2j}, \dots, x_{nj})$; then

$$y_{ij}(t) - y_i(t) = \begin{cases} \int_t^\infty (f_i X_j - f_i X) ds, & t \geq t_0, \\ \int_{t_0}^\infty (f_i X_j - f_i X) ds, & t < t_0. \end{cases}$$

Let

$$H_{ij} = \sup_{-\infty < t < \infty} |y_{ij}(t) - y_i(t)|, \quad 1 \leq i \leq n, \quad j = 1, 2, \dots.$$

Then, if $t_1 \geq t_0$,

$$\begin{aligned} H_{ij} &\leq \int_{t_0}^{t_1} |f_i X_j - f_i X| ds + \left| \int_{t_1}^\infty f_i X_j ds \right| + \left| \int_{t_1}^\infty f_i X ds \right| \\ &\leq \int_{t_0}^{t_1} |f_i X_j - f_i X| ds + 2\rho_i(t_1), \end{aligned}$$

from (8). Since the last integrand is uniformly bounded on $[t_0, t_1]$ for all j and $\rightarrow 0$ pointwise as $t \rightarrow \infty$, the last integral $\rightarrow 0$ as $t \rightarrow \infty$, by the bounded convergence theorem. Hence,

$$\overline{\lim}_{j \rightarrow \infty} H_{ij} \leq 2\rho_i(t_1)$$

for every t_1 . Since $\lim_{t_1 \rightarrow \infty} \rho_i(t_1) = 0$, this implies that $\lim_{j \rightarrow \infty} H_{ij} = 0$ for $1 \leq i \leq n$; that is, $y_{ij}(t) \rightarrow y_i(t)$ uniformly on $(-\infty, \infty)$ as $j \rightarrow \infty$. Now Lemma 1 implies the conclusion.

Theorem 4. *Let \mathcal{S} , $\gamma_1, \gamma_2, \dots, \gamma_n$, M_1, M_2, \dots, M_n and C be as in Theorem 3, and suppose that F satisfies assumptions (i) and (iii) on the set \mathcal{S} of functions $X = (x_1, \dots, x_n)$ in $C_n(t_0)$ such that*

$$|x_i(t) - c_i| \leq M_i \gamma_i(t), \quad t \geq t_0, \quad 1 \leq i \leq n.$$

Suppose further that $\int^\infty FC dt$ converges (perhaps conditionally) and that

$$\sup_{t \geq t_0} (\gamma_i(t))^{-1} \left| \int_t^\infty f_i C ds \right| = A_i < \infty, \quad 1 \leq i \leq n.$$

Suppose also that

$$|(f_i X)(t) - (f_i C)(t)| \leq M_i w_i(t), \quad 1 \leq i \leq n, \quad t \geq t_0,$$

for all X in \mathcal{S} , where

$$\sup_{t \geq t_0} (\gamma_i(t))^{-1} \int_t^\infty w_i ds = \theta_i < 1, \quad 1 \leq i \leq n.$$

Finally, let

$$M_i \geq \frac{A_i}{1 - \theta_i}.$$

Then the conclusion of Theorem 3 holds.

Proof. See [5].

We now apply Theorem 4 to the linear system (5).

Theorem 5. Suppose that $\{a_{ij}\}$ are continuous on $[a, \infty)$ and $\int_a^\infty a_{ij}(t) dt$ converges (perhaps conditionally) for $1 \leq i, j \leq n$. Let $C = (c_1, c_2, \dots, c_n)$ be a given constant vector, and suppose that $\gamma_1, \gamma_2, \dots, \gamma_n$ are nonincreasing positive functions on $[a, \infty)$ such that

$$\int_t^\infty f_i C ds = O(\gamma_i(t)), \quad 1 \leq i \leq n,$$

and define

$$w_i(t) = \sum_{j=1}^n |a_{ij}(t)| \gamma_j(t).$$

Suppose further that

$$\overline{\lim} (\gamma_i(t))^{-1} \int_t^\infty w_i(s) ds = \theta_i < 1, \quad 1 \leq i \leq n.$$

Then the system $X' = AX$ has a solution \hat{X} such that

$$\hat{x}_i(t) = c_i + O(\gamma_i(t)),$$

for $1 \leq i \leq n$; moreover, if $\theta_i = 0$, then this can be replaced by more precise estimate

$$\hat{x}_i(t) = c_i + \int_t^\infty f_i C ds + O\left(\int_t^\infty w_i ds\right).$$

Proof. See [5].

Example 1 (Continuation). As mentioned above, a linear system $X' = A(t)X$ (with A continuous on $[0, \infty)$) has a solution \hat{X} satisfying an arbitrary final condition $\lim_{t \rightarrow \infty} \hat{x}(t) = C$ if $\int_0^\infty \|A(t)\| dt < \infty$. This result can be obtained from Theorem 5 by simply taking $\gamma_i = 1$, $1 \leq i \leq n$. The system (6) with $b_1 \neq 0$ does not satisfy this integrability condition if $\alpha \leq 1$; moreover, even if $\alpha > 1$, the standard theorem merely implies that if c_1 and c_2 are given constants, then (6) has a solution (\hat{x}_1, \hat{x}_2) such that $\lim_{t \rightarrow \infty} \hat{x}_i(t) = c_i$ ($i = 1, 2$). However, Theorem 5 implies that if $\alpha > 0$ and (c_1, c_2) is arbitrary, then (6) has a solution (\hat{x}_1, \hat{x}_2) such that

$$\hat{x}_1(t) = c_1(1 - a_1 S_{\alpha+1}(t)) - b_1 c_2 S_\alpha(t) + 0(t^{-2\alpha})$$

and

$$\hat{x}_2(t) = -a_2 c_1 S_{\alpha+2}(t) + c_2(1 - b_2 S_{\alpha+1}(t)) + 0(t^{-2\alpha-1}),$$

where

$$S_\beta(t) = \int_t^\infty s^{-\beta} \sin s ds = 0(t^{-\beta}), \quad \beta > 0.$$

This conclusion is obtained by letting $\gamma_1(t) = t^{-\alpha}$ and $\gamma_2(t) = t^{-\alpha-1}$. A sharper result is available if $c_2 = 0$; i.e., for every constant c_1 , (6) has a solution (\hat{x}_1, \hat{x}_2) such that

$$\hat{x}_1(t) = c_1(1 - a_1 S_{\alpha+1}(t)) + 0(t^{-2\alpha-1}),$$

and

$$\hat{x}_2(t) = -a_2 c_1 S_{\alpha+2}(t) + 0(t^{-2\alpha-2}).$$

This is obtained by letting $\gamma_1(t) = t^{-\alpha-1}$ and $\gamma_2(t) = t^{-\alpha-2}$.

We will now obtain a global result for the nonlinear integral equation

$$(10) \quad x'(t) = g(t)(x(t))^\alpha + \int_0^t P(t, \tau)(x(\tau))^\beta d\tau, \quad t > 0,$$

where $g \in C[0, \infty)$ and P is continuous on $[0, \infty) \times [0, \infty)$.

Theorem 6. *Suppose that*

$$(11) \quad \int_t^\infty g(s) ds = O(\gamma(t)),$$

$$(12) \quad \int_t^\infty |g(s)|\gamma(s) ds = O(\gamma(t)),$$

$$(13) \quad \int_t^\infty \int_0^s P(s, \tau) d\tau ds = O(\gamma(t)),$$

and

$$(14) \quad \int_t^\infty \int_0^s |P(s, \tau)|\gamma(\tau) d\tau ds = O(\gamma(t)),$$

where γ is positive and nonincreasing on $[0, \infty)$, and $\lim_{t \rightarrow \infty} \gamma(t) = 0$. Suppose also that $0 < \theta < 1$. Then there is a constant $c_0 > 0$ such that (10) has a solution \hat{x} on $[0, \infty)$ which satisfies the following conditions:

$$|\hat{x}(t) - c| \leq \theta c \quad (t \geq 0), \quad \hat{x}(t) = c + O(\gamma(t)).$$

provided that either

(a) $\alpha, \beta > 1$ and $0 < c < c_0$; or

(b) $\alpha, \beta < 1$ and $c > c_0$.

(Notice that (11) and (12) do not imply that $\int_0^\infty |g(s)| ds < \infty$, nor do (13) and (14) imply that $\int_0^\infty \int_0^s |P(s, \tau)| d\tau ds < \infty$.)

Proof. For convenience, normalize γ so that $\gamma(0) = 1$. Here

$$(Fx)(t) = g(t)(x(t))^\alpha + \int_0^t P(t, \tau)(x(\tau))^\beta d\tau.$$

and

$$(Fc)(t) = c^\alpha \int_t^\infty g(s) ds + c^\beta \int_0^t P(t, \tau) d\tau.$$

If $c > 0$, let

$$\mathcal{S} = \{x \in C[0, \infty) \mid |x(t) - c| \leq \theta c \gamma(t), t \geq 0\}.$$

Then, if $x \in \mathcal{S}$, $|x(t) - c| \leq \theta c$ ($t \geq 0$), since γ is nonincreasing. Obviously, F satisfies assumptions (i), (ii), and (iii) of Theorem 3. Now,

$$\begin{aligned} \int_t^\infty Fx ds &= \int_t^\infty Fc ds + \int_t^\infty (Fx - Fc) ds \\ &= c^\alpha \int_t^\infty g(s) ds + c^\beta \int_t^\infty \int_0^s P(s, \tau) d\tau ds \\ &\quad + \int_t^\infty g(s)[(x(s))^\alpha - c^\alpha] ds \int_t^\infty \int_0^s P(s, \tau)[(x(s))^\beta - c^\beta] ds. \end{aligned}$$

By the mean value theorem,

$$|x^\alpha - c^\alpha| \leq K(\alpha) =_{df} |\alpha|[(1 \pm \theta)c]^{\alpha-1}|x - c|$$

if $|x - c| \leq \theta c$ (with “+” if $\alpha > 1$, “−” if $\alpha < 1$). Since $|x(t) - c| \leq \theta c \gamma(t)$ if $x \in \mathcal{S}$, this means that

$$\left| \int_t^\infty Fx ds \right| \leq c^\alpha \left| \int_t^\infty g(s) ds \right| + c^\beta \left| \int_t^\infty \int_0^s P(s, \tau) d\tau ds \right|$$

$$Kc^\alpha \int_t^\infty |g(s)|\gamma(s) ds + Kc^\beta \int_t^\infty \int_0^s |P(s, \tau)|\gamma(\tau) d\tau ds,$$

where K is a constant which does not depend on c . Since all four integrals on the right are $O(\gamma(t))$, this means that

$$(\gamma(t))^{-1} \left| \int_t^\infty Fx \, ds \right| \leq Ac^\alpha + Bc^\beta, \quad x \in \mathcal{S}, \quad t \geq 0,$$

where A and B are constants which do not depend on c . Since our requirement is that

$$(\gamma(t))^{-1} \left| \int_t^\infty Fx \, ds \right| \leq \theta c, \quad x \in \mathcal{S}, \quad t \geq 0,$$

we have only to choose c so that

$$Ac^{\alpha-1} + Bc^{\beta-1} \leq \theta.$$

This is true for c sufficiently large if $\alpha, \beta < 1$, or for c sufficiently small if $\alpha, \beta > 1$.

1. W.B. Coppel, Stability and asymptotic behavior of differential equations, D.C. Heath, Boston.
2. W.F. Trench, Systems of differential equations subject to mild integral smallness conditions, Proc. Amer. Math. Soc. 87 (1983), 263–270.
3. W.F. Trench, Asymptotics of differential systems with deviating arguments, Proc. Amer. Math. Soc. **92** (1984), 219–224.
4. W.F. Trench, Extensions of a theorem of Wintner on systems with asymptotically constant solutions, Trans. Amer. Math. Soc. **293** (1986), 477–483.
5. W.F. Trench, Efficient application of the Schauder–Tychonoff theorem to systems of functional differential equations. J. Math. Anal. Appl. (to appear).

TRINITY UNIVERSITY
SAN ANTONIO, TEXAS 78212
USA